

References

A

- Abramoff, M.D., Magelhaes, P.J. and Ram, S.J. 2004. Image processing with ImageJ. *Biophotonics Intl.* 11: 36-42.
- Adams, G.M., Falke, S., Goldberg, A.L., Slaughter, C.A., DeMartino, G.N. and Gogol, P.E. 1997. Structural and functional effects of PA700 and modulator protein on proteasomes. *J. Mol. Biol.* 273: 646-657.
- Adams, D.J. 2004. Fungal cell wall chitinases and glucanases. *Microbiology.* 150: 2029-2035.
- Adams, J. 2003a. Potential for proteasome inhibition in the treatment of cancer. *Drug Discovery Today.* 8: 307-315.
- Adams, J. 2003b. The proteasome: structure, function and role in the cell. *Cancer Treatment Rev.* 29: 3-9.
- Amerik, A.Y., Nowak, J., Swaminathan, S. and Hochstrasser, M. 2000. The Doa4 deubiquitinating enzyme is functionally linked to the vacuolar protein-sorting and endocytic pathways. *Mol Biol Cell.* 11: 3365-3380.
- Amerik, A., Sindhi, N. and Hochstrasser, M. 2006. A conserved late endosome-targeting signal required for Doa4 deubiquitylating enzyme function. *J. Cell Biol.* 175: 825-835.
- Amir, S., Alin, F., Neta, S. and Ido, H. 2009. Fungal apoptosis: function, genes and gene function. *FEMS Microbiology Reviews.* 33: 833-854.
- Archer, C.T., Burdine, L., Liu, B., Ferdous, A., Johnston, S.A. and Kodadek, T. 2008. Physical and functional interactions of monoubiquitylated transactivators with the proteasome. *J. Biol. Chem.* 283: 21789-21798.
- Archer, D.B. and Peberdy, J.F. 1997. The molecular biology of secreted enzyme production by fungi. *Crit. Rev. Biotechnol.* 17: 273-306.
- Aro, N., Saloheimo, A., Ilmén, M. and Penttilä, M. 2001. ACEII, a novel transcriptional activator involved in regulation of cellulase and xylanase genes of *Trichoderma reesei*. *J. Biol. Chem.* 276: 24309-24314.
- Aro, N., Ilmén, M., Saloheimo, A. and Penttilä, M. 2003. ACEI of *Trichoderma reesei* is a repressor of cellulase and xylanase expression. *Appl. Environ. Microbiol.* 69: 56-65.
- Arvas, M., Pakula, T., Lanthaler, K., Saloheimo, M., Valkonen, M., Suortti, T., Robson, G. and Penttilä, M. 2006. Common features and interesting differences in

transcriptional responses to secretion stress in the fungi *Trichoderma reesei* and *Saccharomyces cerevisiae*. BMC Genomics. 7: 32-50.

Asano, K., Kinzy, T. G., Merrick, W.C. and Hershey, J.W. 1997. Conservation and diversity of eukaryotic translation initiation factor eIF3. J. Biol. Chem. 272: 1101-1109.

Askolin, S., Penttilä, M., Wörsten, H.A.B. and Nakari-Setälä, T. 2005. The *Trichoderma reesei* hydrophobin genes *hfb1* and *hfb2* have diverse functions in fungal development. FEMS 253: 281-288.

Altschuler, G.M., Dekker, C., McCormack, E.A., Morris, E.P., Klug, D.R. and Willison, K.R. 2009. A single amino acid residue is responsible for species-specific incompatibility between CCT and alpha-actin. FEBS Lett. 583: 782-726.

Attaix D., Combaret L., Pouch M.-N. and Taillandier D. 2002. Cellular control of ubiquitin-proteasome-dependent proteolysis. J. Anim. Sci. 80: E56-E63.

B

Babbitt, S.E., Kiss, A., Deffenbaugh, A.E., Chang, Y., Bailly, E., Erdjument-Bromage, H., Tempst, P., Buranda, T., Sklar, L. A., Baumler, J., Gogol, E. and Skowrya, D. 2005. ATP hydrolysis-dependent disassembly of the 26S proteasome is part of the catalytic cycle. Cell 121: 552-565.

Bailly, E. and Reed, S.I. 1999. Functional characterization of Rpn3 uncovers a distinct 19S proteasomal subunit requirement for ubiquitin-dependent proteolysis of cell cycle regulatory proteins in budding yeast. Mol. Cell Biol. 19: 6872-6890.

Baker, S.P. and Grant, P.A. 2007. The SAGA continues: expanding the cellular role of a transcriptional co-activator complex. Oncogene 26: 5329-5340.

Baker, S. and Grant, P.A. 2005. The Proteasome: Not Just Degrading Anymore. Cell. 4: 361-363.

Bardy, S. L., Ng, S.M.Y., Carnegie, D.S., and Jarrell, K.F. 2005. Site-directed mutagenesis analysis of amino acids critical for activity of the type I signal peptidase of the archaeon *Methanococcus voltae*. J. Bacteriol. 187: 1188-1191.

Bateman, D.A. 2007. Unfolded protein response (UPR): Cellular Control for our errors in life. Hypothesis 33-37.

Bailey, M. J., and Nevalainen, K.M.H. 1981. Induction, isolation and testing of stable *Trichoderma reesei* mutants with improved production of solubilizing cellulase. Enz. Microb. Technol. 3:153-157.

Baumeister, W. Walz, J., Zuhl, F. and Seemuller, E. 1998. The proteasome: paradigm of a self-compartmentalizing protease. Cell 92: 367-380.

- Bays, N.W. and Hampton, R.Y. 2002. Cdc48-Ufd-Npl4: Stuck in the middle with Ub. *Curr. Biol.* 12: R366-R371.
- Beaudenon, S.L., Huacani, M.R., Wang, G., McDonnell, D.P. and Huibregtse, J.M. 1999. Rsp5 ubiquitin-protein ligase mediates DNA damage induced degradation-induced degradation of the large subunit of RNA polymerase II in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 19: 6972-6979.
- Benaroudj, N., Tarsa, E., Cascio, P. and Goldberg, A.L. 2001. The unfolding of substrates and ubiquitin-independent protein degradation by proteasomes. *Biochem.* 83: 1-8.
- Bergquist, P.L., Teodoro V.S.J., Gibbs M.D., Curach N.C. and Nevalainen K.M.H. 2004. Recombinant enzymes from thermophilic micro-organisms produced in fungal hosts. *Biochem. Soc. Trans.* 32: 293-297.
- Bertolotti, A., Zhang, Y., Hendershot, L.M., Harding, H.P. and Ron, D. 2000. Dynamic interaction of BiP and ER stress transducers in the unfolded-protein response. *Nat. Cell Biol.* 2: 326-332.
- de Biesebeke, R., van Biezen, N., deVos, W.M., van den Hondel, C.A. and Punt, P. 2005. Different control mechanisms regulate glucoamylase and protease gene transcription in *Aspergillus oryzae* in solid-state and submerged fermentation. *Appl. Microbiol. Biotechnol.* 67: 75-82.
- Bonaccorsi, E. D., Ferreira, A. J. S., Chambergo, F. S., Ramos, S. P., Mantovani, M. C., Farah, J. P. S., Sorio, C. S., Gombert, A. K., Tonso, A. and El-Dorri, H. 2006. Transcriptional response of the obligatory aerobic *Trichoderma reesei* to hypoxia and transient anoxia: implications for energy production and survival in the absence of oxygen. *Biochemistry.* 45: 3912-3924.
- Bose, S. and Freedman, R. B. 1994. Peptidyl prolyl cis-trans-isomerase activity associated with the lumen of the endoplasmic reticulum. *Biochem. J.* 3: 865-870.
- Bose, S., Stratford, F.L., Broadfoot, K.I., Mason, G.G. and Rivett, A.J. 2004. Phosphorylation of 20S proteasome alpha subunit C8 (alpha7) stabilizes the 26S proteasome and plays a role in the regulation of proteasome complexes by gamma-interferon. *Biochem. J.* 378:177-184.
- Botchler, M., Ditzel, L., Groll, M., Hartmann, C. and Huber, R. 1999. The proteasome. *Annu. Rev. Biophys. Biomol. Struct.* 28: 295-317.
- Bourne, Y. and Henrissat, B. 2001. Glycoside hydrolases and glycosyltransferases: families and functional modules. *Curr. Opin. Struct. Biol.* 11: 593-600.

- Bradford, M. M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72: 248-254.
- Brandina, I., Smirnov, A., Kolesnikova, O., Entelis, N., Krasheninnikov, I.A., Martin, R.P. and Tarassov, I. 2007. tRNA import into yeast mitochondria is regulated by ubiquitin-proteasome system. *FEBS Lett.* 581: 4248-4254.
- Brazma, A., Hingamp, P., Quackenbush, J., Sherlock, G., Spellman, P., Stoecker, C., Aach, J., Ansorge, W., Ball, C.A., Causton, H.C., Gaasterland, T., Glenisson, P., Holstege, F.C.P., Kim, I.F., Markowitz, V., Matese, J.C., Parkinson, H., Robinson, A., Sarkans, U., Schulze-Kremer, S., Stewart, J., Taylor, R., Vilo, J. and Vingron, M. 2001. Minimum information about a microarray experiment (MIAME)- toward standards for microarray data. *Nat. Gen.* 29: 365-371.
- Breakspear, A. and Momany, M. 2007. The first fifty microarray studies in filamentous fungi. *Microbiology.* 153: 7-15.
- Brodsky, J.L., Hamamoto, S., Feldheim, D., Schekman, R. 1993. Reconstitution of protein translocation from solubilized yeast membranes reveals topologically distinct roles for BiP and cytosolic Hsc70. *J. Cell Biol.* 120: 95-102.
- Brodsky, J.L. and McCracken, A.A. 1999. ER protein quality control and proteasome-mediated protein degradation. *Semin. Cell Dev. Biol.* 10: 507-513.
- Brodsky, J.L., Werner, E.D., Dubas, M.E., Goeckeler, J.L., Kruse, K.B. and McCracken, A.A. 1999. The requirement for molecular chaperones during endoplasmic reticulum-associated protein degradation demonstrates that protein export and import are mechanistically distinct. *J. Biol. Chem.* 274: 3453-3460.
- Braun, B., Glickman, M., Kraft, R., Dahlmann, B., Kloetzel, P.-M., Finley, D. and Schmidt, M. 1999. The base of the proteasome regulatory particle exhibits chaperone-like activity. *Nat. Cell Biol.* 1: 221-226.
- Bukau, B. and Horwich, A.L. 1998. The Hsp 70 and Hsp 60 chaperone machines. *Cell* 92: 351-366.
- Bush, K.T., Goldberg, A.L., and Nigam, S.K. 1997. Proteasome inhibition leads to a heat-shock response, induction of endoplasmic reticulum chaperones and thermotolerance. *J. Biol. Chem.* 272: 9086-9092.
- Bustin, S.A. and Mueller, R. 2005. Real-time reverse transcription PCR (qRT-PCR) and its potential use in clinical diagnosis. *Clin. Sci.* 109: 365-379.

C

- Cabral, C.M., Liu Y., Moremen, K. W. and Sifers R.N. 2002. Organizational diversity among distinct glycoprotein endoplasmic reticulum-associated degradation programs. *Mol. Biol. Cell.* 13: 2639-2650.
- Cabral, C.M., Liu, Y. and Sifers, R.N. 2001. Dissecting glycoprotein quality control in the secretory pathway. *Trends Biochem. Sci.* 26: 619-624.
- Caldwell, S.R., Hill, K.J. and Cooper, A.A. 2001. Degradation of endoplasmic reticulum (ER) quality control substrates requires transport between the ER and Golgi. *J. Biol. Chem.* 276: 23296-23303.
- Cascio, P., Call, M., Petre, B.M., Walz, T. and Goldberg, A. 2002. Properties of the hybrid form of the 26S proteasome containing both 19S and PA28 complexes. *EMBO J.* 21: 2636-2645.
- Chalfie, M., Tu, G., Euskirchen, G., Ward, W.W. and Prasher, D.C. 1994. Green fluorescent protein as a marker for gene expression. *Science* 263: 802-806.
- Chen, H., Hayn, M. and Esterbauer, H. 1993. Three forms of cellobiohydrolase I from *Trichoderma reesei*. *Biochem. Mol. Biol. Int.* 30: 901-910.
- Cherry, J.R. and Fidantsef, A.L. 2003. Directed evolution of industrial enzymes: an update. *Curr. Opin. Biotechnol.* 14: 438-443.
- Chevet E., Cameron P.H., Pelletier M.F., Thomas D.Y. and Bergeron J.J. 2001. The endoplasmic reticulum: integration of protein folding, quality control, signaling and degradation. *Curr. Opin. Struct Biol.* 11: 112-120.
- Chung, K.K.K., Dawson, V.L. and Dawson, T.M. 2001. Role of the ubiquitin- proteasomal pathway in Parkinson's disease and other neurodegenerative disorders. *Trends Neurosci.* 24: 7-14.
- Chuaqui, R.F., Bonner, R.F., Best, C.J.M., John W. Gillespie, Flaig, M.J., Hewitt, S.H., Phillips, J.L., Krizman, D.B., Tangrea, M.A, Ahram, M., Linehan, W.M., Knezevic, V. and Emmert-Buck, M.R. 2002. Post-analysis follow-up and validation of microarray experiments. *Nature Genetics.* 32: 509-514.
- Ciccarelli, F.D., Izaurralde, E. and Bork, P. 2003. The PAM domain, a multi-protein complex-associated module with an all-alpha-helix fold. *BMC Bioinformatics.* 4: 64-69.
- Ciechanover, A. 2005. Proteolysis: from the lysosome to ubiquitin and the proteasome. *Nature Rev.* 6: 79-86.

- Ciechanover, A. 2006. The ubiquitin proteolytic system: from a vague idea, through basic mechanisms, and onto human diseases and drug targeting. *Neurology*. 66 : S7-S19.
- Conesa, A., Punt, P. J. and van den Hondel C. A. M. J. J. 2001. The secretion pathway in filamentous fungi: a biotechnological view. *Fungal Gen. Biol.* 33: 155-171.
- Conesa, A., Jeenes, D., Archer, D.B., van den Hondel, C.A.M.J.J. and Punt, P. J. 2002. Calnexin overexpression increases manganese peroxidase production in *Aspergillus niger*. *Appl. Environ. Microbiol.* 68: 846-851.
- Collén, A., Saloheimo, M., Bailey, M., Penttilä, M. and Pakula, T. M. 2005. Protein production and induction of the unfolded protein response in *Trichoderma reesei* strain Rut-C30 and its transformant expressing endoglucanase I with a hydrophobic tag. *Biotechnology and Bioengineering*. 89: 335-344.
- Coux, O., Tanaka, K. and Goldberg, A.L. 1996. Structure and functions of the 20S and 26S proteasomes. *Annu. Rev. Biochem.* 65: 801-847.

D

- DenBoer, L.M., Hardy-Smith, P.W., Hogan, M.R., Cockram, G.P., Audas, T.E. and Lu, R. 2005. Luman is capable of binding and activating transcription from unfolded protein response element. *Biochem. Biophys. Res. Com. (BBRC)*. 331: 113-119.
- DeRisi, J.L., Iyer, V.R. and Brown, P.O. 1997. Exploring metabolic and genetical control of gene expression on a genomic scale. *Science* 278: 680-686.
- Dickson, K.M., Bergeron, J.J.M., Shames, I., Colby, J., Nguyen, D.T., Chevet, E., Thomas, D.Y. and Snipes G.J. 2002. Association of calnexin with mutant peripheral myelin protein-22 *ex vivo*: A basis for □gain-of-function□ER diseases. *Cell Biol.* 99: 9852-9857.
- Divne, C., Ståhlberg, J., Reinikainen, T., Ruohonen, L., Pettersson, G., Knowles, J.K.C., Teeri, T.T. and Jones, A. 1994. The three-dimensional structure of the catalytic core of cellobiohydrolase I from *Trichoderma reesei*. *Science*. 265: 524-528.
- Divne, C., Ståhlberg, J., Teeri, T.T. and Jones, T.A. 1998. High-resolution crystal structures reveal how a cellulose chain is bound in the 50Å long tunnel of cellobiohydrolase I from *Trichoderma reesei*. *J. Mol. Biol.* 275: 309-325.
- Drexler, H.C.A. 2009. Synergistic apoptosis induction in leukemic cells by the phosphatase inhibitor salubrinal and proteasome inhibitors. *PLoS ONE* 4(1): e4161. doi:10.1371/journal.pone.0004161. Published online on Jan 8 2009.
- Dobson, C.M. 1999. Protein misfolding, evolution and disease. *TIBS*. 24: 329-332.
- Dobson, C.M. 2003. Protein folding and misfolding. *Nature*. 426: 884-890.

Derkx, P.M. and Madrid, S. M. 2001. The foldase CYPB is a component of the secretory pathway of *Aspergillus niger* and contains the endoplasmic reticulum retention signal HEEL. Mol. Genet. Genomics. 266: 537-45.

E

Ellgaard, L., Molinari, M. and Helenius, A. 1999. Setting the standards: quality control in the secretory pathway. Science. 286: 1882-1888.

Elsässer, S., Chi, Y., Yang, P. and Campbell, J.L. 1999. Phosphorylation controls timing of Cdc6p destruction: a biochemical analysis. Mol. Biol. Cell 10: 3263-3277.

Elsässer, S., Gali, R.R., Schwickart, M., Larsen, C.N., Leggett, D.S., Müller, B., Feng, M.T., Tübing, F., Dittmar, G.A. and Finley, D. 2002. Proteasome subunit Rpn1 binds ubiquitin-like protein domains. Nat. Cell Biol. 4: 725-730.

Elsässer, S., Chandler-Militello, D., Müller, B., Hanna, J. and Finley, D. 2004. Rad23 and Rpn10 serve alternative ubiquitin receptors for the proteasome. J. Biol. Chem. 279: 26817-26822.

Eneyskaya, E.V., Kulminskaya, A.A., Savel', A. N., Savel'eva, N. V., Shabalin, K. A. and Neustroev, K. N. 1999. Acid protease from *Trichoderma reesei*: limited proteolysis of fungal carbohydrases. Appl. Microbiol. Biotechnol. 52: 226-231.

Eveleigh, D.E. and Montenecourt, B.S. 1979. Increasing yields of extracellular enzymes. Adv. Appl. Microbiol. 25: 57-74.

Eytan, E., Armon, T., Heller, H., Beck, S. and Hershko, A. 1993. Ubiquitin C-terminal hydrolase activity associated with the 26 S protease complex. J. Biol. Chem. 268: 4668-4674.

Ezhkova, E., and Tansey, W.P. 2004. Proteasomal ATPases link ubiquitylation of histone H2B to methylation of histone H3. Mol. Cell 13: 435-442.

F

Farrell, P.A., Hernandez, J.M., Fedele, M.J., Vary, T.C., Kimball, S.R. and Jefferson, L.S. 2000. Eukaryotic initiation factors and protein synthesis after resistance exercise in rats. J. Appl. Physiol. 88:1036-1042.

Fatica, A., Tollervey, D. and Dlakic', M. 2004. PIN domain of Nob1p is required for D-site cleavage in 20S pre-rRNA. RNA 10: 1698-1701.

Feldheim, D., Yoshimura, K., Admon, A. and Schekman R. 1993. Structural and functional characterization of Sec66p, a new subunit of the polypeptide translocation apparatus in the yeast endoplasmic reticulum. Mol. Biol. Cell. 4: 931-939.

- Feng, G., Mellor, R.H., Bernstein, M., Keller-Peck, C., Ngyen, Q.T., Wallace, M., Nerbonne, J.M., Lichtman, J.W. and Sanes, J.R. 2000. Imaging neuronal subsets in transgenic mice expressing multiple spectral variants of GFP. *Neuron*. 28: 41-51.
- Fenteany, G., Staendaert R. F., Lane W.S., Choi S., Corey, E.J. and Schreiber, S.L. 1995. Inhibition of proteasome activities and subunit-specific amino-terminal threonine modification by lactacystin. *Science*. 268: 726-731.
- Ferdous, A., Gonzalez, F., Sun, L., Kodadek, T. and Johnston, S.A. 2001. The 19S regulatory particle of the proteasome is required for efficient transcription elongation by RNA polymerase II. *Mol. Cell* 7: 981-991.
- Fewell, S.W., Travers, K.J., Weissman, J.S. and Brodsky, J.L. 2001. The action of molecular chaperones in the early secretory pathway. *Annu Rev Genet*. 35: 149-191.
- Fischer, R., Schumann, D., Zimmermann, S., Drossard, J., Sack, M. and Schillberg, S. 2001. Expression and characterization of bispecific single-chain Fv fragments produced in transgenic plants. *Eur. J. Biochem*. 262: 810-816.
- Flinn, J.P., Pallaghy, P.K., Lew, M.J., Murphy, R., Angus, J.A. and Norton, R.S. 1999. Role of disulfide bridges in the folding, structure and biological activity of ω -conotoxin GVIA. *Prot. Struct. Mol. Enzymol*. 1434: 177-190.
- Frand, A.R. and Kaiser, C.A. 1999. Ero1p oxidizes protein disulfide isomerase in a pathway for disulfide bond formation in the endoplasmic reticulum. *Mol. Cell*. 4: 469-477.
- Friedländer, R., Jarosch, E., Urban, J., Volkwein, C. and Sommer, T. 2000. A regulatory link between ER-associated protein degradation and the unfolded-protein response. *Nat. Cell Biol*. 2: 379-384.
- Frydman, J., and Hartl, F-U. 1994. The stress response, functions of the proteins, and perspectives. pp. 251-283. In *The biology of heat shock proteins and molecular chaperones*. Eds. Morimoto, R. I., Tissieres, A., and Georgopoulos, C. Cold Spring Harbor Laboratory. Cold Spring Harbor, NY.
- Foreman, P.K., Brown, D., Dankmayer, L., Dean, R., Diener, S., Dunn-Coleman, N.S., Goedegebuur, F., Houfek, T.D., England, G.J., Kelley, A.S., Meerman, H.J., Mitchell, T. Mitchinson, C., Olivares, H.A., Teunissen, P.J.M., Yao, J. and Ward, M. 2003. Transcriptional regulation of biomass-degrading enzymes in the filamentous fungus *Trichoderma reesei*. *J. Biol. Chem*. 278: 31988-31997.
- Froment, C., Uttenweiler-Joseph, S., Bousquet-Dubouch, M.P., Matondo, M., Borges, J.P., Esmenjaud, C., Lacroix, C., Monsarrat, B. and Burlet-Schiltz, O. 2005. A quantitative proteomic approach using two-dimensional gel electrophoresis and

isotope-coded affinity tag labeling for studying human 20S proteasome heterogeneity. *Proteomics* 5: 2351-2363.

Fu, L. and Sztul, E. 2003. Traffic-independent function of the Sar1p/COPII machinery in proteasomal sorting of the cystic fibrosis transmembrane conductance regulator. *Cell Biol.* 160: 157-163.

Fuertes, G., Villarroya, A. and Knecht, E. 2005. Role of proteasomes in the degradation of short-lived proteins in human fibroblasts under various growth conditions. *Int. J. Biochem. Cell Biol.* 35: 651-664.

Fujii, T., Doi, Y., Ueno, H. and Hayashi, R. 2000. Effect of mutagenic replacement of the carboxyl terminal amino acid, Val24, on the properties and regeneration of bovine pancreatic ribonuclease A. *J. Biochem.* 127: 877-881.

Fujinami, K., Tanahashi, N., Tanaka, K., Ichihara, A., Cejka, Z., Baumeister, W., Miyawaki, M., Sato, T. and Nakagawa, H. 1994. Purification and characterisation of the 26S proteasome from spinach leaves. *J. Biol. Chem.* 269: 25905-25910.

Fujita, E., Kouroku, Y., Isoai, A., Kumagai, H., Misutani, A., Matsuda, C., Hayashi, Y.K. and Momoi, T. 2007. Two endoplasmic reticulum-associated degradation (ERAD) systems for the novel variant of the mutant dysferlin: ubiquitin/proteasome ERAD (I) and autophagy/lysosome ERAD (II). *H. Mol. Gen.* 16: 618-629.

G

Gardner, R.G., Swarbrick, G.M., Bays, N.W., Cronin, S.R., Wilhovsky, S., Seelig, L., Kim, C. and Hampton, R.Y. 2000. Endoplasmic reticulum degradation requires lumen to cytosol signalling transmembrane control of Hrd1p by Hrd3p. *J. Cell Biol.* 151: 69-82.

Gasser, G., Maurer, M., Rautio, J., Sauer, M., Bhattacharyya, A., Saloheimo, M., Penttilä, M. and Mattanovich, D. 2007. Monitoring of transcriptional regulation in *Pichia pastoris* under protein production conditions. *BMC Genomics.* 8: 179-197.

Gauss, R., Sommer, T. and Jarosch, E. 2006. The Hrd1p ligase complex forms a linchpin between ER-lumenal substrate selection and CDC48p recruitment. *EMBO J.* 25: 1827-1837.

van Gemeren, I.A., Punt, P.J., Drint-Kuyvenhoven, A., Broekhuijsen, M.P., van 't Hoog, A., Beijersbergen, A., Verrips, C.T. and van den Hondel, C.A.M.J.J. 1997. The ER chaperone encoding *bipA* gene of black *Aspergilli* is induced by heat shock and unfolded proteins. *Gene.* 198: 43-52.

Gerards, W.L.H., de Jong, W.W., Boelens, W. and Bloemendal, H. 1998. Structure and assembly of the 20S proteasome. *CMLS. Cell. Mol. Life Sci.* 54: 253-262.

- Gething, M.J. 1999. Role and regulation of the ER chaperone BiP. *Cell. Dev. Biol.* 10: 465-472.
- Geysens, S., Whyteside, G. And Archer, D.B. 2009. Genomics of protein folding in the endoplasmic reticulum, secretion stress and glycosylation in the aspergilli. *Fungal Genetics and Biology* 46: 121-140.
- Ghislain, M., Dohmen, R.J., Levy, F., and Varshavsky, A. 1996. Cdc48p interacts with Ufd3p, a WD repeat protein required for ubiquitin-mediated proteolysis in *Saccharomyces cerevisiae*. *EMBO J.* 15: 4884-4899.
- Gillette, T.G., Gonzalez, F., Delahodde, A., Johnston, S.A. and Kodadek, T. 2004. Physical and functional association of RNA polymerase II and the proteasome. *Proc. Natl. Acad. Sci. USA* 101: 5904-5909.
- Glick, B.S. 1995. Can Hsp70 proteins act as force-generating motors? *Cell.* 80: 11-14.
- Glickman, M.H. and Adir, N. 2004. The proteasome and the delicate balance between destruction and rescue. *PLoS Biology.* 2: 25-27.
- Glickman, M.H., Rubin, D.M., Coux, O., Wefes, I., Pfeifer, G., Cjeka, Z., Baumeister, W., Fried, V.A. and Finley, D. 1998. A subcomplex of the proteasome regulatory particle required for ubiquitin-conjugate degradation and related to the COP9 signalosome and eIF3. *Cell.* 94: 615-623.
- Glickman, M.H., Rubin, D.M., Fu, H., Larsen, C.N., Coux, O., Wefes, I., Pfeifer G., Cjeka, Z., Vierstra, R., Baumeister, W., Fried, V. and Finley, D. 1999. Functional analysis of the proteasome regulatory particle *Mol Biol Reports* 26: 21-28.
- Glover, J.R. and Lindquist, S. 1998. Hsp104, Hsp70, and Hsp40: A novel chaperone system that rescues previously aggregated proteins. *Cell* 94: 73-82.
- Goldberg, A.L. 2003. Protein degradation and protection against misfolded or damaged proteins. *Nature* 426: 895-899.
- Gordon, C.L., Khalaj, V., Ram, A.F., Archer D.B., Brookman, J.L., Trinci, A.P., Jeenes, D.J., Doonan, J.H., Wells, B., Punt, P.J., van den Hondel, C.A. and Robson, G.D. 2000. Glucoamylase::green fluorescent protein fusions to monitor protein secretion in *Aspergillus niger*. *Microbiology.* 146: 415-426.
- Gouka, R.J., Punt, P.J. and van den Hondel, C.A. 1997. Efficient production of secreted proteins by *Aspergillus*: progress, limitations and prospects. *Appl. Microbiol. Biotechnol.* 47: 1-11.
- Grinyer, J., Kautto, L., Traini, M., Willows, R.D., Teodoro, J., Bergquist, P. and Nevalainen, H. 2007. Proteome mapping of the *Trichoderma reesei* 20S proteasome, *Curr. Genet.* 51: 79-88.

- Gritz, L. and Davies, J. 1983. Plasmid encoded hygromycin B resistance: the sequence of hygromycin B phosphotransferase gene and its expression in *Escherichia coli* and *Saccharomyces cerevisiae*. *Gene*. 25: 179-183.
- Goller, S.P., Gorfer, M., Kubicek, C.P. 1998. *Trichoderma reesei* prs12 encodes a stress- and unfolded-protein-response-inducible regulatory subunit of the fungal 26S proteasome. *Curr. Genet.* 33: 284-290.
- Groll, M., Ditzel, L., Lowe, J., Stock, D., Bochtler, M., Bartunik, H.D. and Huber, R. 1997. Structure of 20S proteasome from yeast at 2.4 Å resolution. *Nature*. 386: 463-471.
- Groll, M., Heinemeyer, W., Jäger, S., Ullrich, T., Bochtler M., Wolf, D.H. and Huber R. 1999. The catalytic sites of 20S proteasomes and their role in subunit maturation: A mutational and crystallographic study. *Proc. Natl. Acad. Sci.* 96: 10976-10983.
- Groll, M., Bajorek, M., Kohler, A., Moroder, L., Rubin, D.M., Huber, R., Glickman, M.H. and Finley, D. 2000. A gated channel into the proteasome core particle. *Nat. Struct. Biol.* 7: 1062-1067.
- Groll, M. and Huber, R. 2003. Substrate access and processing by the 20S proteasome core particle. *Int. J. Biochem. Cell Biol.* 35: 606-616.
- Groll, M., Schellenberg, B., Bachmann, A.S., Archer, C.R., R Huber, R., Powell, T.K., Lindow, S., Kaiser, M. And Dudler, R. 2008. A plant pathogen virulence factor inhibits the eukaryotic proteasome by a novel mechanism. *Nature* 452: 755-759.
- Guerrero, C., Tagwerker, C., Kaiser, P. and Huang, L. 2006 An integrated mass spectrometry base proteomic approach: Quantitative analysis of tandem affinity-purified *in vivo* cross-linked protein complexes (QTAX) to decipher the 26S proteasome-interacting network. *Mol. Cell Proteom.* 5: 366-378.
- Guerrero, C., Milenkovic, T., Pržulj, N., Kaiser, P. and Huang, L. 2008. Characterization of the proteasome interaction network using QTAX-based tag-team strategy and protein interaction network analysis. *PNAS*. 105: 13333-13338.
- Guerrera, I. and Kleiner, O. 2005. Application of mass spectrometry in proteomics. *Bioscience Reports*. 25: 71-93.
- Guillemette, T., van Peij, N.N.M.E, Goosen, T., Lanthaler, K., Robson, G.D., Hondel, C.A.M.J.J. van den, Stam, H. and Archer, D.B. 2007. Genomic analysis of the secretion stress response in the enzyme-producing cell factory *Aspergillus niger*. *BMC Genomics*. 8: 158-175.
- Gupta, G.D. and Heath, I.B. 2000. A Tip-high gradient of a putative plasma membrane SNARE approximates the exocytotic gradient in hyphal apices of the fungus *Neurospora crassa*. *Fungal Gen. Biol.* 29: 187-199.

- Guterman, A. and Glickman, M.H. 2004. Complementary roles for Rpn11 and Ubp6 in deubiquitination and proteolysis by the proteasome. *J. Biol. Chem.* 279: 1729-1738.
- Gülow, K., Bienert, D. and Haas, I.G. 2002. BiP is feed-back regulated by control of protein translation efficiency. *J. Cell Science.* 115: 2443-2452.

H

- Haab, D., Hagspiel, K., Kubicek, C.P. Protease activity and proteolytic modification of cellulases from a *Trichoderma reesei* QM 9414 selectant. *Applied Microbiology and Biotechnology.* 32: 64-67.
- Haigh, N.G. and Johnson, A.E. 2002. A new role for BiP: closing the aqueous translocon pore during protein integration into the ER membrane. *J. Cell Biol.* 156: 261-270.
- Haiping Zhou, Yuying Zhang, Cuijuan Jia and Kaiyu Yang. 2004. Chaperone characteristics of PDI-related protein A from *Aspergillus niger*. *Biochem Biophys Research Comm.* 321: 31-3.
- Harding, H.P., Zhang, Y. and Ron, D. 1999. Protein translation and folding are coupled by an endoplasmic-reticulum-resident kinase. *Nature* 397: 271-274.
- Harding, H.P., Novoa I., Zhang Y., Zeng H., Wek R., Schapira M., and Ron D. 2000. Regulated translation initiation controls stress-induced gene expression in mammalian cells. *Mol. Cell.* 6: 1099-1108.
- Harkki, A., Uusitalo, J., Bailey M., Penttilä, M. and Knowles, C.K.J. 1989. A novel fungal expression system: Secretion of active calf chymosin from the filamentous fungus *Trichoderma reesei*. *BioTechnology.* 7: 596-603.
- Harris, J.R. 1968. Release of a macromolecular protein component from human erythrocyte ghosts. *Biochim. Biophys. Acta.* 150: 530-537.
- Hammond, C. and Helenius, A. 1994. Quality control in the secretory pathway: retention of a misfolded viral membrane glycoprotein involves cycling between the ER, intermediate compartment, and Golgi apparatus. *J. Cell Biol.* 126: 41-52.
- Hampton, R.Y. 2002. ER-associated degradation in protein quality control and cellular regulation. *Curr. Opin. Cell Biol.* 14: 476-482.
- Hann, B. and Walter, P. 1991. The signal recognition particle in *S. cerevisiae*. *Cell* 67: 131-144.
- He, X., Wang, K., Tan, W., Xiao, D., Li, J. and Yang, X. 2001. A novel fluorescent label based on biological fluorescent nanoparticles and its application in cell recognition. *Chinese Bulletin* 41: 1962-1965.

- Hedge, A.N. and Upadhyay, S.C. 2006. Proteasome and transcription: a destroyer goes into construction. *BioEssays* 28: 235-239.
- Heinemeyer, W., Fischer, M., Krimmer, T., Stachon, U. and Wolf, D.H. 1997. The active sites of the eukaryotic 20S proteasome and their involvement in subunit precursor processing. *J. Biol. Chem.* 272: 25200-25209.
- Heinemeyer, W., Ramos, P.C. and Dohmen, R.J. 2004. The ultimate nanoscale mincer: assembly, structure and active sites of 20S proteasome core. *Cell Mol. Life Sci.* 61: 1562-1578.
- Hendil, K.B., Welinder, K.G., Pedersen, D., Uerkevitz, W. and Kristensen, P. 1993. Subunit stoichiometry of human proteasomes. *Enzyme* 47: 232-240.
- Henderson, G.S., Conary, J.T., Davidson, J.M., Stewart, S.J., House, F.S. and McCurley, T.L. 1991. A reliable method for northern blot analysis using synthetic oligonucleotide probes. *Biotechniques* 10: 190-197.
- Helenius, A. and Aeby, M. 2004. Roles of N-linked glycans in the endoplasmic reticulum. *Annu. Rev. Biochem.* 73: 1019-1049.
- Henriksson, H., Muñoz, I.G., Isaksson, R., Pettersson, G. and Johansson, G. 2000. Cellobiohydrolase 58 (P.c. Cel 7D) is complementary to the homologous CBH I (T.r. Cel 7A) in enantioseparations. *J Chromatography A*. 898: 63-74.
- Herbert, B., Hopwood, F., Oxley, D., McCarthy, J., Laver, M., Grinyer, J., Goodall, A., Williams, K., Castagna, A. and Righetti, P.G. 2003. Beta-elimination: an unexpected artefact in proteome analysis. *Proteomics* 3: 826-831.
- Hershko, A. and Ciechanover, A. 1998. The ubiquitin system. *Annu. Rev. Biochem.* 67: 425-479.
- Hijarrubia, M.J., Casqueiro, J., Gutiérrez, S., Fernández, F.J. and Martín, J.F. 1997. Characterization of the *bip* gene of *Aspergillus awamori* encoding a protein with an HDEL retention signal homologous to the mammalian BiP involved in polypeptide secretion. *Curr. Gen.* 32:139-46.
- Higgins, V.J., Beckhouse, A.G., Oliver, A.D., Rogers, P.J. and Dawes, I.W. 2003. Yeast genome-wide expression analysis identifies a strong ergosterol and oxidative stress response during the initial stages of an industrial lager fermentation. *Appl. Environ. Microbiol.* 69: 4777-4787.
- Hiller, M.M., Finger, A., Schweiger, M. and Wolf, D.H. 1996. ER degradation of a misfolded luminal protein by the cytosolic ubiquitin-proteasome pathway. *Science* 273: 1725-1728.
- Hilt, W. and Wolf, D. H. 1999. Proteasome: destruction a programme. *Trends Biochem. Sci.* 21: 96-102.

- Hinnebush, A.G. 1997. Translational regulation of yeast GCN4. A window on factors that control initiator-tRNA binding to ribosome J. Biol. Chem. 272: 21661-21664.
- Hirano, Y., Hendil, K.B., Yashiroda, H., Iemura, S., Nagane, R., Hioki, Y., Natsume, T., Tanaka, K. and Murata, S. 2005. A heterodimeric complex that promotes the assembly of mammalian 20S proteasomes. Nature. 437: 1381-1385.
- Ho, M., Tsai, P-I., and Chien, C-T. 2006. F-box protein: key to protein degradation. J. Biochem. Sci. 13: 181-191.
- Hochstrasser, M. 2000. Evolution and function of ubiquitin-like protein conjugation systems. Nature Cell Biol. 2: E153-E157.
- Hofmann K. and Bucher P. 1998. The PCI domain: a common theme in three multiprotein complexes. Trends Biochem Sci. 23: 204-205
- Hosokawa, N., Wada, I., Natsuka, Y., Nagata, K. 2006. EDEM accelerates ERAD by preventing aberrant dimer formation of misfolded alpha1-antitrypsin. Genes Cells. 11: 465-476.
- Hough, R., Pratt, G. and Rechsteiner, M. 1987. Purification of two high molecular weight proteases from rabbit reticulocyte lysate. J. Biol. Chem. 262: 8303-8313.

I

- Imai, J., Maruya, M., Yashiroda, H., Yahara, I. and Tanaka, K. 2003. The molecular chaperone Hsp90 plays a role in the assembly and maintenance of the 26S proteasome. EBMO J. 22: 3557-3567.

J

- Jensch, S. and Pyrowolakis, G. 2000. Ubiquitin and its kin: how close are the family ties? Trends Cell Biol. 10:335-42.
- Johnston, J.A., Ward, C.L., and Kopito R.R. 1998. Aggresomes: A cellular response to misfolded proteins. J. Cell Biol. 143: 1883-1898.
- Joseph, J. D., Heitman, J. and Means, A.R. 1999. Molecular cloning and characterization of *Aspergillus nidulans* cyclophilin B. Fungal Gen. Biol. 27: 55-66.
- Jørgensen, T.R., Goosen, T., van den Hondel C.A.M.J.J., Ram, A.F.J. and Iversen, J.J.L. 2009. Transcriptomic comparison of *Aspergillus niger* growing on two different sugars reveals coordinated regulation of the secretory pathway. BMC Genomics. 10: 44-56.

K

- Kahle, P.J., Leimer, U. and Haass, C. 2000. Does failure of parkin-mediated ubiquitination cause juvenile parkinsonism? Trends Biochem Sci. 25: 524-527.

- Kanehara, K., Kawaguchi, S. and Ng, D.T.W. 2007. The EDEM and Yos9p families of lectin-like ERAD factors. *Cell Dev. Biol.* 18: 743-750.
- Kaufman, R.J. 1999. Stress signaling from the lumen of the endoplasmic reticulum: coordination of gene transcriptional and translational controls. *Genes Dev.* 13: 1211-1233.
- Kautto, L., Grinyer, J., Birch, D., Kapur, A., Baker, M., Bergquist, P. and Nevalainen, H. 2009. Rapid purification method for the 26S proteasome from the filamentous fungus *Trichoderma reesei*. *Protein Expression and Purification* 67: 156-163.
- Karhunen, T., Mäntylä, A., Nevalainen, K.M.H. and Suominen, P.L. 1993. High frequency one-step gene replacement in *Trichoderma reesei*. I: Endoglucanase overproduction. *Mol. Gen. Genet.* 241: 515-522.
- Karpov D.S., Osipov S.A., Preobrazhenskaya, O.V. and Karpov, V.L. 2008. Rpn4p is a positive and negative transcriptional regulator of the ubiquitin-proteasome system. *Mol. Biol.* 42: 456-462.
- Katakura Y., Ametani, A., Totsuka, M., Nagafuchi, S. and Kaminogawa, S. 1999. Accelerated secretion of mutant β -lactoglobulin in *Saccharomyces cerevisiae* resulting from a single amino acid substitution. *Prot. Struc. Mol. Enz.* 1432: 302-312.
- Kawahara, T., Yanagi, H., Yura, T. and Mori, K. 1997. Endoplasmic reticulum stress-induced mRNA splicing permits synthesis of transcription factor Hac1p/Ern4p that activates the unfolded protein response. *Mol. Biol Cell.* 8: 1845-1862.
- Kawazoe Y., Nakai A., Tanabe M. and Nagata K. 1998. Proteasome inhibition leads to the activation of all members of heat shock factor family. *Eur. J. Biochem.* 255: 356-362.
- Keränen, S. and Penttilä, M. 1995. Production of recombinant proteins in the filamentous fungus *Trichoderma reesei*. *Curr. Opin. Biotech.* 6: 534-537.
- Khalaj, V., Brookman J.L. and Robson, G.D. 2001. A Study of the protein secretory pathway of *Aspergillus niger* using a glucoamylase-GFP fusion protein. *Fungal Gen. Biol.* 32: 55-65.
- Kiiskinen, L-L., Kruus, K., Bailey, M., Ylösmäki, E., Siika-aho, M. and Saloheimo, M. 2004. Expression of *Melanocarpus albomyces* laccase in *Trichoderma reesei* and characterization of the purified enzyme. *Microbiology.* 150: 3045-3074.
- Kim, J.H., Park, K.C., Chung, S.S., Bang, O. and Chung, C.H. 2003. Deubiquitinating enzymes as cellular regulators. *J. Biochem.* 134: 9-18.

- Kim, S., Ahl, I.P., and Lee, Y.H. 2001. Analysis of genes expressed during rice-*Magnaporthe grisea* interactions. *Mol. Plant Microb. Interact.* 14: 1340-1346.
- Kimura, Y., Saeki, Y., Yokosawa, H., Polevoda, B., Sherman, F. and Hirano, H. 2003. N-Terminal modifications of the 19S regulatory particle subunits of the yeast proteasome. *Arch. Biochem. Biophys.* 409: 341-348.
- Kirisako, T., Kamei, K., Murata, S., Kato, M., Fukumoto, H., Kanie, M., Sano, S., Tokunaga, F., Tanaka, K. and Iwai, K. 2006. A ubiquitin ligase complex assembles linear polyubiquitin chains. *EMBO J.* 25: 4877-4887.
- Kleizen, B. and Braakman, I. 2004. Protein folding and quality control in the endoplasmic reticulum. *Curr. Opin. Cell Biol.* 16: 343-349.
- Kleymann, G., Ostermeier, C., Heitmann, K., Haase, W. and Michel, H. 1995. Use of antibody fragments (Fv) in immunocytochemistry. *J. Histochem. Cytochem.* 43: 607-614.
- Knight, B. C. and High, S. 1998. Membrane integration of Sec61 α : a core component of the endoplasmic reticulum translocation complex. *Biochem. J.* 331: 161-167.
- Knop, M., Finger, A., Braun, T., Hellmuth, K. and Wolf, D.H. 1996. Der1, a novel protein specifically required for endoplasmic reticulum degradation in yeast. *EMBO J.* 15: 753-763.
- Knowles T.P.J. and Zahn R. 2006. Enhanced Stability of Human Prion Proteins with Two Disulfide Bridges. *Biophys. J.* 91: 1494-1500.
- Kobayashi, K., Kuwae, S., Ohya, T., Ohda, T., Ohyama, M., Ohi, M., Tomomitsu, K. and Ohmura, T. 2000. High-level expression of recombinant human serum albumin from the methylotrophic yeast *Pichia pastoris* with minimal protease production and activation. *J. Biosci. Bioeng.* 89: 55-61.
- Kohno, H.H. 2008. A genetic link between unfolded protein response and vesicle formation from endoplasmic reticulum. *Biochem. Biophys. Commun.* 296: 568-574.
- Koivula, A., Ruohonen, L., Wohlfahrt, G., Reinikainen, T., Teeri, T.T., Piens, K., Claeysens, M., Weber, B., Vasella, A., Becker, D., Sinnott, M.L., Zou, J., Kleywegt, G., Szardenings, M., Stahlberg, J. and Jones, T.A. 2002. The active site of the cellobiohydrolase Cel6A from *Trichoderma reesei*: The roles of the aspartic acids D221 and D175. *J. Am. Chem. Soc.* 124: 10015-10024.
- Koll, H., Guiard, B., Rassow, J., Ostermann, J., Horwich, A., Neupert, W. and Hartl, F.U. 1992. Antifolding activity of HSP60 couples protein import into the mitochondrial matrix with export to the intermembrane space. *Cell* 68: 1163-1175.

- Kominami K., DeMartino G.N., Moomaw C., Slaughter C.A., Shimbara N., Fujimuro M., Yokosawa H., Hisamatsu H., Tanahashi N., Shimizu Y., Tanaka K., Toh-e A., 1995. Nplp, a regulatory subunit of the 26S proteasome, is necessary for activation of Cdc28p kinase of *Saccharomyces cerevisiae*. EMBO J. 14: 3105-3115.
- Kopp, F., Hendkl, K. B., Dahlmann, B., Kristensen, P., Sobek, A. and Uerkvitz, W. 1997. Subunit arrangement in the human 20S proteasome. Proc. Natl. Acad. Sci. 94: 2939-2944.
- Koushik, S.V., Chen, H., Thaler, C., Puhl, H.L. and Vogel, S.S. 2006. Cerulean, Venus, and VenusY67C FRET Reference Standards. Biophys. J. 91: L99-L101.
- Kornitzer, D., Raboy, B., Kulka, R. G. and Fink, G. R. 1994. Regulated degradation of the transcription factor Gcn4. EMBO J. 13: 6021-6030.
- Kostova, Z. and Wolf, D.H. 2002. Protein quality control in the export pathway: the endoplasmic reticulum and its cytoplasmic proteasome connection. pp. 180-213. In Dalbey R. E. and von Heijne G. (Eds.). Protein Targeting, Transport and Translocation. Academic Press, London, UK.
- Kostova, Z. and Wolf, D.H. 2003. For whom the bell tolls: protein quality control of the endoplasmic reticulum and the ubiquitin-proteasome connection. EMBO J. 22: 2309-2317.
- Kowalski J.M., Parekh, R.N., Mao, J. and Wittrup, K.D. 1998. Protein folding stability can determine the efficiency of escape from endoplasmic reticulum quality control. The Journal of Biological Chemistry 273: 19453-19458.
- Krogan, N. J., Lam, M.H.Y., Fillingham, J., Keogh, M-C., Gebbia, M., Li, J., Datta, N., Cagney, G., Buratowski, S., Emili, A., and Greenblatt J. F. 2004. Proteasome involvement in the repair of DNA double-strand breaks. Mol. Cell. 16: 1027-1034.
- Kuehn, M.J. and Schekman, R. 1997. COPII and secretory cargo capture into transport vesicles. Curr. Opin. Cell Biol. 9: 477-483.
- Kuhlbrodt K., Mouysset J. and Hoppe, T. 2005. Orchestra for assembly and fate of polyubiquitin chains. Essays Biochem. 41: 1-14.
- Kumita J.R., Johnson R.J.K., Alcocer, M.J.C., Dumoulin M., Holmqvist, F., McCammon, M.G., Robinson, C.V., Archer, D.B. and Dobson, C.M. 2006. Impact of the native-state stability of human lysozyme variants on protein secretion by *Pichia pastoris*. FEBS J. 273: 711-720.
- Kuratsu, M., Taura, A., Shoji, J.Y., Kikuchi, S., Arioka, M. and Kitamoto, K. 2007. Systematic analysis of SNARE localization in the filamentous fungus *Aspergillus oryzae*. Fungal Genet. Biol. 44: 1310-1323

- Kurucz, E. A. I., Sumegi, M., Holzl, H., Kapelari, B., Baumeister, W. and Udvardy, A. 2002. Assembly of the Drosophila 26 S proteasome is accompanied by extensive subunit rearrangements. *Biochem. J.* 365: 527-36.
- Kurzatkowski, W., Törrönen, A., Filipek, J., Mach, R.L., Herzog, P., Sowka, S. and Kubicek, C.P. Glucose-Induced Secretion of *Trichoderma reesei* Xylanases. *Appl. Env. Microbiol.* 62: 2859-2865.
- Köhler, A., Cascio, P., Leggett, D.S., Woo, K.M., Goldberg, A.L. and Finley, D. 2001. The axial channel of the proteasome core particle is gated by the Rpt2 ATPase and controls both substrate entry and product release. *Mol. Cell.* 7: 1143-1152.

L

- Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature.* 227: 680-685.
- Lam, Y.A., DeMartino, G.N., Pickart, C.M. and Cohen, R.E. 1997. Specificity of the ubiquitin isopeptidase in the PA700 regulatory complex of 26 S proteasomes. *J. Biol. Chem.* 272: 28438-28446.
- Lam, Y.A., Lawson, T.G., Velayutham, M., Zweier, J.L. and Pickart, C.M. 2002. A proteasomal ATPase subunit recognizes the polyubiquitin degradation signal. *Nature* 416: 763-767.
- Laney, J.D. and Hochstrasser, M. 2003. Ubiquitin-dependent degradation of the yeast Mat α 2 repressor enables a switch in developmental state. *Genes Dev.* 17: 2259-2270.
- Lashkari, D.A., DeRisi, J.L., McCusker, J.H., Namath, A.F., Gentile, C., Hwang, S.Y., Brown, P.O. and Davis, R.W. 1997. Yeast microarrays for genome wide parallel genetic and gene expression analysis. *Proc. Nat. Acad. Sci. USA.* 94: 13057-13062.
- Leber, J. H., S. Bernales, and P. Walter. 2004. IRE1-independent gain control of the unfolded protein response. *PLoS Biol.* 2:1197-1207.
- Lee, D.H. and Goldberg, A.L. 1996. Selective inhibitors of the proteasome-dependent and vacuolar pathways of protein degradation in *Saccharomyces cerevisiae*. *J. Biol. Chem.* 271: 27280-27284.
- Lee, D.H. and Goldberg, A.L. 1998. Proteasome inhibitors: valuable new tools for cell biologists. *Trends Cell Biol.* 8: 397-403.
- Lee, D., Ezhkova, E., Li, B., Pattenden, S.G., Tansey, W.P. and Workman, J.L. 2005. The proteasome regulatory particle alters the SAGA coactivator to enhance its interactions with transcriptional activators. *Cell* 123: 423-436.

- Lee, S.B., Milgroom, M. G., and Taylor, J.W. 1988. A rapid, high yield mini prep method for isolation of total genomic DNA from fungi. *Fungal Genet. Newsl.* 35: 23-24.
- Le Tallec, B., Barrault, M-B., Guérois, R., Carré, T. and Peyroche, A. 2009. Hsm3/S5b participates in the assembly pathway of the 19S regulatory particle of the proteasome. *Mol. Cell.* 33: 389-399.
- Legget, D.S., Glickman, M.H. and Finely, D. 2005. Purification of proteasomes, proteasome subcomplexes, and proteasome-associated proteins from budding yeast. pp 57-70. In: *Methods in Molecular Biology*. Eds. Patterson, C. and Cyr, D.M. Vol. 301, Ubiquitin-Proteasome Protocols. Humana Press Inc., Totowa, NJ.
- Li, B., Brown, E.W., D'Agostino, C., LeClerc J.E. and Cebula T.A. 2005. Structure and distribution of the phosphoprotein phosphatase genes, *prpA* and *prpB*, among shigella subgroups. *Microbiology.* 151: 2671-2683.
- Liao, W., Li, X., Mancini, M. and Chan, L. 2006. Proteasome inhibition induces differential heat shock protein response but not unfolded protein response in HepG2 cells. *J. Cell. Biochem.* 99: 1085-1095.
- Lim, D., Hains, P., Walsh, B., Bergquist P. and Nevalainen, H. 2001. Proteins associated with the cell envelope of *Trichoderma reesei*: a proteomic approach. *Proteomics* 1: 899-910.
- Linder, M.B., Szilvay, G.R., Nakari-Setälä, T. and Penttilä, M. 2005. Hydrophobins: the protein-amphiphiles of filamentous fungi. *FEMS Microb. Rev.* 29: 877-896.
- Lipshutz, R.J., Fodor, S.P., Gingeras, T.R. and Lockhart, D.J. 1999. High density synthetic oligonucleotide arrays. *Nat. Genet.* 21: 20-4.
- Löwe, J., Stock, D., Jap, B., Zwickl, P., Baumeister, W. and Huber, R. 1995. Crystal structure of the 20S proteasome from the archaeon *T. acidophilum* at 3.4 Å resolution. *Science* 268: 533-539.

M

- Ma, Y. and Hendershot, L.M. 2001. The unfolding tale of the unfolded protein response. *Cell* 107: 827-830.
- Mackenzie D.A., Guillemette T., Al-Sheikh H., Watson A. J., Jeenes D.J., Wongwathanarat P., Dunn-Coleman N.S., Van Peij N. and Archer, D.B. 2005. UPR-independent dithiothreitol stress-induced genes in *Aspergillus niger*. *Mol. Gen. Genomics.* 274:410-418.
- MacLaren, A.P., Chapman, R.S., Wyllie, A.H. and Watson, C.J. 2001. p53-dependent apoptosis induced by proteasome inhibition in mammary epithelial cells. *Cell Death Diff.* 8: 210-218.

- Madeo, F., Fröhlich, E. and Fröhlich K-U. 1997. A yeast mutant showing diagnostic markers of early and late apoptosis. *J. Cell Biol.* 139: 729-734.
- Majetshak, M. and Sorell L. 2008. Immunological methods to quantify and characterize proteasome complexes: development and application. *J Immunological Methods* 334: 91-103.
- Malecki, M., Hsu, A., Truong, L. and Sanchez, S. 2001. Molecular immunolabeling with recombinant single-chain variable fragment (scFv) antibodies designed with metal-binding domains. *SNAS.* 99: 213-218.
- Marcolles-Clark, E., Hayes C.K., Harman, G.E. and Penttilä, M. 1996. Improved production of *Trichoderma harzianum* endochitinase by expression in *Trichoderma reesei*. 1996. *Appl. Env. Microbiol.* 62: 2145-2151.
- Marquardt, T., Hebert, D.N. and Helenius, A. 1993. Post-translational folding of influenza hemagglutinin in isolated endoplasmic-reticulum-derived microsomes. *J. Biol. Chem.* 268: 19618-19625.
- Marquardt, T. and Helenius, A. 1992. Misfolding and aggregation of newly synthesized proteins in the endoplasmic reticulum. *J. Cell Biol.* 117: 505-513.
- Martin-Magniette M-L., Aubert, J., Bar-Hen, A., Elftieh, S., Magniette, F., Renou, J-P. and Daudin, J-J. 2008. Normalization for triple-target microarray experiments. *BMC Bioinformatics.* 9: 216.
- Martinez, I.M. and Chrispeels, M.J. 2003. Genomic analysis of the unfolded protein response in *Arabidopsis* shows its connection to important cellular processes. *Plant Cell.* 15:561-576.
- Martinez, D., Berka, R.M., Henrissat, B., Saloheimo, M., Arvas, M., Baker, S.E., Chapman, J., Chertkov, O., Coutinho, P.M, Cullen, D., Danchin, E.G.J., Grigoriev, I.V., Harris, P., Jackson, M., Kubicek, C.P., Han, C.S., Ho, I., Larrondo, L.F., de Leon, A.L., Magnuson, J.K., Merino, S., Misra, M., Nelson, B., Putnam, N., Robbertse, B., Salamov, A.A., Schmoll, M., Terry, A., Thayer, N., Westerholm-Parvinen, A., Schoch, C.L., Yao, J., Barabote, R., Nelson, M.A., Detter, C., Bruce, D., Kuske, C.R., Xie, G., Richardson, P., Rokhsar, D.K., Lucas, S.M., Rubin, E.M., Dunn-Coleman, N., Ward, M., and Brettin, T.S. 2008. Genome sequencing and analysis of the biomass-degrading fungus *Trichoderma reesei* (syn. *Hypocrea jecorina*). *Nature Biotech.* 26: 553-560.
- Maruyama J. and Kitamoto, K. 2006. Differential distribution of the endoplasmic reticulum network in filamentous fungi. *FEMS.* 272: 1-7.
- Masai, K., Maryama, J., Nakajima, H. and Kitamoto, K. 2003. *In vivo* visualization of the distribution of a secretory protein in *Aspergillus oryzae*. *Fungal Genet. Biol.* 43: 642-654.

- Mast, S.W., Diekman, K., Karaveg, K., Davis, A., Sifers, R.N. and Moremen, K.W. 2005. Human EDEM2, a novel homolog of family 47 glycosidases, is involved in ER-associated degradation of glycoproteins. *Glycobiology*. 15: 421-436.
- Mathews, C.K. and van Holde, K.E. 1990. Dynamics of Life: Catalysis and control of biochemical reactions. In *Biochemistry*. The Benjamin/Cummings Publishing Company. Redwood City. CA.USA.
- Matsuura, K. and Takagi, H. 2005. Vacuolar Functions are involved in stress-protective effect of intracellular proline in *Saccharomyces cerevisiae*. *J Bioscience and Bioengineering* 100: 538-544.
- Matouschek, A., Rospert, S., Schmid, K., Glick, B.S. and Schatz, G. 1995. Cyclophilin catalyzes protein folding in yeast mitochondria. *Proc. Natl. Acad. Sci.* 92: 6319-6323.
- Mayer, T.U., Braun, T. and Jentsch, S. 1998. Role of the proteasome in membrane extraction of a short-lived ER-transmembrane protein. *EMBO J.* 12: 3251-3257.
- Mayor, T., Lipford, R.J., Graumann, J., Smith, G.T. and Deshaies, R.J. 2005. Analysis of polyubiquitin conjugates reveals that the Rpn10 substrate receptor contributes to the turnover of multiple proteasome targets. *Mol. Cell. Proteomics*. 4: 741-751.
- McDonough, H. and Patterson, C. 2003. CHIP: A link between the chaperone and proteasome systems. *Cell Stress Chaperones*. 8: 303-308.
- McCracken, A.A. and Brodsky, J.L. 2000. A molecular portrait of the response to unfolded proteins. *Gen. Biol.* 1: Reviews 1013.1-1013.3
- Merksamer, P.I., Trusina, A. and Papa, F.R. 2008. Real-time redox measurement during endoplasmic reticulum stress reveal interlinked protein folding functions. *Cell* 35: 933-947.
- Meusser, B., Hirsch, C., Jarosch, E. and Sommer, T. 2005. ERAD: the long road to destruction. *Nat. Cell Biol.* 7: 766-772.
- Mezey, E., Rennie-Tankersley, L. and Potter, J.J. 2001. Liver alcohol dehydrogenase is degraded by ubiquitin-proteasome pathway. *Biochem. Biophys. Res. Commun.* 285: 644-648.
- Miettinen-Oinonen, A. 2004. *Trichoderma reesei* strains for production of cellulases for the textile industry. PhD thesis. VTT publication 550. VTT, Espoo, Finland.
- Miyoshi, K., Miyakawa, T. and Mizuta, K. 2001. Repression of rRNA synthesis due to a secretory defect requires the C-terminal silencing domain of Rap1p in *Saccharomyces cerevisiae*. *Nucleic Acids Res.* 29: 3297-3303.

- Mizuta, K., and Warner, J.R. 1994. Continued functioning of the secretory pathway is essential for ribosome synthesis. *Mol. Cell Biol.* 14: 2493-2502.
- Molinari, M., Eriksson, K.K., Calanca, V., Galli, C., Cresswell, P., Michalak, M. and Helenius, A. 2004. Contrasting functions of calreticulin and calnexin in glycoprotein folding and ER quality control. *Mol. Cell* 13: 125-135.
- Monti, M., Orrù, S., Pagnozzi, D. and Pucci, P. 2005. Interaction proteomics. *Biosci. Rep.* 25: 45-56.
- Moralejo, F.J., Cardoza R-E, Gutierrez S, and Martin, J.F. 1999. Thaumatin Production in *Aspergillus awamori* by use of expression cassettes with strong fungal promoters and high gene dosage. *Appl. Environ. Microbiol.* 65: 1168-1174.
- Moralejo, F.J., Watson, A.J., Jeenes, D.J., Archer, D.B., and Martin, J.F. 2001. A defined level of protein disulfide isomerase expression is required for optimal secretion of thaumatin by *Aspergillus awamori*. *Mol. Genet. Genomics.* 266: 246-253.
- Morris, M.C., Kaiser, P., Stanislav Rudyak, S., Baskerville, C., Watson, M.H. and Reed, S.I. 2003. Cks1-dependent proteasome recruitment and activation of *CDC20* transcription in budding yeast. *Nature* 423: 1009-1013.
- Mulder, H.J., Saloheimo, M., Penttilä, M. and Madrid, S.M. 2004. The transcription factor HACA mediates the unfolded protein response in *Aspergillus niger*, and up-regulates its own transcription. *Mol. Genet. Genomics* 271: 130-140.
- Murata, T. and Shimotohno, K. 2006. Ubiquitination and proteasome-dependent degradation of human eukaryotic translation initiation factor 4E. *J. Biol. Chem.* 281: 20788-20800.
- Muratani, M. and Tansey, W.P. 2003. How the ubiquitin-proteasome system controls transcription. *Nat. Rev. Mol. Cell Biol.* 4: 192-201
- Müsch A., Wiedmann M. and Rapoport T.A. 1992. Yeast Sec proteins interact with polypeptides traversing the endoplasmic reticulum membrane. *Cell* 2: 343-352.
- Myung, J., Kim, K.B. and Crews C.M. 2001. The ubiquitin-proteasome pathway and proteasome inhibitors. *Med. Res. Rev.* 21: 245-273.

N

- Nakatsukasa K. and Brodsky J.L. 2008. The recognition and retrotranslocation of misfolded proteins from the endoplasmic reticulum. *Traffic.* 9: 861-870.
- Nagai, T., Ibata, K., Park, E.S., Kubota, M., Mikoshiba, K. and Miawaki, A. 2002. A variant of yellow fluorescent protein with fast and efficient maturation for cell-biological applications. *Nat. Biotech.* 20: 87-90.

- Nandi, D., Tahiliani, P., Kumar, A. and Chandu, D. 2006. The ubiquitin-proteasome system. *J. Biosci.* 31: 137-155.
- Nebenführ, A. and Lomax T.L. 2004. Multiplex titration RT-PCR: Rapid determination of gene expression patterns for a large number of genes. *Plant Mol. Biol. Rep.* 16: 323-339.
- Ng, D.T., Brown, J.D. and Walter, P. 1996. Signal sequences specify the targeting route to the endoplasmic reticulum membrane. *J. Cell Biol.* 134: 269-278.
- Ngiam, C., Jeenes, D.J. and Archer, D.B. 1997. Isolation and characterization of a gene encoding protein disulphide isomerase, *pdiA*, from *Aspergillus niger*. *Curr. Genet.* 31: 133-138.
- Nguyen, T.H., Law, D.T. and Williams, D.B. 1991. Binding protein BiP is required for translocation of secretory proteins into the endoplasmic reticulum in *Saccharomyces cerevisiae*. *PNAS.* 88:1565-1569.
- Nishikawa, S., Hirata, A., and Nakano, A. 1994. Inhibition of endoplasmic reticulum (ER) to-Golgi transport induces relocation of binding protein (BiP) within the ER to form the BiP bodies. *Mol. Biol. Cell* 5: 1129-1143.
- Nolan, T., Hands, R.E. and Bustin, S.E. 2009. Quantification of mRNA using real-time RT-PCR. *Nature protocols.* 1: 1559-1582.
- Novick, P., Garrett, M.D., Brennwald, P., Luring, A., Finger, F.P., Collins, R. and TerBush, D.R. 1995. Control of exocytosis in yeast. *Cold Spring Harb. Symp. Quant. Biol.* 60: 171-177.
- Nykänen, M., Saarelainen, R., Raudaskoski, M., Nevalainen K.H.M. and Mikkonen, A. 1997. Expression and secretion of barley cysteine endopeptidase B and cellobiohydrolase I in *Trichoderma reesei*. *Appl. Env. Microbiol.* 63: 4929-4937.
- Nyyssönen, E., Penttilä, M., Harkki, A., Saloheimo, A., Knowles J. K. C. and Keränen, S. 1993. Efficient production of antibody fragments by the filamentous fungus *Trichoderma reesei*. *Biotech.* 11: 591-595.

O

- Oda Y., Okada T., Yoshida H., Kaufman R.J., Nagata K., and Mori K. 2006. Derlin-2 and derlin -3 are regulated by the mammalian unfolded protein response and are required for the ER-associated degradation. *J. Cell Biol.* 3: 383-393.
- Ogawa, N. and Mori, K. 2004, Autoregulation of the HAC1 gene is required for sustained activation of the yeast unfolded protein response. *Genes Cells* 9: 95-104.

- Oka M., Yanagawa Y., Asada T., Yoneda A., Hasezawa S., Sato T. and Nakakawa H. 2004. Inhibition of proteasome by MG-132 treatment causes extra phragmoplast formation and cortical microtubule disorganisation during M/G1 transition in synchronised tobacco cells. *Plant Cell Physiol.* 45: 1623-1632.
- Okada, T., Yoshida, H., Akazawa, R., Negishi, M. and Mori, K. 2002. Distinct roles of Atf6 and PERK in transcription during the mammalian unfolded protein response. *Biochem. J.* 366: 585-594.
- Oliver, J.D., Roderick, H.L., Llewellyn, D.H. and High, S. 1999. ERp57 Functions as a subunit of specific complexes formed with the ER lectins calreticulin and calnexin. *Mol. Biol. Cell.* 10: 2573-2582.
- Osteryoung, K.W. and Vierling, E. 1994. Dynamics of small heat shock protein distribution within the chloroplasts of higher plants. *J. Biol. Chem.* 269: 28676-28682.
- Ouyang, J., Yan, M., Kong D. and Xu L. 2006. A complete protein pattern of cellulase and hemicellulase genes in filamentous fungus *Trichoderma reesei*. *Biotech. J.* 1: 1266-1274.

P

- Pagant, S., Kung, L., Dorrington, M., Lee M. C.S. and Miller E.A. 2007. Inhibiting endoplasmic reticulum (ER)-associated degradation of misfolded Yor1p does not permit ER export despite the presence of a diacidic sorting signal. *Mol. Biol Cell.* 18: 3398-3413.
- Pakula, T., Laxell, M., Huuskonen, A., Uusitalo, J., Saloheimo, M. and Penttilä, M. 2003. The effects of drugs inhibiting protein secretion in the filamentous fungus *Trichoderma reesei*. Evidence for down-regulation of genes that encode secreted proteins in the stressed cells. *J. Biol. Chem.* 278: 45011-45020.
- Pakula, T., Uusitalo, J., Saloheimo, M., Salonen, K., Aarts, R.J. and Penttilä, M. 2000. Monitoring the kinetics of glycoprotein synthesis and secretion in the filamentous fungus *Trichoderma reesei*: cellobiohydrolase I (CBHI) as a model protein. *Microbiol.* 146: 223-232.
- Palombella, V.J., Rando, O.J., Goldberg, A.L. and Maniatis, T. 1994. The ubiquitin-proteasome pathway is required for processing the NF-kappa B1 precursor protein and the activation of NF-kappa B. *Cell* 78: 773-785.
- Parales, R. Ju, K., Rollefson, J.B. and Ditty, J.L. 2008. Bioavailability, chemotaxis and transport of organic pollutants. pp. 145-188. In *Microbial biodegradation: Genomics and Molecular Biology*. Ed. Diaz Eduardo. Horizon Scientific Press. UK.
- Pappin, D.J.C., Højrup, P. and Bleasby, A.J. 1993. Rapid identification of proteins by peptide mass fingerprinting. *Curr. Biol.* 3: 327-332.

- Payne, T., Hanfrey, C., Bishop, A.L., Michael, A.J., Avery, S.V. and Archer, D.B. 2008. Transcript-specific translational regulation in the unfolded protein response of *Saccharomyces cerevisiae*. FEBS Lett. 582: 503-509.
- Penttilä, M., Nevalainen, H., Rättö, M., Salminen, E., and Knowles, J. 1987. A versatile transformation system for the cellulolytic filamentous fungus *Trichoderma reesei*. Gene 61:155-164.
- Penttilä, M. 1998. Heterologous protein production in *Trichoderma*. In *Trichoderma & Gliocladium*, pp. 365-382. Eds. G.E. Harman and C.P. Kubicek. London: Taylor and Francis.
- Perkins, G., Drury, L.S. and Diffley, J.F. 2001. Separate SCF(CDC4) recognition elements target Cdc6p for proteolysis in budding yeast. EMBO J. 20: 4836-4845.
- Pickart, C.M. 1997. Targeting of substrates to the 26S proteasome. FASEB J. 11: 1055-1066.
- Pickart, C.M. 2001. Mechanisms underlying ubiquitination. Ann. Rev. Biochem. 70: 503-533.
- Pickart, C.M. and Eddins, M.J. 2004. Ubiquitin: structures, functions, mechanisms. Biochem. Biophys. Acta. 1695: 55-72.
- Pickart, C. and Fushman, D. 2004. Polyubiquitin chains: polymeric protein signals. Curr. Opin. Chem. Biol. 8: 610-616.
- Plath, K., Wilkinson, B. M., Colin J., Stirling, C.J. and Tom A. Rapoport, T.A. 2004. Interactions between Sec complex and Prepro- α -factor during posttranslational protein transport into the endoplasmic reticulum. Mol. Biol. Cell 15: 1-10.
- Plempner, R.K. and Wolf, D.H. 1999. Retrograde protein translocation: ERADication of secretory proteins in health and disease. Trends Biochem. Sci. 24: 266-270.
- Plempner, R.K., Bordallo, J., Deak, P.M., Taxis, C., Hitt, R. and Wolf, D.H. 1999. Genetic interactions of Hrd3p and Der3p/Hrd1p with Sec61p suggest a retro-translocation complex mediating protein transport for ER degradation. J. Cell Sci. 112: 4123-4134.
- Pleofsky-Vig, N. and Brambl, R. 1998. Characterisation of an 88-kDa heat shock protein of *Neurospora crassa* that interacts with Hsp30. J. Biol Chem. 273: 11335-11341.
- Pollack, J.K., Harris, S.D., and Marten, M. 2009. Autophagy in filamentous fungi. Fungal Gen. Biol. 46: 1-8.

Punt, P.J., van Gemenen, I.A., Drink-Kuijvenhoven, J., Hessing, J.G.M., van Muijlwijk-Harteveld, G.M., Beijersbergen, A., Verrips, C.T. and van den Hodel, C.A.M.J.J. 1998. Analysis of the role of the gene *bipA*, encoding the major endoplasmic reticulum chaperone protein in the secretion of homologous and heterologous protein in black *Aspergilli*. Appl. Microbiol. Biotech. 50: 447-454.

R

Ramos, P.C., Höckendorff, J., Johnson, E.S., Varshavsky, A. and Dohmen, R.J. 1998. Ump1p is required for proper maturation of the 20S proteasome and becomes its substrate upon completion of the assembly. Cell 92: 489-499.

Raturi, A. and Mutus, B. 2007. Characterization of redox state and reductase activity of disulfide isomerase under different environments using a sensitive fluorescent assay. Free Radical Biol. Med. 43: 62-70.

Rautio, J., Smit, B.A., Wiebe, M., Penttilä, M. and Saloheimo, M. 2006. Transcriptional monitoring of steady state and effects of anaerobic phases in chemostat cultures of the filamentous fungus *Trichoderma reesei*. BMC Genomics 7: 247-262.

Rautio, J., Satokari, R., Vehmaan-Kreula, P., Serkkola, E. and Söderlund, H. 2008. TRAC in high-content gene expression analysis: application in microbial population studies, process biotechnology and biomedical research. Expert Rev. Mol. Diagn. 8: 379-385.

Rechsteiner, M. 1998. The proteasome. pp. 121-129. In Ubiquitin and biology of the cell. Eds. Peters, J-M., Harris, J.R. and Finley, D. New York. Plenum Press.

Regeimbal J. and Bardwell J.C.A. 2002. Disulfide Bond Formation in Prokaryotes and Eukaryotes, pp.132-149. In Protein Targeting, Transport & Translocation. Eds. Dalbey R.E. and von Heijne, G. Academic Press. London UK..

Reid, J. and Svejstrup, J.Q. 2004. DNA Damage-induced Def1-RNA Polymerase II Interaction and Def1 Requirement for Polymerase Ubiquitylation *in Vitro*. J. Biol. Chem. 279: 29875-29878.

Rekas, A., Alattia, J-R., Nagai, T. Miyawaki, A. and Ikura, M. 2002. Crystal structure of Venus, a yellow fluorescent protein with improved maturation and reduced environmental sensitivity. J. Biol. Chem. 277: 50573-50578.

Richie, D.L., Hartl, L., Aimaganianda, V., Winters, M.S., Fuller, K. K., Miley, M.D., White, S., McCarthy, J. W., Latgé, J-P., Feldmesser, M., Rhodes, J.C. and Askew, D.S. 2009. A role for the unfolded protein response (UPR) in virulence and antifungal susceptibility in *Aspergillus fumigatus*. PLoS Pathog. 5:e1000258. Epub 2009 Jan 9.

- Riordan, J.R. 1999. Cystic fibrosis as a disease of misprocessing of the cystic fibrosis transmembrane conductance regulator glycoprotein. *A. J. H. Genetics* 64: 1499-1504.
- Ritter, C. and Helenius, A. 2000. Recognition of local glycoprotein misfolding by the ER folding sensor UDP-glucose: glycoprotein glucosyltransferase. *Nat. Struct. Biol.* 7:278-280.
- Rocha, T. L., Paterson, G., Crimmins, K., Boyd, A., Sawyer, L. and Fothergill-Gilmore, L. A. 1996. Expression and secretion of recombinant ovine beta-lactoglobulin in *Saccharomyces cerevisiae* and *Kluyveromyces lactis*. *Biochem J.* 313: 927-932.
- Rock, K.L., Gramm, C., Rothstein, L., Clark, K., Stein, R., Dick, L., Hwang, D. and Goldberg, A.L. 1994. Inhibitors of the proteasome block the degradation of most cell proteins and the generation of peptides presented on MHC class I molecules. *Cell* 78: 761-771.
- Rockel, T.D., Stuhlmann, D. von Mickecz, A. 2005. Proteasome degrade proteins in focal subdomains of human cell nucleus. *J. Cell Sci.* 118: 5231-5242.
- Rogers, S., Wells, R., and Rechsteiner, M. 1986. Amino acids sequences common to rapidly degraded proteins: the PEST hypothesis. *Science* 234: 364-368.
- Rosenzweig, R. and Glickman, M.H. 2008. Chaperone-driven proteasome assembly. *Biochem. Soc. Trans.* 36: 807-812.
- Russell, S.J., Ruddock, L.W., Salo, K.E.H., Oliver, J.D., Roebuck, Q.P., Llewellyn, D.H., Roderick, H.L., Koivunen, P., Myllyharju, J. and High, S. 2004. The primary substrate binding site in the b' domain of ERp57 is adapted for endoplasmic reticulum lectin association. *J. Biol. Chem.* 279: 18861-18869.

S

- Saeki, Y., Toh-e, A., and Yokosawa, H. 2000. Rapid isolation and characterisation of the yeast proteasome regulatory complex. *Biochem. Biophys. Res. Commun.* 273: 509-515.
- Saeki, Y., Saitoh, A., Toh-e, A. and Yokosawa, H. 2002. Ubiquitin-like proteins and Rpn10 play cooperative role in ubiquitin-dependent proteolysis. *Biochem. Biophys. Res. Commun.* 293: 986-992.
- Sagt, C.M.J., Müller, W.H., van der Heide, L., Boonstra, J., Verkleij, A.J. and Verrips C.T. 2002. Impaired cutinase secretion in *Saccharomyces cerevisiae* induces irregular endoplasmic reticulum (ER) membrane proliferation, oxidative stress, and ER-associated degradation. *Appl. Env. Microbiol.* 68: 2155-2160.
- Saloheimo, M., Lund, M. and Penttilä, M.E. 1999. The protein disulfide isomerase gene of fungus *Trichoderma reesei* is induced by endoplasmic reticulum stress and regulated by carbon source. *Mol. Gen. Genet.* 262: 35-45.

- Saloheimo, M., Valkonen, M. and Penttilä, M. 2003. Activation mechanisms of the HAcI-mediated unfolded protein response in filamentous fungi. *Mol. Microbiol.* 47: 1149-61.
- Saloheimo, M., Wang, H., Valkonen, M., Vasara, T., Huuskonen, A., Riikonen, M., Pakula, T., Ward, M., and Penttilä, M., 2004. Characterization of secretory genes *ypt1/yptA* and *nsf1/nsfA* from two filamentous fungi: induction of secretory pathway genes of *Trichoderma reesei* under secretion stress conditions. *Appl. Env. Microbiol.* 70: 459-467.
- Sambrook, J. and Russel, D.W. 2001. *Molecular Cloning, a laboratory manual*. Gold Spring harbour Laboratory Press. New York.
- Scharf, A., Rockel, T.D. and von Mikecz, A. 2006. Localization of proteasomes and proteasomal proteolysis in mammalian interphase cell nucleus by systematic application of immunochemistry. *Histochem. Cell Biol.* 127: 591-601.
- Schawalder, S.B., Kabani M., Howald, I., Choudhury, U., Werner, M. and Shore, D. 2004. Growth-regulated recruitment of the essential yeast ribosomal protein gene activator Ifh1. *Nature* 432: 1058-1061.
- Scheffner, M., Smith, S. and Jentsch, S. 1998. The ubiquitin-conjugation system, pp.65-98. In: *Ubiquitin and the Biology of the Cell*. Peters, J.-M., Harris, J.R. and Finley, D. (Eds.), Plenum Press, New York, NY.
- Scheen, J. 1998. Mutational analysis of protein phosphatase 2C involved in abscisic acid signal transduction in higher plants. *Proc Natl. Acad Sci.* 95: 975-980.
- Schindler, M., Mach, R., Vollenhofer, S., Hodits, R., Gruber, F., de Graaff, J. and Kubicek, D. 1993. Characterization of the pyruvate kinase gene (*pk1*) of *Trichoderma reesei*. *Gene* 130: 271-275.
- Schrag, J.D., Propopio, D.O., Cygler, M., Thomas, D.Y. and Bergeron, J.J. 2003. Lectin control of protein folding and sorting in the secretory pathway. *Trends Biochem. Sci.* 28: 49-57.
- Schmidtke, G., Kraft, R., Kostka, S., Henklein, P., Frommel, C., Lowe, J., Huber, R., Kloetzel, P. M. and Schmidt, M. 1996. Analysis of mammalian 20S proteasome biogenesis: the maturation of beta- subunits is an ordered two-step mechanism involving autocatalysis. *EMBO J.* 15: 6887-6898.
- Schmidtke, G., Schmidt, M. and Kloetzel, P.M. 1997. Maturation of mammalian 20 S proteasome: purification and characterization of 13S and 16S proteasome precursor complexes. *J. Mol. Biol.* 268: 95-106.

- Schmitt, M., Neupert, W. and Langer, T. 1995. Hsp78, a Clp homologue within mitochondria, can substitute for chaperone functions of mt-hsp70. *EMBO J.* 14: 3434-3444.
- Schmidt, T., Chen, P.S. and Pellegrini, M. 1985. The induction of ribosome biosynthesis in a nonmitotic secretory tissue. *J. Biol. Chem.* 260: 7645-7650.
- Schmidt, M., Schmidtke, G. and Kloezel, P-M. 1997. Structure and structure formation of the 20S proteasome. *Mol. Biol. Reports.* 24: 103-112.
- Schröder, M. 2008. Endoplasmic reticulum stress responses. *Cell Mol Life Sci.* 65: 862-894.
- Schwartz, D.C. and Hochstrasser, M. 2003. A superfamily of protein tags: ubiquitin, SUMO and related modifiers. *Trends Biochem. Sci.* 28: 321-328.
- Schwarz, S.E., Rosa, J.L. and Scheffner, M. 1998. Characterization of human hect domain family members and their interaction with UbcH5 and UbcH7. *J. Biol. Chem.* Vol. 273: 12148-12154.
- Schuberth, C., Richly, H., Rumpf, S. and Buchberger, A. 2004. Shp1 and Ubx2 are adaptors of Cdc48 involved in ubiquitin-dependent protein degradation. *EMBO Rep.* 5: 818-824.
- Schuberth C. and Buchberger, A. 2005. Membrane-bound Ubx2 recruits Cdc48 to ubiquitin ligases and their substrates to ensure efficient ER-associated protein degradation. *Nat. Cell Biol.* 7: 999-1006.
- Schönbrunner, E. R. and Schmid, F. X. 1992. Peptidyl-prolyl cis-trans isomerase improves the efficiency of protein disulfide isomerase as a catalyst of protein folding. *Proc. Natl. Acad. Sci.* 89: 4510-4513.
- Scott, D.C. and Schekman R. 2008. Role of Sec61p in the ER-associated degradation of short-lived transmembrane proteins. *J. Cell Biol.* 181: 1095-1105.
- Seeger, M., Kraft, R., Ferrell, K., Bech-Otschir, D., Dumdey, R., Schade, R., Gordon, C., Naumann, M. and Dubiel, W. 1998. A novel protein complex involved in signal transduction possessing similarities to 26S proteasome subunits. *FASEB J.* 12: 469-478.
- Seemüller, E., Lupas, A., Stock D., Lowe, J., Huber, R., and Baumeister, W. 1995. Proteasome from *Thermoplasma acidophilum*: a threonine protease. *Science* 268: 579-582.
- Seidl, V., Gamauf, C., Druzhinina, I.S., Seiboth, B., Hartl, L. and Kubicek C.P. 2008. The *Hypocrea jecorina* (*Trichoderma reesei*) hypercellulolytic mutant Rut-C30 lacks a 85 kb (29 gene-encoding) region of the wild-type genome. *BMC Genomics* 9: 327-342.

- Sharon, M., Taverner, T., Ambroggio, X.I., Deshaies, R.J., Robinson, C.V. Structural organization of the 19S proteasome lid: Insights from MS of intact complexes. *PLoS Biol.* 4: 1314-1323.
- Shen, M.Y.J., Gropeanu, R., Pflugfelder, G.O., Weil, T. and Müllen, K. 2008. Fluorescent core/shell nanoparticles for specific cell-nucleus staining. *Small.* 4: 894-898.
- Sheikh, M.S. and Fornace Jr, A.J. 1999. Regulation of translation initiation following stress. *Oncogene* 18: 6121-6128.
- Shoji, J., Arioka, M. and Kitamoto K. 2008. Dissecting cellular components of the secretory pathway in filamentous fungi: insights into application for protein production. *Biotech. Lett.* 30: 7-14.
- Sidrauski, C., Brickner, J.H. and Walter, P. 2002 The unfolded protein response, pp.151-179. In *Protein Targeting, Transport and Translocation*. Eds. Dalbey, R.E. and von Heijne, G. Academic Press, London, UK.
- Sims, A.H., Gent, M.E., Lanthaler, K., Dunn-Coleman, N.S., Oliver, S.G. and Robson, G.D. 2005. Transcriptome analysis of recombinant protein secretion by *Aspergillus nidulans* and the unfolded-protein response *in vivo*. *Appl. Environ. Microbiol.* 71: 2737-2747.
- Sitia, R. and Braakman, I. 2003. Quality control in the endoplasmic reticulum protein factory. *Nature* 426: 891-894.
- Spiro, R. G. 2004. Role of N-linked polymannose oligosaccharides in targeting glycoproteins for endoplasmic reticulum-associated degradation. *Cell Mol. Life Sci.* 61: 1025-1041.
- Sprey, B. 1988. Cellular and extracellular localisation of endocellulase in *Trichoderma reesei*. *FEMS Microbiol. Lett.* 55: 283-294.
- St Leger, R.J., Staples, R.C. and Roberts, D.W. 1992. Cloning and regulatory analysis of starvation stress gene, *ssg4*, encoding a hydrophobin-like protein from the entomopathogenic fungus, *Metarhizium anisopliae*. *Gene* 120: 119-124.
- Staszczak, M. 2002. Proteasomal degradation pathways in *Trametes versicolor* and *Phlebia radiata*. *Enzyme Microb. Technol.* 30: 537-541.
- Steglich, G., Neupert, W. and Langer, T. 1999. Prohibitins regulate membrane protein degradation by the m-AAA protease in mitochondria. 1999. *Mol. Cell. Biol.* 19: 3435-3442.
- Stein, R.L., Melandri, F. and Dick, L. 1996. Kinetic characterization of the chymotryptic activity of the 20S proteasome. *Biochemistry* 35: 3899-3908.

Streit, S., Michalski, C.W., Erkan, M., Kleeff, J. and Friess, H. 2009. Northern blot analysis for detection and quantification of RNA in pancreatic cancer cells and tissues. *Nat. Protoc.* 4: 37-43.

Ståhlberg, J., Divne, C., Koivula, A., Piens, K., Claeysens, M., Teeri, T.T. and Jones, T.A. 1996. Activity studies and crystal structures of catalytically deficient mutants of cellobiohydrolase I from *Trichoderma reesei*. *J Mol Biol.* 264: 337-349.

T

Takeuchi, J. and Toh-e, A. 1999a. Genetic evidence for interaction between components of the yeast 26S proteasome: combination of a mutation in RPN12 (a lid component gene) with mutations in RPT1 (an ATPase gene) causes synthetic lethality. *Mol Gene Genet.* 262: 145-153.

Takeuchi, J., Fujimoro, M., Yokosawa, H., Tanaka, K. and Toh-E, A. 1999b. Rpn9 is required for efficient assembly of the yeast 26S proteasome. *Mol Cell Biol.* 19: 6575-6584.

Tansey, W.P. 2001. Transcriptional activation: risky business. *Genes Dev.* 15: 1045-1050.

Techel, D., Hafker, T., Muschner, S., Reimann, M., Li, Y., Monnerjahn, C. and Rensing, I. 1998. Molecular analysis of a glucose $\square$ regulated gene (*grp78*) of *Neurospora crassa*. *Biocem Biophys Acta.* 1397: 21-26.

Te'o, V.S.J., Cziferszky, A.E., Bergquist, P.L., and Nevalainen, K.M.H. 2000. Codon optimization of xylanase gene *xynB* from the thermophilic bacterium *Dictyoglomus thermophilum* for expression in the filamentous fungus *Trichoderma reesei*. *FEMS Microbiol Letters.* 190: 13-19.

Thrower, J.S., Hoffman, L., Rechsteiner, M. and Pickart, C.M. 2000. Recognition of the polyubiquitin proteolytic signal. *EMBO J.* 19: 94-102.

Tian, W., Zhang, L.V., Taşan, M., Gibbons, F.D., King, O.D., Park, J., Wunderlich, Z., Cherry, J.M. and Roth, F.P. 2008. Combining guilt-by-association and guilt-by $\square$ profiling to predict *Saccharomyces cerevisiae* gene function *Genome Biology*, 9: S7.

Tripler, C.P., Hutchon, S.P., Hendil, K., Tanaka, K., Fishel, S. and Mayer, R.J. 1997. Purification and characterisation of the 26S proteasomes from human and mouse spermatozoa, *Mol. Human Reprod.* 3: 1053-1060.

Tokumoto, M., Yamaguchi, A., Nagahama, Y. and Tokumoto, T. 2000. Identification of the goldfish 20S proteasome beta6 subunit bound to nuclear matrix. *FEBS Lett.* 472: 62-66.

- Tone, Y., Tanahashi, N., Tanaka, K., Fujimuro, M., Yokosawa H., and Toh-e A. 2000. Nob1p, a new essential protein, associates with the 26S proteasome of growing *Saccharomyces cerevisiae* cells. *Gene* 243: 37-45.
- Tone, Y. and Toh-E, A. 2002. Nob1p is required for biogenesis of the 26S proteasome and degraded upon its maturation in *Saccharomyces cerevisiae*. *Genes Dev.* 16: 3142-57.
- Travers, K.J., Patil, C.K., Wodicka, L., Lockhart, D.J., Weissman, J.S. and Walter, P. 2000. Functional and genomic analyses reveal an essential coordination between the unfolded protein response and ER-associated degradation. *Cell* 101: 249-258.
- Trombetta, E.S. and Parodi, A.J. 2003. Quality control and protein folding in the secretory pathway. *Annu. Rev. Cell Dev. Biol.* 19: 649-676.
- Tu, B.P. and Weissman, J.S. 2004. Oxidative protein folding in eukaryotes: Mechanisms and consequences. *J. Cell Biol.* 164: 341-346.
- Turano, C., Coppari, S., Altieri, F. and Ferrano, A. 2002. Proteins of PDI family: unpredicted non-ER locations and functions. *J. Cell Physiol.* 193: 154-163.
- Tyson J.R. and Stirling C.J. 2000. *LHS1* and *SIL1* provide a luminal function that is essential for protein translocation into the endoplasmic reticulum. *EMBO J.* 19: 6440-6452.

U

- Uchanska-Ziegler, B., Xugang, Q., Volz, A., Nissim, A. and Ziegler A. 2000. Human Single chain Fv fragments specific for natural killer cell receptors from phage display libraries. *Methods in Molecular Biology.* 121 pp. 219-290. In *Natural Killer Cell: Protocols Cellular and Molecular Methods*. Eds. Campbell, K.S. and Colonna, M. Humana Press Inc. Totowa. NJ.
- Udvardy, A. 1993. Purification and characterisation of a multiprotein component of the *Drosophila* 26S (1500 kDa) proteolytic complex. *J. Biol. Chem.* 268: 9055-9062.
- Ugai, S., Tamura, T., Tanahashi, N., Takai, S., Komi, N., Chung, C., Tanaka K. and Ichihara A. 1993. Purification and characterisation of the 26S proteasome complex catalysing ATP-dependent breakdown of ubiquitin-ligated proteins from rat liver. *J. Biochem.* 113: 754-768.
- Umebayashi, K., Hirata, A., Horiuchi, H., Ohta, A., and Takagi, M. 1999. Unfolded protein response-induced BiP/Kar2p production protects cell growth against accumulation of misfolded protein aggregates in the yeast endoplasmic reticulum. *Eur. J. Cell Biol.* 78: 726-738.
- Urano, T., Bertolotti, A. and Ron, D. 2000. IRE1 and effect signalling from endoplasmic reticulum. *J. Cell Sci.* 21: 3697-3702.

Uusitalo, J.M., Nevalainen, K.M.H., Harkki, A.M., Knowles, J.K. and Penttilä, M.E. 1991. Enzyme production by recombinant *Trichoderma reesei* strains. J Biotechnol. 17: 35-49.

V

Vainberg, I.E., Lewis, S.A., Rommelaere, H., Ampe, C., Vandekerckhove, J., Klein, H. L. and Cowan, N. J. 1998. Prefoldin, a chaperone that delivers unfolded proteins to cytosolic chaperonin. Cell 93: 863-873.

Valkonen, M. 2003. Functional studies of the secretory pathway of filamentous fungi. The effect of unfolded protein response on protein production. PhD thesis. VTT Publications 505. VTT. Espoo. Finland.

Valkonen, M., Penttilä, M., and Saloheimo, M. 2003a. Effects of inactivation and constitutive expression of the unfolded- protein response pathway on protein production in the yeast *Saccharomyces cerevisiae*. Appl. Env. Microbiol. 69: 2065-72.

Valkonen, M., Ward, M., Wang, H., Penttilä, M., and Saloheimo, M., 2003b. Improvement of foreign-protein production in *Aspergillus niger* var. *awamori* by constitutive induction of the unfolded-protein response. Appl. Env. Microbiol. 69: 6979-6986.

Valkonen, M., Penttilä, M. and Saloheimo, M., 2004. The *ire1* and *ptc2* genes involved in the unfolded protein response pathway in the filamentous fungus *Trichoderma reesei*. Mol. Genet. Genomics. 272: 443-451.

Valkonen, M., Kalkman, E.R., Saloheimo, M., Penttilä, M., Read, N.D. and Duncan, R.R. 2007. Spatially segregated SNARE protein interactions in living fungal cells. J. Biol Chem. 282: 22775-22785.

Vashist, S., Kim, W., Belden, W.J., Spear, E.D., Barlowe, C. and Ng, D.T. 2001. Distinct retrieval and retention mechanisms are required for the quality control of endoplasmic reticulum protein folding. J. Cell Biol. 155: 355-368.

Verdoes, J.C., Punt, P.J., Schrickx, J.M., Van Verseveld, H.W., Stouthamer, A.H. and van den Hondel, C.A.M.J.J. 1993. Glucoamylase overexpression in *Aspergillus niger*. Molecular genetic analysis of strains containing multiple copies of the *glaA* gene. Transgenic Res. 2: 84-92.

Verma, R., Chen, S., Feldman, R., Schieltz, D., Yates, J., Dohmen, J. and Deshaies, R.J. 2000. Proteasomal Proteomics: Identification of nucleotide-sensitive proteasome-interacting proteins by mass spectrometric analysis of affinity-purified proteasomes. Mol. Biol. Cell. 11: 3425-3439.

- Verma, R., Aravind, L., Oania, R., McDonald, W.H., Yates, I.J., Koonin, E.V. and Deshaies, R.J. 2002. Role of Rpn11 metalloprotease in deubiquitination and degradation by the 26S proteasome. *Science* 298: 611-615.
- Virgilio, M., Kitzmuller, C., Schweiger, E., Klein, M., Kreibich, G., Ivessa, N. E. 1999. Degradation of a short-lived glycoprotein from the lumen of the endoplasmic reticulum: the role of N-linked glycans and the unfolded protein response. *Mol. Biol. Cell.* 10: 4059-4073.
- Viswanathan, A., You, H.J. and Doetsch, P.W. 1999. Phenotypic change caused by transcriptional bypass of uracil in nondividing cells. *Science* 284: 159 □162.
- Voges, D., Zwicki, P. and Baumeister, W. 1999. The 26S proteasome: a molecular machine designed for controlled proteolysis. *Annu. Rev. Biochem.* 68: 1015-1068.
- W**
- Wagner, I., Arlt, H., van Dyck, L., Langer, T. and Neupert, W. 1994. Molecular chaperones cooperate with PIM1 protease in the degradation of misfolded proteins in mitochondria. *EMBO J.* 13: 5135-5145.
- Walz, J., Erdmann, A., Kania, M., Typke, D., Koster, A.J. and Baumeister, W. 1998. 26S proteasome structure revealed by three-dimensional electron microscopy. *J. Struct. Biol* 121: 19-29.
- Wang, C., Xi, J., Begley, T.P., and Nicholson, L.K. 2001. Solution structure of ThiS and implications for the evolutionary roots of ubiquitin. *Nat. Struct. Biol.* 8: 47-51.
- Wang, X. and Johnsson, N. 2005. Protein kinase CK2 phosphorylates Sec63p stimulate the assembly of the endoplasmic reticulum protein translocon apparatus. *J. Cell Sci.* 118: 723-732.
- Wang, T. and Hebert, D.N. 2003. EDEM and ER quality control receptor. *Nat. Structure Biol.* 5: 319-321.
- Wang, H. and Ward M. 2000. Molecular characterization of a PDI-related gene *prpA* in *Aspergillus niger* var. *awamori*. *Curr. Genet.* 37: 57-64.
- Warner, J.R. 1999. The economics of ribosome biosynthesis in yeast. *Trends Biochem. Sci.* 24: 437-440.
- Warner, J.R., Vilardell, J. and Sohn, J.H. 2001. Economics of ribosome biosynthesis. *Cold Spring Harbor Symp. Quant. Biol.* 66: 567-574.
- Wei, N. and Deng, X.W. 1998. Characterization and purification of the mammalian COP9 complex, a conserved nuclear regulator initially identified as a repressor of photomorphogenesis in higher plants. *Photochem. Photobiol.* 68: 237-241

- Wei, N. and Deng, X.W. 2003. The COP signalosome. *Annu. Rev. Cell Dev. Biol.* 19: 261-286.
- Weissman, A.M. 2001. Themes and variations on ubiquitylation. *Nat. Rev. Mol. Cell Biol.* 2: 69-178.
- Welchman, R.L., Gordon, C. and Mayer, J. 2005. Ubiquitin and ubiquitin-like proteins as multifunctional signals. *Nat. Rev. Mol. Cell Biol.* 6: 599-609.
- Welihinda, A.A., Tirasophon, W., Green, S.R. and Kaufman, R.J. 1998. Protein Serine/Threonine Phosphatase Ptc2p negatively regulates the unfolded-protein response by dephosphorylating Ire1p kinase. *Mol. Cell Biol.* 18: 1967-1977.
- Werner, E.D., Brodsky, J.L. and McCracken, A.A. 1996. Proteasome-dependent endoplasmic reticulum-associated protein degradation: an unconventional route to a familiar fate. *Proc. Natl. Acad. Sci. USA.* 93: 13797-13801.
- Werten, M.W., van den Bosch, T.J., Wind, R.D., Mooibroek, H. and de Wolf, F.A. 1999. High-yield secretion of recombinant gelatins by *Pichia pastoris*. *Yeast.* 15: 1087-1096.
- Wiebe, M.G., Rintala, E., Tamminen, A., Simolin, H., Salusjärvi, L., Toivari, M., Kokkonen, J.T., Kiuru, J., Ketola, R.A., Jouhten, P., Huuskonen, A., Maaheimo, H., Ruohonen, L. and Penttilä, M. 2008. Central carbon metabolism of *Saccharomyces cerevisiae* in anaerobic, oxygen limited and fully aerobic steady state conditions and following the shift to anaerobic conditions. *FEMS Yeast Res.* 8: 140-154
- Wiertz, E.J., Tortorella, D., Bogoy, M., Yu, J., Mothes, W., Jones, T.R., Rapoport, T.A. and Ploegh, H.L. 1996. Sec61-mediated transfer of a membrane protein from the endoplasmic reticulum to the proteasome for destruction. *Nature* 384: 432-438.
- Wing, S.S. 2003. Deubiquitinating enzymes-the importance of driving in reverse along the ubiquitin-proteasome pathway. *Int. J. Biochem. Cell Biol.* 35: 590-605.
- Wilkinson, K.D. 2000. Ubiquitination and deubiquitination: targeting of proteins for degradation by the proteasome. *Semin. Cell Dev. Biol.* 11:141-148.
- Williams, D.B. 2005. Beyond the lectins: the calnexin/calreticulin chaperone system of the endoplasmic reticulum. *J. Cell Sci.* 119: 615-623.
- Wolf, D.H. and Plemper, R.K. 1999. Endoplasmic reticulum degradation. Reverse protein transport and its end in the proteasome. *Mol. Biol. Reports.* 26: 125-130.
- Wolf, D.H. and Hilt, W. 2004. The proteasome: a proteolytic nanomachine of cell regulation and waste disposal. *Biochim. Biophys. Acta.* 1695: 19-31.

Wong, K.H., Todd ,R.B., Oakley, B.R., Oakley, C.E., Hynes, M.J. and Davis, M.A. 2008. Sumoylation in *Aspergillus nidulans*: *sumO* inactivation, overexpression and live-cell imaging. Fungal Gen. Biol. 45: 728-737.

X

Xu, P. and Peng, J. 2008. Characterization of polyubiquitin chain structure by middle-down mass spectrometry. Anal. Chem. 80: 3438-3444.

Y

Yao, T. and Cohen, R.E. 2002. A cryptic protease couples deubiquitylation and degradation by the proteasome. Nature 419: 403-407.

Yanagawa, Y., Ohhashi, A., Murakami, Y., Saeki, Y., Yokosawa, H., Tanaka, K., Hashimoto, J., Sato, T. and Nakagawa, H. 1999. Purification and characterization of the 26S proteasome from cultured rice (*Oryza sativa*) cells. Plant Sci. 149: 13-41.

Ye, L. and Pan, L. 2008. A comparison of the unfolded protein response in solid-state with submerged cultures of *Aspergillus oryzae*. Biosci. Biotechnol. Biochem. 72: 2998-3001.

Yen, H-C.S., Espiritu C. and Chang, E.C. 2003. Rpn5 is a conserved proteasome subunit and required for proper proteasome localization and assembly. J. Biol. Chem. 278: 30669-30676.

Yerlikaya, A. 2004. Cellular functions of the 26S proteasome. Turkish J. Biol. 28:31-38.

Yoder, W.T. and Lehmbeck, J. 2004. Heterologous expression and protein secretion in filamentous fungi. In Advantages in fungal biotechnology for industry, agriculture and medicine. Eds. Tkacz, J.S. and Leng, L. Springer. NY. USA.

Yoshida, H., Matsu, I.T., Yamamoto, A., Okada, T. and Mori, K. 2001. XBP1 mRNA is induced by ATF6 and spliced by IRE1 in response to ER stress to produce a highly active transcription factor. Cell 107: 881-891.

Z

Zhang, J., Campbell, R.E., Ting, A.Y. and Tsien, R.Y. 2002. Creating new fluorescent proteins for cell biology. Nat. Rev. Mol. Cell Biol. 3: 906-918.

Zhang, K. and Kaufman, R.J. 2004 Signaling the unfolded protein response from the endoplasmic reticulum. J. Biol. Chem. 279: 25935-25938.

Zhao, Y., Sohn, J.H. and Warner, J.R. 2003. Auto regulation in the biosynthesis of ribosomes. Mol. Cell Biol. 23: 699-707.

- Zhao, A. and Rodgers, V.G.J. 2006. Using TEM to couple transient protein distribution and release for PLGA microparticles for potential use as vaccine delivery vehicles. *J. Contr. Rel.* 113: 15-20.
- Zhou, S., and DeWille, J. 2007. Proteasome-mediated CCAAT/enhancer-binding protein δ (C/EBP δ) degradation is ubiquitin-independent. *Biochem. J.* 405: 341-349.
- Zwickl, P., Grziwa, A., Puhler, G., Dahlmann, B., Lottspeich, F. and Baumeister, W. 1992. Primary structure of the *Thermoplasma* proteasome and its implications for the structure, function, and evolution of the multicatalytic proteinase. *Biochem.* 31: 964-972.