

# **Verbs and nouns in awake neurosurgery needs and answers**

by

Adrià Rofes

A Doctoral thesis submitted in partial fulfillment of the requirements  
for the degree of Doctor in Philosophy

at the International Doctorate for Experimental Approaches to Language and Brain  
(IDEALAB)

October, 2015



Under the supervision of

Prof. Dr. Gabriele Miceli, University of Trento

Prof. Dr. Roelien Bastiaanse, University of Groningen

Prof. Dr. Lyndsey Nickels, Macquarie University

Assessment committee

Prof. Dr. Antoni Rodríguez Fornells, University of Barcelona

Prof. Dr. Jenny Crinion, University College London

Prof. Dr. Roberto Zamparelli, University of Trento



UNIVERSITY  
OF TRENTO



university of  
 groningen



MACQUARIE  
University



Newcastle  
University



Universität  
Potsdam



## TABLE OF CONTENTS

ACKNOWLEDGEMENTS.....	ix
LIST OF TABLES.....	xi
LIST OF FIGURES.....	xiii
Chapter 1 - General Introduction.....	1
Chapter 2 – Language mapping with verbs and sentences in awake surgery: A Review ....	5
2.1. Introduction.....	6
2.1.1. Language mapping tests in awake surgery.....	9
2.1.2. Why use verbs in intraoperative language mapping?.....	11
2.2. Methods.....	13
2.3. Results.....	14
2.3.1. Tasks with verbs and sentences.....	14
2.3.2. Language processes assessed by each task.....	17
2.2.3. Areas where cortico-subcortical stimulation inhibited language processing and types of errors elicited by stimulation.....	20
2.4. Discussion.....	23
2.4.1. Towards a differentiated approach to language mapping.....	24
2.4.2. Design of intraoperative tasks.....	26
2.4.3. Relevance of standardized and customizable tasks.....	28
2.4.4. Other neurocognitive components of intraoperative language mapping.....	29
2.5. Conclusion.....	31
Chapter 3 - A minimal standardization setting for language mapping tests:	
An Italian example.....	33
3.1. Introduction.....	34
3.2. VISC (verb production in sentence context).....	39
3.2.1. Picture-name agreement and cut-offs.....	39
3.2.2. Norming of the stimuli.....	40
3.3. ECCO (object naming test).....	41
3.3.1. Picture-name agreement and cut-offs.....	42
3.3.2. Norming of the stimuli.....	42
3.4. Validation on a clinical population.....	43
3.4.1. Participants.....	43
3.4.2. Materials and procedure.....	43
3.4.3. Analyses and results.....	44
3.4.4. Example of data usage in a single-case setting using hypothetical scenarios...	45
3.5. Discussion.....	46
Chapter 4 - Verb production tasks in the measurement of communicative abilities in aphasia.....	49
4.1. Introduction.....	50
4.1.1. Differences between verbs and nouns.....	51
4.1.2. Language properties involved by picture-naming paradigms.....	54

4.1.3. Impairment-level and functional-level measures.....	55
4.1.4. Questions and hypotheses.....	57
4.2. Methods.....	59
4.2.1. Participants.....	59
4.2.2. Materials.....	60
4.2.2.1. Picture naming tasks.....	60
4.2.2.2. Functional measures of language.....	62
4.2.3. Procedure and scoring.....	63
4.3. Results.....	64
4.4. Discussion.....	66
4.4.1. An advantage for producing a finite verb in sentence context over other tasks?.....	67
4.4.2. Both object and action naming tasks may be relevant in clinical practice.....	70
4.4.3. CADL-2, CETI, and CAL as measures of communicative abilities in aphasia.....	72
4.4.4. Relevance and limitations of the study.....	73
4.5. Conclusion.....	75
Chapter 5 - Advantages and disadvantages of intraoperative language tasks in awake surgery.....	77
5.1. Introduction.....	78
5.1.1. Language testing in awake surgery.....	79
5.1.2. Models of language processing.....	80
5.1.3. Advantages and disadvantages of tests reported in the literature.....	82
5.1.3.1. Non-semantic processes.....	83
5.1.3.2. Lexico-semantic processes.....	85
5.1.3.3. Grammatical processes.....	91
5.1.4. Administration of tasks during surgery.....	95
5.2. A three-task approach for prefrontal tumors.....	97
5.2.1. Participants.....	97
5.2.2. Materials and procedure.....	98
5.2.3. Results.....	99
5.3. Discussion: future steps of language tests in awake surgery.....	101
5.4. Conclusion.....	103
Chapter 6 - Surgical mapping of language production in sentence context	
A single case series.....	105
6.1. Introduction.....	106
6.1.1. Aims and predictions.....	112
6.2. Methods.....	113
6.2.2. Participants.....	113
6.2.3. Materials.....	113
6.2.4. Procedure.....	115
6.3. Results.....	116
6.3.1. Quantitative results.....	116
6.3.2. Error analyses.....	121

6.4. Discussion.....	124
6.4.1. DES v no-DES (collapsing across nouns and verbs).....	124
6.4.2. Object naming and finite verbs (independently of stimulation site).....	125
6.4.3. Object naming and finite verbs (by stimulation site).....	127
6.4.4. Error analyses.....	129
6.5. Conclusion.....	131
 Chapter 7 - General discussion.....	 133
8.1. Applying language knowledge to functional neurosurgery.....	136
8.2. Developing new and standardized tasks in awake surgery.....	148
8.3. The production of finite verbs (in awake surgery).....	141
8.4. Conclusion.....	144
 REFERENCES.....	 145
 APPENDICES.....	 181
APPENDIX A: Cortico-subcortical areas where language processing was detected and error types during electrical stimulation mapping (Chapter 2).....	181
APPENDIX B Description of psycholinguistic variables of the tests, instructions for on-line questionnaires, and MATLAB code to calculate the H-STATISTIC (Chapter 3).....	184
APPENDIX C: Matching for linguistic variables, post-hoc tests, and individual results (Chapter 4).....	198
 EPILOGUE.....	 203
ABOUT THE AUTHOR.....	203
PUBLICATIONS, PUBLISHED ABSTRACTS, AWARDS.....	204





## ACKNOWLEDGEMENTS

I express my deepest gratitude to Prof. Dr. Gabriele Miceli for his enthusiasm and honest support on this project, for his networking knack, and for the finest advice and revisions - working with you provided me strong means to become an independent researcher, with robust problem-solving capacities, and with resilient aptitude to work in group-based environments. I sincerely thank Prof. Dr. Roelien Bastiaanse for supporting my application to this program, providing an excellent continuation of my career, and fuelling my interest in the academic conversation - neither verbs or clinical linguistics would have passed through my mind without your input. I am indebted to Prof. Dr. Lyndsey Nickels for the clearest feedback, organizational skills, and extensive support on single case methodology and cognitive neuropsychology - I will certainly keep all that knowledge in a good place.

I will not forget working with Dr. Barbara Santini - thank you for paving the way to join forces with the Hospital Borgo Trento in Verona, and for being critically open to new assessment methods. I thank Prof. Andrea Talacchi and Prof. Giampietro Pinna, for their work in the operation room, and Dr. Pasquale Mirtuono, Dr. Giuseppe Kenneth Ricciardi, Dr. Giada Zocatelli, and Dr. Franco Alessandrini, for all the radiological insights. I feel very rewarded to have worked with Dr. Giannantonio Spena - you have always received me with open arms and have been very available. I am grateful to Dr. Antonio Miozzo for his efficiency following patients at the Hospital in Brescia, to Prof. Sergio Carlomagno for helping me with functional language assessment materials in Italian, and to Dr. Rita Capasso for following patients in Rome and welcoming me in Rovereto with a huge booklet of articles! I will not forget the very valuable opportunities that Dr. Silvio Sarubbo and Dr. Monica Dallabona brought me at the Hospital Santa Chiara in Trento. I am also very thankful to Dr. Emmanuel Mandonnet and Prof. Stephney Whillier to allow me to sit in their neuroanatomy and brain dissection courses at the Lariboisière Hospital in Paris and at Macquarie University in Sydney. To Prof. Dr. Angelika Lingnau for following my intern work on fMRI data acquisition and analysis - it was certainly a pleasure to learn from you. I am grateful to Prof. Dr. Gabriella Vigliocco and to Prof. Dr. Judit Druks for welcoming me at their labs in London and to discuss parts of this thesis with me. I am indebted to Prof. Marco Baroni, Dr. Alberto Abad, Germán Kruszewski, and Vânia de Aguiar for passing on useful computational programming skills, and for fuelling the scientific discussion. I also thank Giovanna Cappelletti for being an excellent master's student - *110 e lode!*

I thank the academic staff of the consortium of universities from the International Doctorate for Experimental Approaches to Language and Brain (IDEALAB), all the scholars and colleagues that were interested in this work and who have put me to test in conferences and colloquia, and the administrative personnel in Trento, Groningen, and Macquarie; in particular, Anja Papke, Ulla Behr, Lesley McKnight, Alice Poomstra, and Leah Mercanti. Last but not least, I thank all my friends and family - life would have been much more difficult without you!

This work has been funded by the Erasmus Mundus PhD Program IDEALAB (Macquarie University, Newcastle University, University of Groningen, University of Trento and University of Potsdam: agreement number 2014-0025).



## LIST OF TABLES

Table 2.1 Language processes assessed by each task.....	19
Table 3.1 Age-based cut-off scores VISC and ECCO.....	40
Table 3.2 Convergent and divergent validation: summary of results.....	44
Table 4.1 Language processes assessed by each task.....	55
Table 4.2 Main demographic and neurological features of participants.....	59
Table 5.1 Language processes assessed by each task.....	83
Table 6.1 Subject characteristics and tumor variables.....	113
Table 6.2 Preoperative differences in psycholinguistic dimensions (subsets).....	116
Table 6.3 Individual comparisons per condition.....	120
Table A2.1 Cortical areas in language dominant hemisphere where language processing was detected with each task and error types per area and per test.....	181
Table A2.2 Subcortical areas in the language-dominant hemisphere where language processing was detected by action naming and error types produced by patients during stimulation of that area.....	183
Table C4.1 Matching of the tests for relevant linguistic variables including imageability...	198
Table C4.2 Individual behavioral data (percentage correct).....	199
Table C4.3 Post-hoc tests of picture-naming tasks (Z and FDR corrected p-values).....	200
Table C4.4 Spearman correlations of picture-naming tasks and functional-measures (rho and p-value).....	200
Table C4.5 Comparisons of overlapping correlations (Steiger's Z).....	201
Table C4.6 Paired tests on picture-naming tasks (Fisher Exact – two tailed).....	201
Table C4.7 Major error types per task and participant.....	202



## LIST OF FIGURES

Figure 2.1. Greater implication of the left inferior frontal regions in verb processing.....	13
Figure 2.2. Cortical and subcortical structures in which DES interfered with language processing and error types.....	22
Figure 3.1. The use of picture naming tasks in awake surgery.....	35
Figure 3.2. Examples of picture stimuli.....	36
Figure 3.3. The indicative value of psycholinguistic variables detecting damage to language and the brain .....	38
Figure 4.1.Examples of items from each of the tasks structure of experimental trial.....	63
Figure 4.2. Spearman correlations between CADL2 and picture-naming tasks.....	69
Figure 4.3. Performance patterns in individual participants.....	71
Figure 5.1. Schematic representation of language components.....	81
Figure 5.2. Summary of language maps.....	99
Figure 6.1. Significant DES sites per subject.....	117
Figure 6.2. Error types during DES per subject.....	121



*To my dearest wife*





# Chapter 1

## General introduction

Awake neurosurgery (henceforth, awake surgery) is a medical procedure recommended for patients with brain tumors or pharmacologically intractable epilepsy. It borrows its name from the intraoperative stage in which patients are awake, so that neuropsychological and/or language tasks can be administered (e.g., Berger 1996; Duffau et al. 1999; Huncke, Van de Wiele, Itzhak, & Rubinstein, 1998; Kilbride, 2013; Ojemann & Mateer, 1979; Penfield & Boldrey, 1937). Awake surgery has gained popularity over the last years, because in more classical approaches - where patients are not awake and language tests cannot be administered - less tumor tissue may be removed and worse cognitive outcomes may be obtained (see, for a meta-analysis, De Witt Hamer, Gil-Robles, Zwinderman, Duffau, & Berger, 2012). Language testing is usually also performed shortly before and after surgery (the so-called “perioperative stages”). Awake surgery finds its way within a multidisciplinary field that includes a range of specialists – radiologist, anesthesiologist, neurosurgeon, neuropathologist, neurophysiologist and neuropsychologist (or speech/language therapist or clinical linguist). All these professionals have a specific role in the assessment and treatment of each individual patient. As researchers, we should strive to understand the needs of these professionals, and provide answers that help them reach their goals, while enabling us to gain knowledge on language and the brain, and to improve current standards of patient care.

A recurrent topic throughout this dissertation is that the language capacities of patients with brain tumors have been traditionally assessed with few tasks, and sometimes with non-standardized materials (e.g., Finch & Copland, 2014; Rofes & Miceli, 2014). Object naming tasks have been a key component of these assessments (for a review see De Witte & Mariën,

2013). Object naming has much to commend it, as seeing a picture and producing a noun (“apple”) or a determiner phrase (“the apple”) engages lexico-semantic input and output processes that are indispensable for everyday communication (Goodglass & Wingfield, 1997). For example, perioperative processing speed on object naming has been shown to correlate with postsurgical chances of return to work in patients undergoing this type of surgery (Moritz-Gasser, Herbet, Maldonado, & Duffau, 2012). Furthermore, object naming tasks were introduced by pioneers in the field and have maintained their role as a gold standard in the identification of areas relevant for language processing during surgery (Hamberger & Tamny, 1999; Ojemann & Mateer, 1979; Ojemann, Ojemann, & Lettich, 2002; Sanai & Berger, 2009).

Despite these positive results, having object naming as the main component of assessment in awake surgery may not be always sufficient. This is because (1) the language capacities required for object naming may be relatively spared in spite of damage to other language abilities such as reading, writing, comprehension, or naming of actions (Santini, Talacchi, Squintani, Casagrande, Capasso, & Miceli, 2012; Satoer et al., 2014); (2) there are language processes typically used in everyday language (e.g., for sentence formulation, to refer to actions, or to moments in time) that cannot be assessed with object naming (e.g., Rofes & Miceli, 2014; for a review); and (3) the neural correlates of object naming may be partially segregated from those of other processes such as action-naming (e.g., Mätzig, Druks, Masterson, & Vigliocco, 2009; Vigliocco, Vinson, Druks, Barber, & Cappa, 2011; for reviews). These are important issues in aphasiology, neurolinguistics, and cognitive neuroscience, which we revisited to work on critical improvements for awake surgery: the use of standardized tasks for language mapping, more specifically, the use of verbs as isolated words and finite verbs in sentences.

Verbs differ from nouns in a number of ways: they typically refer to actions (Gentner, 1982), and may weigh differently on psycholinguistic variables that affect performance in individuals with aphasia and non-brain-damaged people (e.g., Whitworth, Webster, & Howard, 2005; Vigliocco, Vinson, Druks, Barber, & Cappa, 2011). Verbs also entail a predicate-argument and a thematic structure (Carlson & Tanenhaus, 1989), and when they are used in the context of a sentence they require that agreement relations between the subject and the verb be fulfilled (Hale & Keyser, 1998). Verbs also allow the positioning of events in time and reference to events within a specific time frame (Bastiaanse, Bamyaci, Hsu, Lee, Yarbay Duman, & Thompson, 2011). Furthermore, they may be processed in different brain regions compared to nouns. It has been shown that verb processing elicits activation in the frontal and inferior-parietal lobe, whereas nouns trigger more activation in the temporal lobe (e.g., subjects KJ-1360, AN-1033 and Boswell in Damasio & Tranel, 1993; Lubrano, Filleron, Démonet, & Roux, 2014; Miceli, Silveri, Noncentini, & Caramazza, 1988).

Overall, earlier research suggests that the current assessments in awake surgery are pertinent but not sufficient. In this dissertation, we contribute to the efforts that are underway to meet the needs of awake surgery, both in terms of behavioral and anatomical knowledge. We approach the issue from the perspective of aphasiology, neurolinguistics and cognitive neuroscience (e.g., Coltheart 2011; Withworth et al., 2005). The chapters of this thesis each correspond to an article published in an international peer-reviewed journal or to a final draft that is (or will be) submitted for publication.

The thesis begins with a review of language mapping with tests that use verbs and sentences (Chapter 2). This chapter serves as a critical introduction to some of the needs that neurosurgeons and people responsible for the assessment of language skills in individuals with

brain gliomas face in everyday practice. More specifically, it talks about renewing the consideration of tasks that use verbs, an approach already tackled in classical papers such as Ojemann and Mateer (1979). After this introduction, we deal with the design and development of an object naming task and a task with finite verbs in sentences, specifically designed to assess Italian-speakers undergoing awake surgery (Chapter 3). Such tools did not exist for the Italian language. The standardization of the tasks is described so that both the task methodology and the standardization of the tasks can be easily adopted by other research or awake surgery teams.

In Chapter 4, the validity of the noun and verb picture-naming tasks are assessed, as well as two other tasks typically used to assess language deficits in communicative contexts. By evaluating the performance of people suffering from post-stroke aphasia, we assess how well these naming tasks correlate with the ability to use language in an everyday context. The next two chapters describe the implementation of these tasks in surgical rooms and report the neurofunctional results of perioperative tests, their relation to the current neuroscience of language literature, and a critical review of the advantages and disadvantages that language mapping tasks may entail for awake surgery (Chapter 5 and Chapter 6).

We finish the dissertation with a general discussion, aimed at describing theoretical and clinical reasons why a rigorous approach may improve awake surgery procedures. We also discuss some future implications of this line of work (Chapter 7).

**Chapters 2 & 3 (pp. 5-48) of this thesis have been removed as they contain published material under copyright. Removed contents published as:**

Rofes, A., Miceli, G. (2014) Language Mapping with Verbs and Sentences in Awake Surgery: A Review. *Neuropsychology Review*, vol. 24, pp. 185–199.  
<https://doi.org/10.1007/s11065-014-9258-5>

Rofes, A., de Aguiar, V. & Miceli, G. (2015) A minimal standardization setting for language mapping tests: an Italian example. *Neurol Sci* 36, 1113–1119 (2015). <https://doi.org/10.1007/s10072-015-2192-3>

# Chapter 4

## Verb production tasks in the measurement of communicative abilities in aphasia<sup>4</sup>

### Abstract

*Background:* The neurofunctional correlates of verbs and nouns have been the focus of many theoretically oriented studies. In clinical practice, however, more attention is typically paid to nouns, and the relative usefulness of tasks probing nouns and verbs is unclear. The routine administration of tasks that use verbs could be a relevant addition to current batteries. Evaluating performance on both noun and verb tasks may provide a more reliable account of everyday language abilities than an evaluation restricted to nouns.

*Aims:* To assess the benefits of administering verb tasks in addition to noun tasks, and their relation to three functional measures of language.

*Method and procedure:* Twenty-one subjects with poststroke language disorders completed four picture-naming tasks and a role-playing test (Communicative Abilities in Daily Living, Second Edition, CADL-2), commonly used as measure of everyday language abilities. Two questionnaires (Communicative Effectiveness Index, CETI, and Communicative Activity Log, CAL) were completed by caregivers. Picture-naming tasks were matched for psycholinguistic variables to avoid lexicosemantic and morphosyntactic confounds.

*Results:* No significant differences emerged across picture-naming tasks. Scores on the role-playing test and the two questionnaires differed; scores between the two questionnaires did not. The four naming tasks correlated significantly with CADL-2, CETI, and CAL. The strength of the correlation with CADL-2 was significantly greater for Producing a finite verb in sentence context than for Object Naming. Thirteen participants showed no differences in performance between tasks, 6 fared significantly worse on verb tasks than on Object Naming, 1 fared better at Producing a finite verb in sentence context though his performance was poor overall, and 1 was significantly more impaired on verbs.

*Conclusions:* Performance on tasks that use verbs, and especially Producing a finite verb in sentence context, may provide a more accurate estimate of language abilities in daily living than Object Naming alone. Administering both verb and noun tasks may be recommended in clinical practice.

---

<sup>4</sup> Published manuscript. Rofes, A., Capasso, R., & Miceli, G. (in press). Verb production tasks in the measurement of communicative abilities in aphasia. *Journal of Clinical and Experimental Neuropsychology*, 37(5), 483-502. doi:10.1080/13803395.2015.1025709

#### **4.1. Introduction**

Aphasic disorders can affect the subject's ability to communicate, to various extents and in various forms. A well-known phenomenon in aphasic speech is the separability of noun and verb processing (for reviews, see Black & Chiat, 2003; Damasio & Tranel, 1993; Kiefer & Pulvermüller, 2012; Mätzig, Druks, Masterson, & Vigliocco, 2009; Pulvermüller, 2005; Shapiro & Caramazza, 2003b; Vigliocco, Vinson, Druks, Barber, & Cappa, 2011). Such distinction is the object of a growing and controversial body of neuropsychological investigations, trying to establish the neurofunctional correlates of the two word types. Studies have reached contrasting conclusions, and the patterns of associations/dissociations documented in language-impaired individuals have been attributed to lexical variables (e.g., Shapiro & Caramazza, 2003b), to supramodal semantic variables (e.g., Bird, Howard, & Franklin, 2000), or to sensorimotor semantic properties (e.g., Pulvermüller, 2005). Regardless of where they locate the critical distinctions, however, these experimental studies have devoted comparable attention to nouns and verbs and have contributed to a better understanding of the representation/processing of nouns and verbs, and of their neural underpinnings (e.g., Damasio & Tranel, 1993).

By contrast, in clinical practice object naming is part-and-parcel of most aphasia screening tests (e.g., Goodglass & Kaplan, 1983; Huber, Poeck, & Willmes, 1983; Kertesz, 1982; Swinburn, Porter, & Howard, 2004; Riddoch & Humphreys, 1993) but relatively little attention has been devoted to verbs (Bastiaanse, Edwards, Maas, & Rispens, 2003; Conroy, Sage, & Lambon Ralph, 2009; de Aguiar, Paolazzi, & Miceli, 2015; Rofes & Miceli, 2014). As a consequence, to date information on the relative usefulness of noun and verb production tasks in assessing communicative abilities in everyday life is still scarce. To evaluate the benefits of systematically administering verb tasks in clinical evaluations, especially when measuring



communicative difficulties in everyday life, we studied the performance of people with aphasia on four picture-naming tests that use verbs and nouns (Object Naming, Verb Generation, Producing a finite verb in sentence context, and Action Naming) and the relation of such tasks with three functional measures of language—a role-playing test administered to the patient, and two questionnaires filled by the caregiver.

#### **4.1.1. Differences between verbs and nouns**

Verbs and nouns contain many lexicosemantic variables that must be controlled to avoid unwanted confounds (Kemmerer & Tranel, 2000; Whitworth, Webster, & Howard, 2005; for a review). For example, verbs used in experimental designs typically refer to actions and are less referential than nouns, which refer to objects (Gentner, 1982). The different distributions of nouns and verbs as regards frequency and imageability values have also been the source of a recurrent argument in the neuropsychological literature (Bird et al., 2000; Luzzatti et al., 2002; for contrasting results see Berndt, Haendiges, Burton, & Mitchum, 2002; Black & Chiat, 2003). When nouns and verbs are studied in the context of language disorders, further problems may arise. In the case of nouns, an additional difficulty may stem from the overlap of deficits for the grammatical category “noun” with semantic category-specific effects, selectively affecting or sparing conspecifics, animals, plant life, or tools (Damasio, Grabowski, Tranel, Hichwa, & Damasio, 1996; Hart, Berndt, & Caramazza, 1985; Warrington & McCarthy, 1983; Warrington & Shallice, 1984). In the case of verbs, difficulties may arise from diverse syntactic structure complexities (transitive vs. intransitive verbs) or from similarity to a noun (Jonkers & Bastiaanse, 2007; Thompson, 2003). Additional issues may arise with verbs denoting actions performed with specific body parts (Finocchiaro & Miceli, 2002; Pulvermüller, 2005), or with

nouns referring to a manipulable object (Aggujaro, Crepaldi, Pistarini, Taricco, & Luzzatti, 2006; Bub & Masson, 2012; Rueschemeyer, van Rooij, Lindemann, Willems, & Bekkering, 2010). Number of phonemes, stress patterns (Black & Chiat, 2003; Nickels & Howard, 2004), and age of acquisition (Bell, Davies, Hermann, & Walters, 2000; Morrison, Ellis, & Quinlan, 1992) may also play a relevant role.

Verbs and nouns also differ along morphosyntactic dimensions. Verbs are the main component of the predicate: They entail a complex argument structure (Thompson, 2003), thematic role assignment (Carlson & Tanenhaus, 1989), subject–verb agreement (Hale & Keyser, 1998), and other processes relevant to communication, such as time reference (Berndt, Haendiges, Mitchum, & Sandson, 1997). By contrast, nouns typically function as arguments of the predicate: They complete its meaning by filling in the grammatical requirements of the verb (e.g., subject/agent, object/theme, locative/goal). For example, in order to convey that “someone” (“the boys”) finished the action of making “something” (“the car”) clean, we may use the sentence “The boys have washed the car,” where the verb “have washed” indicates what happened (verb meaning) and when it happened (time reference), and tells who performed the action and to whom or what (reference to person, number, argument structure, thematic roles). The nouns “boys” and “car” indicate the agent and theme of the action, as projected by the grammatical properties of the verb. Another potentially critical difference between nouns and verbs is that different languages may take very diverse sets of verbal and nominal inflections. For example, English verbs have four possible inflections, but English nouns only have two. Italian verbs have 46 inflected forms, and nouns two, or in rare cases four (e.g., *bambino*, *bambini*, *bambina*, *bambine*; boy, boys, girl, girls). The opposite is the case in Chinese, where verb morphology is less complex than noun morphology (Bastiaanse et al., 2011; Tsapkini, Jarema, &

Kehayia, 2002). Other morphosyntactic variables with potential impact on performance are the number of internal arguments (Bastiaanse & Van Zonneveld, 2004; Thompson, Lange, Schneider, & Shapiro, 1997), transitivity (De Bleser & Kauschke, 2003; Luzzatti et al., 2002), and regularity (Morrison et al., 1992; Pinker & Prince, 1994; Sach, Seitz, & Indefrey, 2004).

Verb and noun differences have behavioral, neuroanatomical, and neuroimaging correlates. Verbs are associated with longer reaction times than nouns in psycholinguistic paradigms, including picture naming (Arévalo, 2002; Székely et al., 2005; cf. Tsigka, Papadelis, Braun, & Miceli, 2014). People with aphasia frequently fare worse at production tasks that use verbs than at tasks that use nouns (Bastiaanse & Jonkers, 1998; Kemmerer, Rudrauf, Manzel & Tranel, 2012; Luzzatti et al., 2002; Mätzig et al., 2009; Tranel, Adolphs, Damasio, & Damasio, 2001; Vigliocco et al., 2011), although the opposite pattern has also been documented (e.g., Miceli, Silveri, Nocentini, & Caramazza, 1988; Miceli, Silveri, Villa, & Caramazza, 1984; Damasio & Tranel, 1993; Goldberg & Goldfarb, 2005). Dissociations between verbs and nouns raised the question of whether these word categories may be represented in at least partly separable neural networks. In neuroimaging studies, large sources of activity in the left inferior frontal gyrus have been detected in processing unambiguous verbs as compared to nouns (Federmeier, Segal, Lombrozo, & Kutas, 2000) and inflectional verb affixes as compared to noun affixes (Den Ouden, Fix, Parrish, & Thompson, 2009; Finocchiaro, Basso, Giovenzana, & Caramazza, 2010; Pulvermüller & Shtyrov, 2009; Tsigka et al., 2014; Tyler, Bright, Fletcher, & Stamatakis, 2004; Yokohama et al., 2006). A greater involvement of frontal regions for verbs than nouns has also been reported in several brain-damaged populations, including stroke, frontotemporal dementia, Alzheimer's disease, HIV-1 infection, brain tumors, and epilepsy (Conner, Chen, Pieters, & Tandon, 2014; Corina et al., 2005; Damasio & Tranel, 1993; Hillis &

Caramazza, 1995; Kemmerer et al., 2012; Kemmerer & Tranel, 2000; Lubrano, Filleron, Démonet & Roux, 2014; Miceli, Silveri, Romani, & Caramazza, 1989; Shapiro & Caramazza, 2003a, 2003b; Tranel et al., 2001; Woods, Carey, Tröster, & Grant, 2005).

#### **4.1.2. Language properties involved by picture-naming paradigms**

Response accuracy on a given task can be differentially affected by many variables, including the types of knowledge (semantic vs. lexical) tapped by the stimuli, the lexicosemantic and morphosyntactic characteristics of the target words, and whether the target word appears in isolation or in sentence context (Nickels & Howard, 1995). The picture-naming tasks used in the present study require participants to respond to drawings of actions by producing the name of the action with a verb in the infinitive form (i.e., Action Naming), or with a finite verb in the third person singular (i.e., Producing a finite verb in sentence context). We also administered tasks that require participants to generate verbs in response to object drawings (i.e., Verb Generation), or by producing the article and the corresponding noun (i.e., Object Naming). Each of these tasks taps on related, yet distinguishable, language levels. The language processes assessed by each task are schematically represented in Table 4.1.

Table 4.1

*Language processes assessed by each task*

	<u>ObNam</u>	<u>VGen</u>	<u>FinVerb</u>	<u>ActNam</u>
Phonological lexicon (Input)				
Structural description (Input)	✓	✓	✓	✓
Orthographic lexicon (Input)				
Semantic level	✓	✓	✓	✓
Phonological lexicon (Output)	✓	✓	✓	✓
Orthographic lexicon (Output)				
Syntactic level (morpho-syntactic features)	*		✓	

*Notes.* ActNam: Action Naming; ObNam: Object Naming; FinVerb: Producing a finite verb in sentence context; Vgen: Verb generation. Check marks (✓) indicate that the process is assessed in the task. Asterisks (\*) indicate that the process may be assessed in the task. A blank space indicates that the process is not assessed by the task.

#### 4.1.3. Impairment-level and functional-level measures

In brain-damaged individuals, language abilities may be measured with a focus on diagnosing the processing/representational level(s) affected by the lesion, or with emphasis on the subject's ability to use language to communicate. Picture-naming tasks are typically used to obtain impairment-level measures, since they provide objective quantitative (and sometimes qualitative) measures of specific linguistic processes (Katz et al., 2000), such as the ability to retrieve a target word in response to its visual portrayal. In aphasic patients, performance accuracy is usually higher on picture-naming tasks than on tasks that require linguistically more complex responses. For example, Barton, Maruszewski, and Urrea (1969) showed that people with aphasia fare better in picture naming and sentence completion than in naming to verbal description.

Goodglass and Stuss (1979) reported that individuals with aphasia respond more accurately to picture naming than to naming to description.

Role-playing tests, rating scales, and questionnaires are functional-level measures, frequently administered to chronic outpatients. They provide the examiner with a broad picture of the subjects' ability to use language in everyday life (Katz et al., 2000). Role-playing tasks require participants to act out in several fictitious circumstances, mimicking everyday-life events. They may be used to assess how proficiently the patient looks up a number in an agenda, dials a phone number, reports a message, and so on, and how effectively the aphasic speaker uses language when responding to a threat, in a humorous situation, when making a doctor's appointment, or writing down a grocery list. A well-known example of these functional tests is the Communicative Abilities in Daily Living (CADL, Holland, 1980; Second Edition, CADL-2, Holland, Frattali, & Fromm, 1999), which has been shown to measure the severity of language impairment (Fromm & Holland, 1989) and its improvement following treatment (Aten, Caligiuri, & Holland, 1982; see the section Materials). Examples of questionnaires are the Communicative Effectiveness Index (CETI, Lomas et al., 1989) and the Communicative Activity Log (CAL, Pulvermüller et al., 2001). Functional-level measures are valuable tools for language assessment, as the context in which language is produced and the purpose of the communicative interaction affect the choice of words and syntactic structures (Williams & Canter, 1982). The use of synonyms and circumlocutions, or just knowing that responses can be provided without time constraints, may also ease language production or mask a specific deficit, while allowing successful communication (Williams, 1983, for review). Context effects are also reported when using picture-naming tests (Zingeser & Berndt, 1988) and in comparisons between picture naming and spontaneous speech (Herbert, Hickin, Howard, Osborne, & Best, 2008).

Some work has been directed to finding relations between impairment-level and functional-level measures. Aftonomos, Steele, Appelbaum, and Harris (2001) and Bakheit, Carrington, Griffiths, and Searle (2005) found a significant correlation between the patients' scores on the Western Aphasia Battery (WAB; Kertesz, 1982) and caregivers' opinion on their language abilities at different stages of speech and language therapy protocols, as measured by the CETI (Lomas et al., 1989). Best, Greenwood, Grassly, and Hickin (2008) argued for a relationship between the outcome of therapy targeting word retrieval and the patients' opinions on their communicative abilities. In subjects with acquired dysgraphia, Carlomagno, Pandolfi, Labruna, Colombo, and Razzano (2001) showed that improvement of writing performance after treatment correlated with better performance on the reading and writing subtest of the CADL (Holland, 1980). Fucetola et al. (2006) found a positive correlation between comprehension tasks that require semantic processing and the CADL-2 (Holland, Frattali & Fromm, 1998). In a less direct manner, other studies suggested that difficulties producing verbs significantly interfere with connected speech in aphasia (e.g., Bastiaanse & Jonkers, 1998; Crepaldi et al., 2011; Miceli et al., 1989; Saffran, Berndt & Schwartz, 1989; Zingeser & Berndt, 1990).

#### **4.1.4. Questions and hypotheses**

To evaluate the potential benefit of administering verb tasks, we asked (a) whether three commonly used verb tasks provide a sensitive account of (i.e., significantly correlate with) functional measures of language abilities in daily living; and (b) whether any of these tasks provide a more sensitive account of language abilities in daily living than an object naming task matched for difficulty (i.e., if the strength of the correlation between any of the verb tasks and communicative abilities is significantly different to the strength of the correlation between

Object Naming and communicative abilities). To verify that our subject sample was not biased by the inclusion of a greater than normal number of language-impaired individuals with noun- or verb-specific deficits, we compared each individual's accuracy on Object Naming to that on Verb Generation, Producing a finite verb in sentence context, and Action Naming. To ensure that findings could be attributed to the impairment of different processing levels and/or sources of information engaged by verbs and nouns, rather than to an imbalance between psycholinguistic variables shared by these two word categories, stimuli included in all tasks were matched for 12 lexicosemantic and morphosyntactic variables (see Table C4.1).

We predicted that verb tasks would provide a sensitive account of communicative difficulties, as measured by CADL-2 and by the two questionnaires. In addition, we expected that the task requiring the production of finite verb forms would measure language difficulty more reliably than Object Naming and other verb production tasks. This is because only Producing a finite verb in sentence context entails morphosyntactic operations (e.g., argument structure, thematic role assignment, time reference, verb inflection, etc.) in addition to lexicosemantic processes. Morphosyntactic operations are essential for sentence formation (e.g., Berndt et al., 1997; Carlson & Tanenhaus, 1989; Hale & Keyser, 1998; Thompson, 2003) and are frequently affected in aphasia (e.g., Bastiaanse & Jonkers, 1998; Mätzig et al., 2009; Rofes, Bastiaanse, & Martínez- Ferreiro, 2014; Vigliocco et al., 2011). Of course, we did not expect verb tasks to be systematically more impaired than Object Naming. Worse performance on objects than on actions has been reported in patients with temporal lobe lesions (e.g., subjects AN-1033 and Boswell, in A. R. Damasio & Tranel, 1993; see also Daniele, Giustolisi, Silveri, Colosimo, & Gainotti, 1994), while patients with prefrontal lesions frequently show greater difficulty with verb tasks (e.g., subject KJ-1360 in A. R. Damasio & Tranel, 1993; Finocchiaro et



al., 2010; Kemmerer & Tranel, 2000; Lubrano et al., 2014; Rofes, De Witte, Mariën, & Bastiaanse, 2013; Tranel, Kemmerer, Adolphs, Damasio, & Damasio, 2003). This study reports the first attempt to correlate action and object picture-naming tasks, matched for lexicosemantic and morphosyntactic variables, with functional measures of language.

## **4.2. Methods**

### **4.2.1. Participants**

Twenty-one adult, right-handed Italian-speaking aphasic volunteers were recruited (10 female; mean age = 63 years,  $sd = 15$ ; mean years of education = 13,  $sd = 4$ ). They were all at least 6 months post onset of a left-hemisphere stroke (mean months = 40,  $sd = 47$ ) and had moderate-to-severe naming impairment, as shown by performance on the naming subtests of the Battery for the Analysis of Aphasic Disorders (BADA; Miceli, Laudanna, Burani, & Capasso, 1994). Subjects were excluded if they were under age 18, completed fewer than 5 years of formal education, had psychiatric disorders, or were taking drugs that could affect cognitive performance. Participants were relatively heterogeneous as for age, education, lesion site, and time post onset. They were tested as outpatients. Table 4.2 provides information on the main demographic and neurological features of participants. This study was approved by the Ethics Committee of the University of Trento.

Table 4.2

*Main demographic and neurological features of participants*

	<u>Gender</u>	<u>Age</u>	<u>Years of schooling</u>	<u>Etiology</u>	<u>MPO</u>	<u>Lesion site</u>
AL	m	40	17	ICVA	20	LH FT
BL	m	75	18	ICVA	6	LH O
BS	f	43	18	CVST + HCVA	24	LH OT
CB	f	52	13	ICVA	19	LH FTI
CC	f	77	8	HCVA	12	LH P
CK	f	76	19	HCVA	46	LH NB
ES	f	72	8	ICVA	10	LH IC, RH F
FT	f	42	15	ICVA	153	LH FTP
GC	m	68	8	ICVA	30	LH FTPO
GM	f	62	10	ICVA	50	LH FTPB
LF	m	44	15	2 ICVA	40	LH FT, RH OC
OS	f	71	13	ICVA	42	LH TPI
OT	m	65	8	ICVA	7	LH FTI
PG	m	73	10	ICVA	176	LH FTPI
PI	f	51	8	HCVA	6	LH TP
PS	m	76	18	ICVA	15	LH TP
RA	m	57	14	ICVA	8	LH FTP
RG	m	76	14	HCVA	29	LH TP
RL	f	44	13	ICVA	100	LH TI
TG	m	91	11	ICVA	18	LH TPI CR
UA	m	58	13	HCVA	29	LH F, RH FP
m(sd)		63(15)	13(4)		40(47)	

*Notes.* MPO: Months post-onset. ICVA: ischemic cerebrovascular accident; HCVA: hemorrhagic cerebrovascular accident; CVST: Cerebral venous sinus thrombosis; LH: left-hemisphere; CR: Corona radiata; F: frontal; I: insular; NB: Nucleus basalis; O: occipital; P: parietal; T: temporal. m(sd): mean and standard deviation.

## 4.2.2. Materials

### 4.2.2.1. Picture-naming tasks

Each participant completed four picture-naming tasks:

*Object Naming.* Patients see a black-and-white drawing of an object or an animal. The introductory sentence “Here . . .” (Ecco . . .) is written above the drawing. Participants are asked to respond by producing its corresponding noun, with the appropriate determiner. For the picture

of an “apple,” participants are expected to say “Here the apple” (Ecco la mela). This structure is commonly used and fully grammatical in Italian. A Google search for the structure “Ecco la mela” resulted in 17,000 hits.

*Verb Generation.* Patients are shown the black-and-white drawing of an object and must name an action that can be performed with the object, using a verb in the infinitival form. For example, for a drawing of a book, patients are expected to say “to read” (leggere).

*Producing a finite verb in sentence context.* Patients are shown a black-and- white drawing of an action. The subject of the sentence “He/she . . . ” (Lui/lei . . . ) is written above the drawing. Patients are asked to read the pronoun and finish the sentence with the verb in the correct inflected form. For the drawing of “a woman painting,” patients have to say “She paints” (Lei dipinge).

*Action Naming.* A black-and-white drawing of an action is presented. Patients are asked to produce the infinitival form of the verb represented in the drawing. For example, for a picture of “a man kicking a football,” patients are expected to say the verb “to kick” (calciare, in Italian).

Each task contained 20 items with more than 80% picture name agreement, as assessed by the responses of 30 Italian-speaking healthy adults (15 male, 15 female) matched to brain-damaged participants for age (mean = 53.43 years, sd = 15.2) and years of education (mean = 13.93, sd = 3.35). Across tasks, items were matched for lexicosemantic and morphosyntactic variables that affect language production (see Table C4.1 in the Appendices; and the section on the differences between verbs and nouns for a description). These variables may affect processing at the semantic level (imageability, frequency, whether or not objects correspond to biological vs. nonbiological entities, manipulability), at the lexical level (length in phonemes, age of acquisition, name relation to a noun), at the lexical–grammatical level (instrumentality,

transitivity, regularity), and at other levels (e.g., actions performed with face, arm, or leg muscles).

#### **4.2.2.2. Functional measures of language**

Three functional measures of language were used:

*Communicative Abilities in Daily Living, Second Edition* (CADL-2; Carlomagno et al., 2013, Italian version). This role-playing test consists of 50 questions revolving around fictitious environments (e.g., going to the doctor's office, grocery shopping, making a phone call, looking for directions, driving a car). It is used to assess skills such as reading, writing, and calculation; social interactions; divergent communication; contextual communication; nonverbal communication; humor and metaphors; and sequential relations. For example, participants are asked to read a notice and complete a form, to write down three things they may need from a supermarket, and to buy a medicine with the change left after buying a drink.


*Communicative Effectiveness Index* (CETI; Lomas et al., 1989). This questionnaire is completed by the caregiver. It contains 16 questions that resulted from a study in which patients with aphasia and their spouses were asked which communicative situations were relevant in their everyday life. Caregivers rate whether “never, rarely, sometimes, often, or always the patient is able to” get somebody's attention, participate in conversations, communicate emotions, understand writing, describe or discuss something in depth, and so on.

*Communicative Activity Log* (CAL; Pulvermüller et al., 2001). This questionnaire is also completed by the caregiver. It contains 18 questions that revolve around how frequently the patient communicates with friends, in a group, and with one or more strangers; how frequently

she or he uses the phone, listens to or watches the news, writes notes, does simple mathematical operations (e.g., getting change at the supermarket), asks/answers questions appropriately, etc.

#### 4.2.3. Procedure and scoring

Test administration was balanced following a Latin square design. The assessment lasted one hour on average. The CADL-2 was administered and scored following recommendations (Carlomagno et al., 2013). The picture-naming tasks were administered on a laptop computer. To ensure consistent stimulus presentation across tasks, each item was shown for 4000-ms, preceded by a 500-ms beep. This paradigm is used in clinical practice (Brookshire, 1971; Kayama, 2012; Rofes & Miceli, 2014). See Figure 4.1.



*Figure 4.1.* (a) Examples of items from each of the tasks (20 items per task). (b) Structure of an experimental trial: Each image is shown for 4000 ms; a beep is played 500 ms before picture presentation.

Four practice items were included at the beginning of each picture-naming task, to ensure that participants understood the procedure. Incorrect responses were classified as lexical errors (semantic paraphasias were included in this category), part/whole errors (patient says “wheel” instead of car), anomias, circumlocutions, and unrelated words. Formal errors included fragments (“medit... ” instead of “meditate”), phonemic paraphasias, and neologisms. Morphosyntactically incorrect responses included errors of time, person, and number, and morphologically decomposable, derivational neologisms (e.g., “\*fooding” for “eating” where “\*fooding” can be interpreted as being derived from “food”; \*pietanzare for mangiare in Italian). Since subjects had been instructed to produce verbs in the infinitival form in Action Naming and Verb Generation, responses consisting of a verb in the third singular form were counted as errors in these tasks. Additional error types included incorrect responses produced outside the 4-s window. *Conduites d’approche* (effort to approximate to the target item) and hesitations were not scored as errors, as long as the target word was produced within the allotted response time.

Morphosyntactic errors with correct semantic and lexical information were counted as incorrect—for example, “\*Lui mangiare” (\*He to eat) where the verb is in the infinitive instead of the correct inflected form (i.e., Lui mangia, He eats), or \*Ecco la<sub>f.sg.</sub> piede<sub>m.sg.</sub> (This is the<sub>f.sg.</sub> foot<sub>m.sg.</sub>) where a feminine instead of a masculine determiner is used.

### 4.3. Results

Scores were normalized to 100. Nonparametric Friedman’s test  $\chi^2$  was used to check for differences across conditions. Post hoc two-tailed Wilcoxon signed rank tests were also used, when needed. False discovery rate (FDR)-adjusted p-values were systematically calculated to correct for multiple comparisons (Benjamini & Hochberg, 1995). At the group level, correct

response rates on the naming tasks were significantly different (50% for Object Naming, 41% for Verb Generation, 48% for Producing a finite verb in sentence context, and 37% for Action Naming), Friedman's test  $\chi^2(3) = 8.18$ ,  $p = .042$ . However, post hoc tests revealed no significant differences between conditions ( $p < .05$ ). Also at the group level, the patients' performance on CADL-2 was 81% (range = 49–100%)—where lower values denote diminished or poor communicative abilities. On the same scale, the caregiver's opinion of the communicative abilities of the patient's was 72% (range = 42.5–93.8%) for CETI and 71% (range 41.1–96.7%) for CAL. Significant differences were detected between CADL-2, CETI, and CAL, Friedman's test  $\chi^2(2) = 12.10$ ,  $p = .002$ . Post hoc tests revealed significant differences between CADL-2 and CETI ( $Z = 2.36$ ,  $p = .036$ ) and between CADL-2 and CAL ( $Z = 2.43$ ,  $p = .036$ ), but not between CETI and CAL ( $Z = 0.24$ ,  $p = .808$ ). The four naming tasks correlated significantly with CADL-2, CETI, and CAL ( $p \leq .05$ ). Spearman nonparametric correlations were used instead of multiple regression, due to sample size (VanVoorhis & Morgan, 2007; Wampold & Freund, 1987). Even though both Producing a finite verb in sentence context and Object Naming correlated significantly with CADL-2, the strength of the correlation was significantly different (Steiger's  $Z = -2.34918$ ,  $p = .028$ ), being greater for Producing a finite verb in sentence context. No other differences in correlation strength were detected ( $p < .05$ ). Test scores and correlation values are reported in the Appendices (Tables A2 to A5).

Three main response patterns were detected at the individual level. In 13/21 participants (A.L., B.L., B. S., C.C., E.S., F.T., G.M., O.S., P.G., P.I., R.A., R. G., and R.L.), or 61.9% of the sample, performance across tasks was statistically indistinguishable. Subjects A.L., B.S., C.C., E.S., F.T., O.S., P.I., and R.L. performed with relatively high accuracy, and patients B.L., G.M., P.G., R.A., and R.G. fared rather poorly, in all picture-naming tests. Six out of 21 subjects (C.B.,

C.K., O.T., P.S., T.G., and U.A.), or 28.6% of the sample, scored significantly higher on Object Naming than on the verb-naming tasks. P.S. is a clear example of this pattern: He scored 13/20 correct in Object Naming, but scored below 4/20 correct for the three verb tasks. By contrast, one subject (G.C.) performed relatively better on verb tasks than on Object Naming. He provided 5 to 8/20 correct responses in the three verb tasks, but only 1/20 in Object Naming. A last subject (L.F.) performed relatively poorly in all tasks, although he scored significantly better in Producing a finite verb in sentence context than in Verb Generation. Significance tests for each participant are reported in the Appendices (Table C4.6).

Qualitative analysis reveal five major error types. Anomias are the most frequent error type across tasks. Phonemic and semantic paraphasias and responses produced after more than 4 s were also common. Category substitution errors (from verb to noun) were the second most common error type for the Verb Generation task. Error frequency varied slightly across tasks. The major error types per task and participant are reported in the Appendices (Table C4.7).

#### **4.4. Discussion**

The anatomical and behavioral correlates of verbs and nouns have been the focus of many studies in the neuroscience of language (e.g., Black & Chiat, 2003; Shapiro & Caramazza, 2003b; Damasio & Tranel, 1993; Daniele et al., 1994; Mätzig et al., 2009; Pulvermüller, 2005; Vigliocco et al., 2011). This knowledge has made its way in clinical practice only occasionally, and greater attention is still being paid to noun than to verb tasks (Bastiaanse et al., 2003; Conroy et al., 2009; de Aguiar, et al., 2015; Rofes & Miceli, 2014). In this study, we assessed the benefit of administering verb tasks in relation to three functional language measures: the CADL-2




(Carlomagno et al., 2013), the CETI (Lomas et al., 1989), and the CAL (Pülvermüller et al., 2001). Individual analyses were also carried out to check for noun- or verb-specific deficits.

All verb tasks (Verb Generation, Producing a finite verb in sentence context, and Action Naming) significantly correlated with CADL-2, CETI, and CAL ( $p \leq .05$ ). Object Naming also correlated with the three functional level measures ( $p \leq .05$ ). Thus, both verb and object picture-naming tasks would seem to provide reliable indices of language abilities in daily living. More specifically, the processes required to understand a structural description of a picture, to access the corresponding word meaning at the semantic level, and to retrieve its sound elements in the phonological lexicon are good predictors of communicative abilities in everyday life. These processes are shared by the four naming tasks, are commonly used in everyday language, and are typically damaged in people with aphasia (Damasio & Tranel, 1993; Miceli et al., 1988; Miceli et al., 1984; Nickels & Howard, 1995; Tsapkini et al., 2002).

#### **4.4.1. An advantage for producing a finite verb in sentence context over other tasks?**

Interesting differences emerge from the analysis of the relative sensitivity of verb tasks in measuring language difficulties. Producing a finite verb in sentence context strongly correlates with the CADL-2, while only a moderate correlation with CADL-2 is observed for Verb Generation, Action Naming, and Object Naming (see Figure 4.2). In light of the differences between verbs and nouns, it is worth enquiring whether the strength of the correlation between each task and the CADL-2 also differs. As expected, the strength of the correlation between CADL-2 and Producing a finite verb in sentence context is significantly greater than the strength of the correlation between CADL-2 and Object Naming. This indicates that the processes involved in Producing a finite verb in sentence context (i.e., verb retrieval and the

morphosyntactic processes required by naming verbs in sentence context) are better predictors of functional communication skills than the language processes required by Object Naming (see Table 4.1). This is because producing an inflected verb requires lexicosemantic and overt morphosyntactic processes that are not required (at least, not overtly) by Object Naming. For example, producing “Lui mangia” (He eats) in the Producing a finite verb in sentence context task involves at least four steps that are not necessary for Object Naming. The participant must compute (a) a predicate- argument structure indicating which arguments the verb requires to bear out of its meaning (in the example, a subject and an implicit object); and (b) a thematic structure (Carlson & Tanenhaus, 1989) to indicate that the subject position is filled by the agent performing the action (e.g., “He” in “He eats”). Furthermore, the participant is required (c) to establish an agreement relation (Hale & Keyser, 1998) between subject and verb, which in Italian is overtly marked by the morpheme “-a” in “Lui mangia” (i.e., the “-s” in “He eats”); and (d) to locate the action in time by considering tense and time reference (e.g., Bastiaanse et al., 2011). In addition, all these steps also involve working memory processes (Hartsuiker & Barkhuysen, 2006).



*Figure 4.2.* Spearman correlations between Communicative Abilities in Daily Living, Second Edition (CADL-2) and picture-naming tasks. All tasks correlate with the CADL-2. The correlation value of Producing a finite verb in sentence context and CADL-2 is significantly different to that of Object Naming and CADL-2 ( $p = .0324^*$ ).


On these premises, the need to use overt morphology could be a key factor in making Producing a finite verb in sentence context a better predictor of language abilities in daily living than Object Naming and the other verb production tasks. Finding a great number of subject–verb agreement errors in Producing a finite verb in sentence context would strengthen this hypothesis. However, qualitative analyses indicate that overt morphology per se may not be the main factor, as only 3/21 participants produced a maximum of two morphosyntactic errors in this task. Therefore, it is likely that the sensitivity of the Producing a finite verb in sentence context task stems from the need to carry out the complex set of computations listed above (retrieving predicate–argument

structure, assigning thematic roles, encoding agreement relation, tense and time reference, while engaging working memory processes), rather than exclusively or mainly from difficulty with morphosyntactic processes.

However, the strength of the correlations between naming tasks and CADL-2 results from the specific nature of the processes engaged by each naming task, rather than merely from their number. This is suggested by the fact that the correlation between Verb Generation and CADL-2 is strong ( $\rho = .649$ ) but not significantly different to that between Object Naming and CADL-2. If engaging a greater number of steps were sufficient to yield stronger correlation with the CADL-2, Verb Generation should be differentially linked with the CADL-2 as compared with Object Naming, since it requires not only processing of the portrayed object (just like Object Naming), but also the ability to retrieve an action related to the object.

#### **4.4.2. Both object and action naming tasks may be relevant in clinical practice**

Individual-level analyses indicate that the results obtained in our sample do not stem from recruiting a biased number of subjects with noun- or verb specific deficits. Overall, 13/21 participants (61.9%) failed to show significant differences across tasks. Of these, 8 fared relatively well, and 5 relatively poorly in all tasks. Of the participants showing across-task differences, only G.C. was particularly inaccurate in Object Naming (1/20) when compared to Producing a finite verb in sentence context (8/20) and Action Naming (8/20; see Figure 4.3). L.F. scored significantly better at Producing a finite verb in sentence context, although his performance was low overall. The remaining six subjects fared worse on at least one of the verb tasks than on Object Naming.



*Figure 4.3.* Performance patterns in individual participants. Subjects P.I. and P.G. present no difference across tasks (Figures 4.3a and 4.3b). G.C. fares worse in Object Naming than in the other three tasks (Figure 4.3c). P.S. fares worse in the three verb tasks than in Object Naming (Figure 4.3d). ObNam = Object Naming; VGen = Verb Generation; FinVerb = Producing a finite verb in sentence context; ActNam = Action Naming.

These findings alert us to the possibility that clinical language assessments restricted to noun or to verb tasks yield misleading results (e.g., Rofes et al., 2013), by failing to identify individuals whose impairments are limited to one word type, and/or by underestimating communicative problems in a sizeable proportion of aphasic individuals. More sensitive assessments of language functions may be granted by test batteries that tap both noun and verb processing. In the present study, production of finite verbs proved to be a particularly useful addition.

#### **4.4.3. CADL-2, CETI, and CAL as measures of communicative abilities in aphasia**

In the present study, all the functional measures (CADL-2, CETI, and CAL) significantly correlated with all the picture-naming tasks. CADL-2 scores differed from CETI and CAL scores, which in turn did not differ from each other. Furthermore, even though all the picture-naming tests significantly correlated with CADL-2, CETI, and CAL, differences in correlation strength between Producing a finite verb in sentence context and Object Naming reached significance only for CADL-2. Consideration of the structure of CADL-2 leads us to prefer this task to CETI and CAL as a functional measure of everyday communicative abilities. This is because the participant's score relies on an objective analysis of his or her linguistic/communicative skills as they emerge in a number of tasks that are not directly tied to performance in any specific cognitive area (e.g., reading, writing, speech, etc.). This is in contrast with the scores that can be obtained from CETI and CAL, which are based on the subjective opinion of an observer, emotionally involved in the subject's difficulties.

Contrary to our conclusions, other researchers argued that questionnaires such as CETI and CAL are better measures of functional impairment than CADL, because scores are provided by someone who is familiar with the patient's premorbid communicative style (Aftonomos et al., 2001; Bakheit et al., 2005). It could be argued that in our study there was an intrinsic bias for CADL-2 to provide a better fit than CETI and CAL. In the first place, naming accuracy and CADL-2 scores are often collected in the same testing session, and by an external observer who does not have prolonged experience with the subject's current (as opposed to premorbid) communicative difficulty. Secondly, 32/50 questions in CADL-2 use picture stimuli, just like object and action naming tasks. These two objections, however, fail to account for the significantly greater correlation between CADL-2 and Producing a finite verb in sentence

context than between CADL-2 and Object Naming— both tasks are collected essentially simultaneously by an external observer and use picture stimuli. Different correlation strength is likely due to the different linguistic demands of the two naming tasks.

#### **4.4.4. Relevance and limitations of the study**

This investigation is open to questions related to task choice, especially considering that the relative value of the three functional measures used in this study is not established. For example, one might question whether CADL-2 or the two questionnaires provide the more adequate measure of communicative abilities in everyday life. Of course, CADL-2 is not the only objective functional measure of everyday language difficulties—although both CADL and CADL-2 have been frequently used for this purpose (see Fromm & Holland, 1989; Fucetola et al., 2006, for studies on CADL; and Carlomagno et al., 2001, for studies on CADL-2).

Measuring several variables that may influence the narratives elicited during picture description (e.g., Cookie Theft; Goodglass & Kaplan, 1972) could be a suitable alternative, as spontaneous speech would diminish the need for impersonation inherent in CADL-2 (the patient is required to pretend being at the doctor's office, at the supermarket, etc.). Different methods for collecting and analyzing spontaneous speech have been proposed (Craig et al., 1993; Miceli et al., 1989; Nicholas & Brookshire, 1993; Prins & Bastiaanse, 2004; Saffran et al., 1989; Yorkston & Beukelman, 1980), with heterogeneous conclusions. Yorkston and Beukelman (1980) probed the validity of different variables in the spontaneous speech of an aphasic individual, showing that the participant conveyed information informatively (as indicated by the total number of content units), but less efficiently than controls (as indicated by the number of syllables and content units per minute). Similarly, Nicholas and Brookshire (1993) showed that people with aphasia produce fewer words, words per minute, content units, and content units per minute than

non-brain-damaged participants. Jacobs (2001) found a correlation between such measures and the ratings of naïve listeners on the informativeness, efficiency of communication, and listening comfort of spontaneous speech samples before and after specific language treatment. Herbert et al. (2008) found significant correlations between picture-naming accuracy and the proportion of content words as speech units. Berndt et al. (1997) found that people with verb deficits relied on simple sentence structures with light verbs (e.g., give, do, have, be, etc.) and on verbs that do not require inflections (or require zero inflection). Bastiaanse and Jonkers (1998) and Crepaldi et al. (2011) reported that people with nonfluent aphasia produce a lower proportion of verb types in spontaneous speech than non-braindamaged speakers. Goral and Kempler (2009) indicated that naïve listeners perceived a positive change (e.g., less stress, less awkwardness) in the communicative abilities of a person with chronic nonfluent aphasia after training of verb production.

Thus, CADL–2 and spontaneous speech analyses can be considered as good predictors of the patient’s communicative abilities in daily life, as both rely on objective measures of the patient’s behavior and therefore provide more reliable results than subjective rating scales, exposed to potential biases on the examiner’s part. On the other hand, both are exposed to the same criticism— namely, that it is not yet completely understood whether they provide sufficient information on functional communicative abilities. Our (slight) preference for CADL–2 stems from the fact that scores on this task are obtained via a procedure shared by the users of this test. By contrast, further work is needed to establish which features of spontaneous speech provide the most sensitive measure of effective communication in daily life (e.g., content units, fluency, mean length of utterance, proportion or ratio of specific word categories, number of syllables, type–token ratio, proportion of inflected verbs, etc.).



Regarding impairment-level measures, we administered four picture-naming tasks that require the ability to produce nouns that refer to objects (e.g., apple, arm, trumpet) and verbs that refer to actions (e.g., to count, to moo, to squeeze). Further work could be directed towards finding correlations of functional-level measures with nouns that do not refer to objects (e.g., month, future, realm) or verbs that do not refer to actions (to love, to trust, to shine). Other tasks requiring spoken output, such as repetition and connected speech, or comprehension tasks could be administered in addition to picture-naming tasks.

As in any correlational study, results are indicative. The statistics employed were adequate in the light of sample size and number of items included in each task (Benjamini & Hochberg, 1995; VanVoorhis & Morgan, 2007; Wampold & Freund, 1987). Further work on a larger subject sample should correlate the patterns observed here with lesion site and with the underlying language deficit. Most participants in this study suffered from single focal lesions in the middle cerebral artery territory (e.g., middle and inferior frontal gyrus, insula, inferior parietal lobule, superior and middle temporal gyrus). Further insights on this issue may be gained by studying subjects who fare worse in Object Naming than in verb tasks (such as our case G.C.), whose lesions typically affect the posterior cerebral artery territory, thus damaging inferomesial temporo-occipital areas (cases B.L., B.S.), or subjects with etiologies other than stroke (e.g., degenerative or slowly growing lesions, in which brain plasticity may modify anatomo-behavioral correlates).

#### **4.5. Conclusion**

Verb tasks correlated with abilities in daily living, as measured by CADL-2. Producing a finite verb in sentence context and Object Naming differently correlated with CADL-2. More subjects fared worse on one or more verb tasks than on Object Naming, but the opposite pattern was also reported. These patterns should be further investigated by focusing on lesion site, etiology, and underlying language deficit. Results strongly encourage the use of both verb tasks and noun tasks in clinical practice, when evaluating subjects with language disorders.

## Chapter 5

# Advantages and disadvantages of intraoperative language tasks in awake surgery: a three-task approach for prefrontal tumors<sup>5</sup>

### Abstract

*Introduction:* Multidisciplinary efforts are being made to provide surgical teams with sensitive and specific tasks for language mapping in awake surgery. Researchers and clinicians have elaborated different tasks over time. A fair amount of work has been directed to study the neurofunctional correlates of some of these tasks, and there is recent interest in their standardization. However, little discussion exists on the advantages and disadvantages that each task poses from the perspective of the cognitive neuroscience of language. Such an approach may be a relevant step to assess task validity, to avoid using tasks that tap onto similar processes, and to provide patients with a surgical treatment that ensures maximal tumor resection while avoiding postoperative language deficits. An understanding of the language components that each task entails may also be relevant to improve the current assessments and the ways in which tasks are administered, and to disentangle neurofunctional questions.

*Methods:* We review 17 language mapping tasks that have been used in awake surgery. We provide examples of a three-task approach we are administering to patients with prefrontal lesions.

*Results:* Overt production tasks have been a preferred choice over comprehension tasks. Tasks tapping lexico-semantic processes, particularly object naming, maintain their role as gold standards. Automated speech tasks are used to detect speech errors and to set the amplitude of the stimulator. Comprehension tasks, reading and writing tasks, and tasks that assess grammatical aspects of language may be regularly administered in the near future.

*Conclusions:* Future advances in this area are contingent upon reviewing gold standards and introducing new assessment tools. Our three-task approach is feasible and useful.

*Keywords:* brain mapping, language tests, semantics, quality of life

---

<sup>5</sup> Published manuscript. Rofes, A., Spena, G., Miozzo, A., Fontanella, M.M., & Miceli, G. (in press). Advantages and disadvantages of intraoperative language tasks in awake surgery: a three-task approach for prefrontal tumors. *Journal of Neurosurgical Sciences*.

## 5.1. Introduction

There is increasing interest in the neurocognitive components of language mapping in awake surgery: efforts are being directed towards providing surgical teams with appropriate assessment tasks (De Witte, Satoer, Robert, Colle, Verheyen, Visch-Brink et al., 2015; Polczyńska 2009; Rofes, de Aguiar, Miceli, 2015); literature exists regarding the selection of tasks based on lesion localisation and functional networks (e.g., Chang, Raygor, & Berger, 2014; Coello, Moritz-Gasser, Martino, Martinoni, Matsuda, & Duffau, 2013; Hamberger, 2015; Mesulam, 1990); and relevant work has been devoted to studying the neural correlates of language tests in relation to direct electrical stimulation (DES) and brain plasticity (e.g., Binder, Swanson, Hammeke, & Sabsevitz, 2008; Desmurget, Song, Mottolese, & Sirigu, 2013; Foki, Gartus, Geissler, & Beisteiner, 2008; Kristo, Raemaekers, Rutten, de Gelder, & Ramsey, 2015; Picht, Krieg, Sollmann, Rösler, Niraula, Neuvonen, et al., 2013; Zacà, Nickerson, Deib, & Pillai, 2012; Zacà, Jarso, & Pillai, 2013). The field is blooming with new and increasingly refined approaches, encouraged by the fact that language mapping in awake surgery provides better outcomes for the patient than fully-anesthetized surgeries (De Witt Hamer, Gil-Robles, Zwinderman, Duffau, & Berger, 2012). However, little discussion exists on the advantages and disadvantages that currently used tasks entail for the assessment of language during surgery.

Here we review 17 language mapping tasks commonly reported in the literature. This may help awake surgery teams to choose which task(s) may better suit their surgical needs, to avoid administering tests that assess too similar processes, and to preserve the patient's language abilities and quality of life. We discuss the language components assessed by each task from a cognitive neuropsychological perspective (Coltheart, 2001; Whitworth, Webster, & Howard, 2005). We take into account language components that are typically agreed-upon in language

processing models (e.g., Caramazza 1997; Levelt 1989). We consider the semantic level as a central component, as its damage may affect production and comprehension processes, with consequent impact in the patient's quality of life (Goodglass & Wingfield, 1997; Weitzner, Meyers, & Byrne, 1996). We also discuss the order of administration of tasks and provide an overview of an intraoperative three-task approach we are successfully implementing. Finally, we point toward some future directions in this field.

Discussion over non-language factors such as type and timing of electrical stimulation, histological factors, lesion localisation, extent of tumor resection, patient's profile, variability, anxiety, responsiveness, etc. is outside the scope of this paper. Attention has been often paid to these issues (e.g. Borchers, Himmelbach, Logothetis & Karnath, 2011; Gil-Robles & Duffau, 2010; Hamberger 2015; Ojemann & Mateer, 1979; Ruis, Wajer, Robe, & van Zandvoort, 2014; Santini, Talacchi, Squintani, Casagrande, Capasso, & Miceli, 2012; Skirboll, Ojemann, Berger, Lettich, & Winn, 1996; Talacchi, Taphoorn, & Miceli, 2012). Similar work could be aimed at reviewing tasks that are used to assess memory (Feudio & Van Buren, 1975; Ojemann, 1978; Ojemann & Dodrill, 1985; Teixidor, Gatignol, Leroy, Masuet-Aumatell, Capelle, & Duffau, 2007), executive functions (Wager et al., 2013), visual processing (Duffau, Velut, Mitchel, Gatignol, & Capelle, 2004), mentalizing (Herbet, Lafargue, Moritz-Gasser, Bonnetblanc, & Duffau, 2014), numerosity and calculation (Roux, Boukhatem, Draper, Sacko, & Démonet, 2009), etc.

### **5.1.1. Language testing in awake surgery**


Before being operated upon, patients undergoing awake surgery may be assessed with batteries that allow an evaluation of language in its different input and output components (e.g.,

reading, writing, speaking, and listening). Tasks prepared for intraoperative use are administered at this stage to decide which items or item-categories can be used during surgery. Only those items (or subsets of items) that patients can produce flawlessly before surgery are administered intraoperatively, as this is thought to minimize false positives (i.e., instances in which the patient is unable to answer in the absence of electrical stimulation) and to provide reliable language maps (Rofes, de Aguiar, & Miceli, 2015). During surgery, low-intensity electrical trains are directly applied to the cortex and subcortical areas, while the patient is asked to respond to each language stimulus (e.g., Ojemann & Mateer, 1979). The intraoperative occurrence of speech errors (i.e., anomia, paraphasias, latencies, etc.) during language mapping is taken as an indication that the stimulated region is a relevant part of the language network (see Borchers et al., 2011, Desmurget et al., 2013, for contrasting opinions). The types of errors that emerge during electrical stimulation of an area may reveal the role that area plays in the language network. For example, stimulation to the superficial layer of the inferior fronto-occipital fasciculus (IFOF) may trigger semantic errors in production (e.g., saying “tiger” in response to the picture of a lion) and in comprehension (e.g., stating that “pyramid” is related with “palm tree”, but not with “pine tree”), suggesting that this fiber bundle plays a role in semantic processing (Moritz-Gasser, Herbert, & Duffau, 2013).

### **5.1.2. Models of language processing**

Models of language processing are the operationalization of theories of language production and comprehension. They are based on studies in healthy people and on behavioral dissociations documented in neurological populations (e.g., patient A is worse at writing nouns than producing nouns aloud, while patient B has the opposite profile). The articulation of these

models may be schematically represented in box-and-arrow diagrams, where boxes represent separable memory/language components and arrows correspond to connections between these components. A schematic representation of one of such model can be found in Figure 5.1.



*Figure 5.1.* Schematic representation of language components. The schema includes input and output phonological and orthographic lexical components that revolve around a semantic level. It also includes a visuo-perceptual, and a (lexico-) syntactic component.

On a broad level, input lexical components ensure that the subject recognizes orthographic or phonological sequences as familiar (i.e., as words stored in orthographic/phonological long-term memory). The output lexicons provide access to orthographic or phonological sequences that are familiar to the subject. The semantic system represents meanings (i.e., the word /dog/ stands for a typically domesticated mammal that has a good sense of smell, howls and barks, has a long snout, etc.) and is accessed both for production and for comprehension. Syntactic processes allow us to glue words into sentences to express complex meanings, to refer to time and place, etc. (i.e., “My wife has two dogs”). Visuo-perceptual processes are needed for the interpretation of pictorial stimuli (e.g., Gibson, 2013). Finally, articulatory processes allow the production of speech sounds through the correct movement of the speech organs (e.g., tongue, lips, larynx, etc.).

### **5.1.3. Advantages and disadvantages of tests reported in the literature**

The language processes and modalities tested, the time needed to perform the task (ideally, no more than 4 seconds per item), and the patient's pre-surgical profile (cognitive abilities, age, education, pathology, etc.) determine advantages and disadvantages of intraoperative tasks. We briefly describe each task, with a focus on the language processes and modalities tested. We comment on other factors and contrast various tasks, where necessary. We discuss the tasks in subgroups, according to whether they pose greater demands on non-semantic/sublexical, lexico-semantic, or grammatical processes. This does not imply that some tasks grouped among those tackling lexico-semantic processes, for example, may not be useful to assess also grammatical processes (e.g., action-naming). A summary of the language processes assessed by each task can be found in Table 5.1.



Table 5.1

*Language processes assessed by each task.*

	<u>PhL</u> (Input)	<u>StrD</u> (Input)	<u>OrL</u> (Input)	<u>SemL</u>	<u>PhL</u> (Output)	<u>OrL</u> (Output)	<u>SynL</u>
Non-semantic processes							
Automated speech				*	✓		
Phonemic identification	✓				✓		
Word/non-word repetition	✓			*	✓		
Lexico-semantic processes							
Action-naming		✓		✓	✓		
(Visual) Object naming		✓		✓	✓		
(Auditory) Object naming	✓			✓	✓		
Naming famous faces		✓		✓	✓		
PPTT (with drawings)		✓		✓			
PPTT (with written words)			✓	✓			
PPTT (with words and drawings)	✓	✓	✓				
Reading words/non-words aloud	*		✓	✓	✓		
Verb generation		✓		✓	✓		
Writing words			✓	✓		✓	
Grammatical processes							
Naming actions with finite verbs		✓		✓	✓		✓
Reading sentences aloud			✓	*	✓		*
Sentence completion			✓	✓	✓		✓
Sentence repetition	✓			*	✓		✓
Translating paragraphs			✓	✓	✓		✓
Writing sentences			✓	✓		✓	✓
Writing sentences (to dictation)	✓		✓	✓		✓	✓

*Notes.* Check marks (✓) indicate that the process is assessed in the task, asterisks (\*) that it may be assessed, and blank spaces that it is not assessed. PPTT = Pyramids and Palm Trees test, PhL = Phonological lexicon, StrD = Structural description, OrL = Orthographical lexicon; SemL = Semantic level; SynL = Syntactic level (morphosyntactic features).

### 5.1.3.1. Non-semantic processes

*Automatic speech tasks* require motor planning and articulatory processing. Patients can be asked to count from one to ten, to recall the months of the year, the days of the week, etc. These language stimuli do not pose high demands on linguistic knowledge as they typically contain overlearned sequences of words (Bookheimer, Zeffiro, Blaxton, Gaillard, & Theodore,

2000). Such tasks are administered to set the intensity of the electrical stimulator and to check for effects on the peripheral nervous system (Duffau et al., 1999). Some teams have asked patients to count from one to fifty (e.g., Sanai, Mirzadeh, & Berger, 2008; Chacko et al., 2013). Oral diadochokinesis or fast naming of sequences of phonemes (e.g., pa-ta-ka, pa-ta-ka, pa-ta-ka) could also be used, as they do not pose high demands on lexico-semantic processes (e.g., Hukrmans, Jonkers, Boonstra, Stewart, & Reinders-Messelink, 2012). Ojemann and Mateer (1979) asked patients to make orofacial movements, such as mimicking postures of lip protrusion when seeing the action performed on a picture. Special attention may be paid to the rate at which the patient is asked to produce automatic speech, as a fast speech rate may induce undesired spontaneous errors, leading intraoperatively to false positives.

*Phonemic identification* was used by Ojemann and Mateer (1979). Patients were presented with plosive consonants embedded in a nonsense syllable (e.g., a[p]ma vs a[b]ma), and were asked to decide which plosive consonant they heard. The task may detect the inability to recognize and isolate phonemes within a word, which may in turn prevent access to phonological word forms (e.g., [b]all vs [t]all), and hence, their comprehension. We are not aware of other studies administering this task during awake surgery. Caplan, Gow and Makris (1995) reported the scores of people with acoustic-phonetic processing deficits after stroke. They reported that patients with lesions in the left posterior supramarginal gyrus as well as in the parietal operculum are more likely to present poor phonemic identification. The question remains as to whether it may be useful to modify the original task proposed by Ojemann and Mateer (1979), and ask patients to identify target phonemes in different word positions (e.g., onset vs coda position), or to assess specific contrasts (e.g., voicing, manner and place of articulation, etc.).

*Word/non-word repetition* may be useful to tap on language routes that do not require semantic access or to identify the language level(s) affected by electrical stimulation. Tomasino et al. (2015) asked Italian-speaking patients to repeat non-words such as “bolata”. Moritz-Gasser and Duffau (2013) asked patients to repeat the target word whenever they produced it incorrectly in an object naming context during stimulation mapping. Successfully repeating the target word after a previous naming error during stimulation may indicate a temporary failure to access the phonological output lexicon. Similarly, an inability to name the target word may signal disruption in processing its structural description (if images are used), or in accessing the phonological/orthographic input lexicon or the semantic level (if spoken/written words are used as stimuli). This may be particularly useful in perioperative assessments.

### **5.1.3.2. Lexico-semantic processes**

*Action-naming* assesses the ability to produce a verb as the label for the action depicted in a drawing/picture. The task has been used, although not extensively (see Rofes & Miceli, 2014 for a review). Patients are given a black-and-white drawing or a video of an action, and they are asked to answer with a verb in the infinitive (e.g., “to jump”) or in the progressive form (“jumping”). Bogka et al. (2003) ran three studies on healthy people and showed that naming actions may be slower than naming objects – but, still within the 4-second window allowed in awake surgery (e.g., mean=1051 milliseconds, sd= 171 milliseconds). The processes engaged by action-naming are similar to those of object naming. The main limitation of action-naming resides in the fact that it does not require the production of inflectional features which, in many languages, encode time and person information, and are involved in lexical and grammatical processes in discourse (Tyler, Bright, Fletcher & Stamatakis, 2004). This relevant linguistic

information may be left unassessed when an action-naming task is used instead of a task that requires production of a finite verb (see later in this section).

*Visual object naming* assesses the ability to produce a noun (e.g., “book”) as the label for the object depicted in a drawing/picture. This task has been used by the pioneers of awake surgery (Ojemann & Mateer, 1979), is mentioned in current surgical guidelines (Kayama, 2012), and has been reported as a good predictor of the patient’s likelihood to go back to work, and of overall quality of life (Moritz-Gasser, Herbet, Maldonado, & Duffau, 2012). In versions of this test for Dutch, English, French, Italian, and Japanese an introductory sentence has been added above the image to induce the production of the determiner with the noun (e.g. “This is... *an* apple” instead of “apple”). As a result, patients are asked to produce a sentence with a light verb “to be” (This is... *an apple*) or a determiner phrase (*Ecco la mela*, Here the apple). This latter task requires the overt production of grammatical information, as the selection of the appropriate determiner requires the patient to encode number (i.e., *an* apple vs *some* apples), reference to the noun in previous discourse (i.e., *the* apple vs *an* apple), and in some languages the noun’s grammatical gender (i.e., “*la*.*f.sg.* *mela*” instead of “*the*.*m.sg.* apple”). As stressed by Ojemann and Mateer (1979), the introductory sentence “this is” allows to distinguish anomic errors (patients read the introductory sentence but are unable to produce the determiner and the noun) from speech arrest, in which patients are unable to speak. Similar arguments are found in recent reports (e.g., Zemmoura, Herbet, Moritz-Gasser, & Duffau, 2015). This interpretation is open to discussion, as the inability to produce the introductory sentence is not necessarily due to problems in motor production or planning. It can also result from other problems, such as the inability to read the introductory sentence, due to difficulty accessing the orthographic input lexicon.

*Auditory object naming* (or naming objects to visual description) is similar to visual object naming, insofar as the response required of the patient is the name of an object. However, instead of being presented with a drawing/picture, patients hear a spoken description of an object, of which they have to provide the name (e.g., “a hand-held instrument used for cutting paper” for scissors). This task requires accurate auditory word and sentence comprehension (not necessary in visual object naming), and the ability to retrieve the target name that matches the stimulus sentence. It may be particularly sensitive in patients with left temporal lobe epilepsy, compared to visual object naming (Hamberger & Tamny, 1999; Hamberger et al., 2005). Bird, Howard and Franklin (2000) successfully used this task in people with post-stroke aphasia. By not using visual stimuli, the task can be relevant to assess the semantic level in patients who cannot read, or when the patient position may difficult the presentation of visual stimuli such as in midline (sub)occipital craniotomies. Also, it is suitable to assess word types that cannot be easily depicted (e.g. ego, welfare, attractive, deny, promote, scent, noticeably, etc.)

*Naming famous faces* requires similar processes to object naming, as well as facial feature encoding and accessing semantic/biographic information of the person that is portrayed. The final output required from the patient is a proper name such as “Chaplin” or “Madonna”. Giusanni et al. (2009) used a task with 30 famous faces during surgery. The authors argue that famous faces are more complex than objects, as the visual and lexico-semantic characteristics required to process a famous face correspond to a unique entity, while those required for the correct processing of an object correspond to a category. As any other task, a standardization process is required to decide which famous figures are better known to patients – age and cultural background may influence these results. Giusanni et al. (2009) were able to find specific areas for this task compared to object naming in the left frontal and anterior temporal gyri, and

propose that naming tasks should be adapted to the brain region studied. In perioperative studies of people with brain tumors, temporal lobe epilepsy, and other neurological disorders, damage to the left anterior temporal lobe can disrupt access to the names of famous people, particularly when damage extends to the uncinate fasciculus (e.g., for a review, see Papagno, 2011)

The *Pyramids and palm trees test* assesses “the person’s ability to access detailed semantic representations from words and from pictures” (Howard & Patterson, 1992). The test consists of triads of pictures or written/spoken words. A stimulus item is presented, centered at the top of the page (e.g., a pyramid) and two alternatives are presented below it: a target (e.g., palm tree) and a distracter (e.g., pine tree). The patient’s task is to choose which of the two items is associated with the item presented on top. To do so, patients need to access information from the three items in each triad (i.e., recognize the items, retrieve information, perform the association). Moritz-Gasser et al. (2013) and Motomura, Fujii, Maesawa, Kuramitsu, Natsume, and Wakabayashi (2014) used the version containing three images during stimulation of the IFOF and as a complement to object naming. The authors argue that this task allows the identification of brain regions related to non-verbal semantic memory. They report that during DES patients were unable to make a semantic choice, looked startled and perplexed, and sometimes answered with sentences like “I don’t know at all”, “I do not understand anything” “what do I have to do?”. Howard and Patterson (1992) indicated that six other versions of the test are possible (three written words, three spoken words, and combinations of those with pictures as stimuli or as alternative choices). These authors emphasize that patients may not be assessed with all versions routinely. Rather, a specific version may be selected depending on which language level must be further assessed. For example, the version with three auditorily presented words may be recommended to evaluate access to semantics via the phonological input lexicon.

This version of the task should be used in patients with good phonological short-term memory, as it requires the subject to hold information from the three words in memory in order to provide the correct answer.

*Reading words/non-words aloud* requires access to the orthographic input lexicon. Sanai et al. (2008) used this task during surgery. Similarly to auditorily presented tasks, reading tasks can be useful to assess words that are difficult to depict. The task allows the assessment of very specific types of words, if needed, while allowing to control for psycholinguistic variables such as frequency, age of acquisition, imageability (see Brendt, Mitchum, Haendiges, & Sandson, 1997; for an example in people with post-stroke aphasia). Zemmoura et al. (2015) asked patients to read 20 single words shown on a computer screen for 4 seconds, after a beep presented 500ms before item presentation. The task includes regular words (i.e., words with a transparent phoneme-grapheme relationship: *bilatéral*, bilateral), irregular words (i.e., words with opaque orthography-to-phonology relationship: for example, the word *oignon*, onion, is read out /ɔ.ɲɔ̃/) and non-words/pseudowords (i.e., letter sequences that are not represented in the vocabulary: *bafiko*). The three types of stimuli were presented randomly. The patients' performance allowed the assessment of both non-lexical reading mechanisms (non-words) and reading mechanisms that required lexical-semantic processing (irregular words). Since the task uses individual stimuli, pre-stimulus triggers may be administered during electrical stimulation.

*Verb generation* tasks require comprehending a noun in order to produce a conceptually-related verb in the infinitive. Patients are shown the picture or told the name of an object (e.g., “book”) to which they have to associate the first verb that comes to mind (e.g., “to read”). Verb generation involves language processes that impact on the semantic level and can be perturbed by stimulation in multiple brain areas (Ojemann, Ojemann, & Lettich, 2002; Roux, Boetto,

Sacko, Chollet & Trémoulet, 2003). However, this task does not assess syntactic processes. In addition, if electrical stimulation disrupts performance, it is difficult to establish if it interfered with noun comprehension, verb retrieval, or both. This distinction has relevant implications for mapping language processes, as nouns and verbs may have a partially segregated representation in the brain, also as suggested by studies using DES (Corina, Gibson, Martin, Poliakov, Brinkley, & Ojemann, 2005; Havas et al., 2015; Lubrano, Filleron, Démonet, & Roux, 2014). Verb generation tasks should not be confounded with *word generation or fluency tasks*, in which patients are given one minute to produce as many words as possible starting with one specific letter or referring to a specific semantic category (Goldstein, Obrzut, John, Hunter, & Armstong, 2004). These latter tasks may be appropriate for perioperative testing, as a way to assess cognitive control and executive functions in general (Moritz-Gasser & Duffau, 2013). However, the use of fluency tasks to map language during surgery is questionable, as it is hard to establish an appropriate stimulation time to trigger specific language errors. Furthermore, in normal settings subjects do not produce a continuous flow of words, probably due to word search processes.

*Writing words to dictation* may be useful to tap lexical-semantic knowledge of words that are difficult to depict. Since this is a one-stimulus task, instead of a sentence-production task, electrical stimulation may be applied right after word dictation, before the patient provides an answer. The task may be easier to perform and to administer than writing sentences. Magrassi, Bongetta, Bianchini, Berardesca, Arienta (2010) reported on a patient who wrote words correctly under dictation or spontaneously, but was unable to write simple sentences. In some languages, this task can be used to assess different writing systems. Motomura et al. (2014) took advantage of the Japanese writing system and asked a participant to write kanji (morphograms) when cued



with kana (syllabograms). For example, the patient read the word grass in kana (クサ - /kusa/) and was asked to write it down in kanji (草). Similar work could be performed in languages like Serbian, which is normally written in both Latin and Cyrillic alphabets.

### **5.1.3.3. Grammatical processes**

*Naming actions with finite verbs* requires the patient to retrieve an inflected verb – i.e., the verb root that corresponds to the action and the specific morphosyntactic features that relate the subject of the sentence and the verb. Patients are presented with a picture or video of an action. An introductory sentence may be added above the image, consisting either of a determiner phrase referring to the object (i.e., “the woman”) or of a pronoun (“she”). Patients are required to read the introductory sentence and to produce a verb in its correct inflected form. This task targets verb production, involves semantics, lexical retrieval and the syntax required to establish the agreement relation between subject and verb, time reference, etc. Importantly, these processes can be performed within the 4-second window allowed by current electrical stimulation procedures (Kayama, 2012). Rofes, Capasso and Miceli (2015) showed that this task is a good predictor of language abilities in daily living, based on data of individuals with post-stroke aphasia. The authors recommend to associate this task to object naming when assessing language functions, as the production of nouns and verbs may dissociate, even when items are matched for relevant psycholinguistic variables.

*Reading sentences aloud* has typically been implemented by asking patients to read sentences of 8 to 10 words slowly (e.g., Ojemann, Ojemann, Lettich, & Berger, 1989; Roux, Lubrano, Lawers-Cances, Trémoulet, Mascott, & Démonet, 2004). Lubrano et al. (2004) asked patients to read two unrelated sentences that were not semantically related and that had not been

previously trained (e.g., *La chaise est jolie. Le vent du nord souffle fort*; The chair is pretty. The north wind blows hard). A caveat of this task is that the stimulation point is hard to establish. Roux et al. (2004) reported that “stimulation was applied randomly on the cortex while [patients] were reading”. Other than that, due to the non-semantic grapheme-to-phoneme conversion route, patients may perform well on this task without accessing semantics. And obviously, this task is not appropriate for patients who cannot read.

*Sentence completion* involves similar processes to Naming actions with finite verbs. Patients are given a string of 7 to 8 words forming a sentence, in which one component is missing. Ojemann and Mateer (1979) presented sentences of this kind: “If it’s sunny next Saturday she ... to the beach”, where the target is “will go”. When the missing word is the lexical verb (e.g., “to go”), this task directly taps lexical-semantic and syntactic mechanisms, as it requires the ability to retrieve the appropriate verb for the context provided by the sentence. This task also allows the assessment of verb inflection, as well as of different time frames (e.g., past, present, future) and agreements between subject and verb (i.e., first person, second person, etc.). Patients could also be asked to produce a noun (e.g., “If it’s sunny next Saturday she will go to ...; where the target could be “the beach”). The task is not suitable for people who cannot read. Stimulation may be applied before the production of the target word, as at this point in time the processes necessary to complete the sentence may be engaged. The fact that patients need to complete the task with one word poses greater demands on lexico-semantic processing than reading sentences aloud.

*Sentence repetition* adds the grammatical processes that are required in sentence processing to those engaged by word repetition. Tomasino et al. (2015) asked patients to repeat active and passive sentences from an Italian language battery (BADA, Miceli, Capasso,

Laudanna, & Burani, 1994). Patients produced phonemic paraphasias that extended to more than one word in the sentence (\**il cavàno incine il cane* instead of *il cavallo insegue il cane*; the horse chases the dog). Errors like these raise the possibility that electrical stimulation affects speech output not only at the single-word level but also at the sentence level, similar to other tasks that use sentences. Also in this case it is difficult to establish when in time (and where in the brain) electrical stimulation should be applied.

*Translating paragraphs* is only applicable to proficient bilinguals. Borius et al. (2012) asked patients to translate paragraphs from newspapers from L2 to L1 and reported no specific sites for translation (as compared to naming or sentence reading). Stimulation during this task may lead to relevant but difficult-to-interpret results, as translation requires many language processes to be concomitantly activated (i.e., reading and understanding in L2; finding the appropriate words, building the correct syntactic frame, and reading it out in L1). Further work is needed to establish the relevance of this task in mapping protocols with proficient polyglots. Until then, it may still be profitable to assess patients with similar tasks, or parallel versions of the same task, in the two languages (e.g., object naming in French and Occitan).

*Writing sentences* can be used to trigger errors at various stages of written output (semantic level, syntactic level, orthographic word forms, orthographic short-term memory, graphomotor routines, etc.). Since the task requires reading of the text that is being written, in addition to engaging the orthographic input lexicon. Errors arising at this latter level may be difficult to detect, as written word form recognition processes are covert. Stimuli similar to those used in sentence reading tasks can be administered. Patients are asked to write sentences of seven or eight words. They may be active sentences (e.g., *La chaise est jolie*; The chair is pretty; Lubrano et al., 2004; Roux et al. 2003; 2014), or more syntactically-complex constructions (i.e.,

*Quando il gatto non c'è i topi ballano*; When the cat is away the mice will play; *Luca mangia un panino e abbraccia Carlino*; Luca eats a sandwich and hugs Carlino; Magrassi et al., 2010). Sentences are typically dictated by the examiner, thus engaging the phonological input lexicon and allowing in principle the task to be performed without accessing semantic-lexical information (via conversion routes). Roux et al. (2014) reported that in rare cases patients may present difficulty understanding the task. In such instances, the authors complemented it with other tasks tapping the phonological output lexicon and conversion routes (e.g., word repetition), and with tasks that require access to the semantic system such as choosing one picture (e.g., a flower, a dog, a hammer, a boat) and is asked to indicate “where is the boat?” Similar to sentence reading, writing sentences to dictation poses the problem to decide when to apply electrical stimulation. The types of errors that may be encountered during electrical stimulation are: irregular handwriting, drift from the initial line of the sentence, disorganization of the text in space, letter/phoneme displacements or substitutions, repetitions, using upper-case letters instead of lower-case letters within words, inability to write (i.e., writing arrest). Roux et al. (2014) divide errors in two categories: semantic or verbal paraphasias and phonological paraphasias. Further work is needed to improve task administration, as in order to write while being operated upon, patients may need to take an uncomfortable position. Lubrano, Roux, & Démonet (2004) report that “patients used a pencil to write horizontally on A4 sheets of paper, which lay on a stiff pad that a nurse had presented to them vertically”.

All the tasks that require producing sentences, orally or in writing, can be influenced by disorders of working memory. These might be diagnosed in perioperative stages by administering cognitive tasks such as the Digit Span, the Corsi's block-tapping test (e.g. Orsini,

Grossi, Laiacona, Papagno, & Vallar, 1987), the Immediate Visual Memory test (Gainotti, Caltragione, & Miceli, 1978), etc.

#### **5.1.4. Adminsitration of tasks during surgery**

Little research exists onto the order in which the tasks are administered. Automatic speech tasks may be used first to assess non-linguistic motor processes, and to regulate the amperage of the electrical stimulator (e.g., Duffau et al., 1999). Subsequently, the patient may be presented with language tasks (typically object naming) during cortical stimulation, in order to delineate a cortical language map. During tumor resection, some groups engage the patient in naming tasks in the absence of any electrical stimulation. These tasks may be paired with a motor task, such as moving the right arm to obtain further on-line feedback (e.g., Gil-Robles & Duffau, 2010). Another possibility is to engage the patient in a spontaneous conversation. Patient may be asked questions about their lives that may have been learned by the examiner in the preoperative assessment (e.g. “You told me that you have two children. Which is the one who is married?”). Questions related to whether the patient feels pain, is comfortable, thirsty, etc. may also be asked. Language-driven approaches may also be used in which patients are engaged in semi-structured interviews instead of spontaneous conversations, as these allow further control on speech production and can be contrasted with the same patient’s preoperative assessments, and with those of a non-brain-damaged control group (Prins & Bastiaanse, 2004; Saffran, Berndt & Schwartz, 1989). Finally, the same tasks used for cortical stimulation may be used for subcortical stimulation. Some groups used the Pyramids and Palm trees test for subcortical mapping (e.g., Moritz-Gasser et al., 2013; Motomura et al., 2014). Other groups (De Witte et al., 2015) propose

a fast-sequencing of different language and non-language tasks (i.e., word and sentence repetition, naming, calculating, and line bisection).

The ordering of tasks may become relevant when a patient does not respond to one task but responds to the other. For example, Magrassi et al. (2010) asked the patient to write words, as she was unable to write sentences. It could be argued that the task the patient is more respondent to may be used first for language mapping. This raises the question of whether any other task may be more sensitive or specific (either because of the localization of the lesion, or because of the nature of the tasks). If a patient performs poorly in a specific modality (e.g., oral production), another modality may be used (e.g., written production). Whenever the patient unexpectedly fails to respond to all the tasks that engage semantic and lexical processes (e.g., object naming, action-naming by finite verbs), it may be relevant to consider the results of tasks that do not require semantic processing, such as automated speech tasks, non-word repetition, non-word reading, etc. These procedures may diminish the number of spontaneous errors, while increasing the sensitivity and specificity of DES mapping.

Another example where task order may become relevant is when tasks with verbs and sentences are used together with object naming tasks (see below). Task order may differ depending on lesion location and on the hypotheses the surgical team has. Surgical teams can choose to use verb tasks first in the case of prefrontal lesions, as aphasiological and lesion studies indicate a greater involvement of the prefrontal cortex in the processing of verbs, as opposed to the processing of nouns (e.g., Mätzig, Druks, Masterson, & Vigliocco, 2009). The opposite considerations apply in the case of temporal tumors, where object naming may be used first. Irrespective of any considerations on task order, the results of the various tests should be interpreted together.

## **5.2. A three-task approach for prefrontal tumors**

Our current approach to intraoperative language mapping includes an automated speech task (counting from one to ten) and two overt language production tasks: naming objects (determiner + noun) and naming actions by producing finite verbs (pronoun + finite verb). The two naming tasks have been standardized for Italian and validated on an aphasic population (Rofes et al., 2015). The approach builds on modern surgical procedures (Duffau et al., 1999; Hamberger & Tamny 1999; Kayama, 2012; Ojemann & Mateer 1979; Ojemann et al. 2002; Sanai et al. 2008), and stresses the need to administer standardized tasks. The choice of the two naming tasks was driven by current knowledge in the cognitive neuroscience of language, and may be particularly efficient in patients with prefrontal lesions (Kemmerer, Rudrauf, Manzel, & Tranel, 2012; Mätzig et al., 2009; Rofes & Miceli, 2014). The approach we advocate is conservative, insofar as it does not implement too many tasks. At this stage, it is also suitable, as there are no standardized Italian versions available for all the tasks that could be used for language mapping. A good understanding of which tasks are preferable during surgery may require rigorous analyses and contrastive results pointing to the sensitivity, specificity, and predictive values of newer tests. In the final section of the study we describe our experience with this three-task approach.

### **5.2.1. Participants**

Three Italian-speaking subjects (all male) with supratentorial glial lesions in left-dominant prefrontal areas were included. Age ranged from 37 to 63 years, education from 8 to 17 years. TT was operated for an anaplastic astrocytoma (WHO III) occupying the superior frontal gyrus (SFG) and middle frontal gyrus (MFG). PR was operated for a fronto-polar astrocytoma (WHO

III) extending to the MFG. MG was re-operated for an oligodendroglioma (WHO III) in the precentral gyrus and posterior part of the SFT, MFG, and inferior frontal gyrus (IFG).


### **5.2.2. Materials and procedure**

Object naming and naming actions with finite verbs were administered intraoperatively. These were also administered preoperatively, to decide which items could be administered during surgery. The tasks were part of a broader perioperative language and cognition screening protocol. Detailed information on the tasks is reported elsewhere (Rofes et al., 2015). Items were presented for 4 seconds, as proposed in current surgical protocols (Kayama, 2012). Trains of electrical stimulation (60Hz, 0.2msec) lasting 4 seconds were delivered with a bipolar electrode. The amperage of the stimulator was regulated with an automated task (i.e., counting from one to ten). It varied for each participant and ranged from 2.5-4.5mA. All surgeries were video-recorded. We report areas where brain stimulation induced at least two errors out of three stimulations during language mapping with object naming, naming actions with finite verbs, and with both tasks. These areas were considered as critical for language processes and were not removed during surgery. Post-hoc two-tailed Fisher's exact tests (FE) were calculated to assess whether the number of correct and incorrect responses differed between the stimulation and non-stimulation condition in each stimulated area.



### 5.2.3. Results

Results for each patient are summarized in Figure 5.2.



*Figure 5.2.* Summary of language maps. Areas where cortical stimulation induced a significant representative amount of errors and that were therefore considered as responsive during surgery are indicated. *Blue squares* indicate areas where errors were triggered during DES and object naming; *Orange squares* areas where errors were triggered during finite verb naming; and *Green squares* areas where errors were triggered with both object naming and finite verb naming. The area occupying the tumor is marked in red.

In patient TT errors appeared with both object naming and naming actions with finite verbs during stimulation of the posterior part of the MFG. In a more superior area of the MFG, errors were only detected during DES with finite verbs. At the cortical level, 2 errors in 21 stimulations appeared during object naming ( $p=0.1364$ ) and 4 errors in 20 stimulations during naming actions with finite verbs ( $p=0.2213$ ). At the subcortical level, 6 errors in 13 stimulations were elicited during object naming ( $p=0.08$ ) and 6 errors in 16 stimulations during naming actions with finite verbs ( $p=1.000$ ). Response latencies longer than 4 seconds were the most common error type, followed by anomia.

In patient PR, DES to the posterior part of the MFG triggered 2 errors with object naming. These were semantic paraphasias that were self-corrected within the 4-second time window (i.e., *ecco la sedia, il tavolo* – here the chair, the table; *ecco la chitarra, il violino* – here

the guitar, the violin). A more superior area within the MFG was detected during DES with both object naming and naming actions with finite verbs. Incorrect responses resulted in latencies that resolved after removing the electrical dipole. Errors in the posterior part of the SFG were only triggered when naming actions with finite verbs. PR produced the introductory sentence, but failed to produce the verb in the correct inflected form (i.e., *lui... lui...*; he... he...). Subcortical stimulation was delivered to areas corresponding to the fronto-polar cortex. During DES with object naming the patient produced a semantic paraphasia (*ecco la prugna* for *ecco il peperone* – “Here the plum” for “Here the pepper”) and two anomias (*non mi ricordo* – “I don’t remember”). During the task with finite verbs the patient perseverated, while producing the pronoun in the incorrect gender in the introductory sentence (*lui lei pettina, lei pettina, lei lui pettina, lei lui pettina, lui pettina* – he she combs, she combs, she he combs, she he combs, he combs – the target is “she combs”). Statistical analyses are not provided, as this surgery could not be fully videotaped.

In patient MG no errors were triggered during naming actions with finite verbs. During object naming, errors occurred in 3/12 stimulations of the middle part of the IFG ( $p=0.001$ ) and 3/10 stimulations of the MFG ( $p=0.005$ ). Errors were phonological in nature. The patient produced a phonemic paraphasia or a fragment of the target, always followed by the correct response (e.g., *ecco la cigliere, ciliegia* – here the cherry; *ecco la gira, giraffa* – here the giraffe; *ecco il pf, peperone* – here the pepper). Stimulation to the subcortical area corresponding to the inferior part of the prefrontal cortex during naming actions with finite verbs induced a phonological error and anomias, that persisted when no stimulation was applied.

To sum up, mapping with object naming and naming-finite verbs seemed relevant in this population. *Post-hoc* analyses do not indicate a superiority for any of the tasks. The two tests can

complement each other: in TT and PR areas in the posterior MFG and STG were only detectable when naming actions with finite verbs and/or object naming, while in MG errors were triggered only during object naming.

### **5.3. Discussion: future steps of language tests in awake surgery**

We reviewed 17 different task types that have been used in awake surgery. These tasks cover non-semantic, lexico-semantic, and grammatical processes, mostly for overt production. The fact that such a wide range of tasks already exists indicates both the interest and the need to provide surgical teams with sensitive and specific tasks for intraoperative language mapping. In the next years, we expect to see a large number of tests, carefully controlled and standardized for each specific language. In fact, some surgery groups are already using standardized tests instead of home-made tasks (e.g., De Witte et al., 2015; Moritz-Gasser et al., 2012; Połczyńska 2009; Rofes et al., 2015). We also expect to see an increase in the variety of tasks.

Sufficient discussion of the different tasks will need to rise, so that surgical teams have an adequate understanding of the benefits and risks that each task entails. We believe that progress in this area requires that awake surgery teams add one or two new tasks to their current protocols, so that it is possible to learn whether new methodologies provide more specific and reliable results than the current gold standard. Roux et al. (2014), for example, administered three tasks in their patients provided that “brain mapping in awake surgery can be rather time consuming and some patients can obviously be tired (with some loss of attention) after several tasks”. Our team is also administering two-language tasks and an automated speech task. This is similar to the regular surgical procedure adopted by other teams (e.g., Lubrano et al., 2014;

Motomura et al., 2014; Moritz-Gasser et al., 2014; Tomasino et al., 2015). Further discussion on the tasks and paradigms used is essential.

The question arises of whether new tasks may give a direct oncological benefit by enhancing the patient's quality of life and survival time. Certain new tasks may be given priority, since they are more sensitive and specific, and of greater predictive value than the old ones. Histopathological, demographic, neurofunctional and many other factors may play a role in answering this question. Research into relations between functional and impairment-based scores could reveal the extent in which damage to specific components of the language system may affect language abilities in everyday life and, therefore, quality of life (e.g., Moritz-Gasser et al., 2012; Rofes et al, 2015). Further work may also be devoted to understanding the extent in which mild postoperative language damage may be tolerable or acceptable by an individual patient (e.g., Sanai et al., 2008).

At this stage, overt production tasks are more widely used than comprehension tasks. This could be because they are easier to score and provide better on-line feedback (Gil-Robles & Duffau, 2010). Other production tasks (e.g., writing words, producing sentences, naming actions with finite verbs) may be put to regular use in the near future. Reading individual words as well as sentence completion tasks may provide a bridge from the current, noun-centered processing approaches to tasks that require reading and writing, while controlling the timing of electrical stimulation. At the same time, given the paucity of comprehension tasks (note that the Pyramids and Palm Trees task is the only one used currently), we expect researchers and clinicians to be also interested in them. Of course, we do not expect nor wish that automatic speech or object naming tasks be left aside. The former are necessary to establish the amplitude of the stimulator,

and the latter will probably maintain their prominent role in tackling semantic and lexical processing.

Irrespective of the specific task(s), advances in this area are contingent on reporting for each task detailed information on its theoretical background and internal structure. We refer to the types of stimuli used, whether or not stimuli were standardized, which language processes were tapped by the task, etc. Efforts may also be devoted to describing test administration and scoring procedures, such as how long the stimulus was available on-screen, when stimulation was applied, the statistical methods used to identify specific brain areas (e.g., comparison of stimulation sites between two tests, comparison of correct and incorrect responses, etc.), and which error types emerged during stimulation.

#### **5.4. Conclusion**

We reviewed tasks currently used in awake surgery. Advantages and disadvantages of these tasks were assessed within a theoretical framework based on language components that are largely agreed upon in cognitive neuropsychology models of language processing. We reviewed our current approach, based on three tasks for prefrontal gliomas and showed that a task that uses object naming and naming actions with finite verbs is relevant. Progress in this area will be forthcoming if surgical teams work on the standardization of tasks in their respective languages and report their experiences when administering new tasks, compared to current gold standards. This work should involve the collaboration of different professional figures in the surgical team and researchers working in the neurobiology of language.



# Chapter 6

## Surgical mapping of language production in sentence context: a single-case series<sup>6</sup>

### Abstract

*Introduction:* Object naming has been traditionally used to map language production in patients undergoing awake surgery procedures. Tests that use verbs in sentence context can be an appropriate addition to reach the same goal, as they use actions instead of objects and assess lexical verb retrieval mechanisms and sentence formation processes that cannot be tapped by object naming alone.

*Aim:* To investigate the contribution of finite verb production in sentence context for language mapping in awake surgery. Emphasis is placed on individual results, on subcortical mapping and on the types of errors elicited during DES, for which consistent information is currently scarce.

*Methods:* Six Italian-speaking participants with gliomas in the left, language-dominant hemisphere. Two overt production tests were administered: an object naming task (noun task) and a task requiring the production of finite verbs in sentence context (verb task). Factors affecting correct and incorrect responses were evaluated by non-parametric statistics. We contrasted the performance in all sites during DES v no-DES collapsing across nouns and verbs, or considering performance on nouns and verbs separately. Also, we contrasted noun and verb production during DES independently of site, and for each stimulation site separately.

*Results:* More errors were detected during DES v no-DES. The cortico-subcortical representation of nouns and verbs and the error types observed during DES are partially consistent with the current literature. In four participants we detected language-relevant areas only with object naming, in one participant only with the verbs task, and in one participant with both tasks.

*Conclusions:* The production of finite verbs in sentence context may provide complementary information to object naming, as it may permit to map language production when object naming is unaffected by DES.

*Keywords:* brain mapping, language production, verbs, nouns

---

<sup>6</sup>Submitted to Acta Neurochirurgica. Rofes, A., Spina, G., Talacchi, A., Santini, B., Miozzo, A., & Miceli, G. Surgical mapping of language production in sentence context: a single-case series.

## 6.1. Introduction

Direct electrical stimulation (henceforth, DES) is the current gold standard for language mapping in individuals undergoing surgery for glial tumors in left perisylvian areas (e.g., Desmurget, Song, Mottolese, & Sirigu, 2013). Patients are kept awake under local anesthesia and are asked to perform cognitive tasks, while short trains of electricity are delivered in cortical or subcortical structures (e.g., Ojemann & Mateer, 1979; patient 22 in Penfield & Boldrey, 1937). Brain tissue in which DES triggers reproducible errors is not excised, as it is deemed that its removal may affect relevant cognitive functions (cf. Borchers et al., 2011). Keeping the patient awake during surgery offers advantages over fully asleep procedures, as it allows obtaining direct feedback on the patient's cognitive abilities, thus shedding light on the integrity of the underlying representations and processes (e.g., Surbeck, Hildebrandt, & Duffau, 2015).

Various cognitive processes can be assessed during surgery. In the language domain, object naming has been traditionally used to prevent functional damage (e.g., Hamberger & Tamny, 1999; Lubrano, Filleron, Démonet, & Roux, 2014; Ojemann & Mateer, 1979). This task has a long tradition in lesion studies. It is sensitive to lexical-semantic damage, which affects word production and comprehension, and is known to heavily impact on quality of life (Gainotti, Silveri, Villa, & Miceli, 1986; Goodglass & Wingfield, 1997; Moritz-Gasser, Herbet, Maldonado, & Duffau, 2012). Therefore, in all likelihood it will keep a central role in awake surgery. However, a mapping procedure based exclusively on object naming does not provide a sufficiently detailed intraoperative evaluation of language skills. From the strictly cognitive/linguistic viewpoint, object naming does not tap onto other relevant language functions, such as the processing of action words and morphosyntax, which are necessary for sentence formation and correlate with language abilities in daily living more strongly than object



naming itself (e.g., Rofes, Capasso, & Miceli, 2015). From the clinical perspective, it has been repeatedly shown that nouns and verbs can be selectively affected/spared by brain damage (for reviews, see Pillon & d'Honinckthun, 2011; Vigliocco et al., 2011), and that difficulties with verbs may selectively affect verb retrieval or verb morphology (e.g., Shapiro & Caramazza, 2003; Miceli, Mazzucchi, Menn & Goodglass, 1983; Miceli, Silveri, Romani, & Caramazza, 1989), in the context of spared object naming.

A finer-grained mapping of language may be obtained by adding simple intraoperative tasks that require the ability to produce verbs in a phrasal context, such as completing or reading sentences, or producing verbs in sentential context. A test that requires the production of a finite verb in a sentence, in response to a pictured event, for example, requires interpreting the stimulus picture, retrieving the meaning of the verb, building its corresponding argument structure, assigning thematic roles (agent, theme, beneficiary, etc.) to the different components of the sentence, and producing the verb in its correct inflected form (e.g., in agreement with the preceding subject, and in the correct time frame). Adding a finite verb production task to an object naming task would allow us to evaluate intraoperatively, and in a short time, not only semantics, lexicon and phonology, but also lexical-grammatical properties and morphosyntax (for a discussion of intraoperative language tests, see Rofes, Spina, Miozzo, Fontanella, Miceli, 2015).

Studies of neurological populations and of healthy volunteers with non-invasive neuroimaging techniques serve as a stepping stone for the introduction and development of mapping tools in awake surgery. Regardless of the origin of the differences between nouns and verbs, lesion studies indicated that verb processing is more frequently and severely impaired than noun processing (for reviews see Mätzig, Druks, Masterson, & Vigliocco, 2009; Pillon &

d'Honichthun, 2011). Analyses of the anatomoclinical correlates in these studies led to propose that deficits in producing nouns most likely arise from temporal damage, and deficits in producing verbs from frontal and (to a lesser extent) parietal and temporal damage (e.g., Damasio & Tranel, 1993; Goodglass, Klein, Carey, & Jones, 1966; Kemmerer, Rudrauf, Manzel, & Tranel, 2012; Miceli, Silveri, Noncentini, & Caramazza, 1988; Silveri & Di Betta, 1997).

Functional Magnetic Resonance Imaging (fMRI) studies of noun and verb production in healthy volunteers show activations in large perisylvian areas, that are more extensive and widely distributed than previously hypothesized (for reviews see, Cappa & Perani, 2003; Crepaldi, Berlingeri, Paulesu, Luzzatti, 2011; Vigliocco, Vinson, Druks, Barber, & Cappa, 2011). Several accounts can be proposed for these findings. One possibility is that, since fMRI paradigms show task-correlated activation that may not be essential for naming, some hyperactive foci may not be specific to verbs. For example, Thompson-Schill, D'Esposito, Aguirre, and Farah (1997) found significant activation in the left inferior frontal gyrus (IFG) during several semantic tasks (i.e., verb generation, noun classification, noun comparison). They interpreted their results as evidence that the left IFG is critical for "selection of information among competing alternatives from semantic memory", rather than for the retrieval of semantic information, or for verb processing. It is also possible that verbs activate widely distributed neural networks because, unlike nouns, they also entail complex lexical-grammatical properties (e.g., argument structure, subcategorization restrictions, etc.), and are typically inflected for tense, aspect and person, thus playing a crucial role for morphosyntactic processes (e.g., the Italian form *mangia*, refers to a third person singular agent that performs the action of eating in the present moment). Each of these verb properties may involve at least partially distinct neural networks, and therefore be responsible for widespread activations in fMRI studies. Another

possibility is that these diverse observations result from the different focus of various investigations of the noun/verb distinction, and hence on how stimuli were selected in the original studies. For example, Saccuman et al. (2006) found no differences between nouns and verbs, but reported fronto-parietal activations in the contrast between manipulable or hand action words (e.g., the scissors/to comb) v non-manipulable words (e.g., the pyramid/to walk). By contrast, Bedny and Caramazza (2011) found a posterior region of the middle temporal gyrus (MTG) to be more activated by verbs than by nouns, and by syntactic properties of verbs, irrespective of motion properties.

Like fMRI studies, Transcranial Magnetic Stimulation (TMS) investigations failed to document the involvement of specific cortical regions in naming pictures of objects v actions, but demonstrated that object naming tasks are more sensitive than verb naming tasks when mapping language in the perisylvian cortex, and that both tasks induce errors as compared to a no-TMS condition (Hernandez-Pavon, Mäkelä, Lehtinen, Lioumis, & Mäkelä, 2014). Some TMS studies also highlight the role of psycholinguistic dimensions. For example, Oliveri, Finocchiaro, Shapiro, Gangitano, Caramazza and Pascual-Leone (2004) found no differences in the production of singular/plural nouns and third-person singular/plural verbs during TMS of the motor cortex, but reported differences in the production of action-related words, regardless of grammatical category (e.g., the axe/to bite v the cloud/to adore).

Even though agreement on the underpinnings of the noun/verb distinction is only partial, mapping the processes underlying noun and verb production during surgery, and understanding the neural mechanisms underlying such distinction would be relevant to surgical teams. In fact, DES of brain sites that are more (or, selectively) involved in processing one or the other word type may increase the probability to detect functionally relevant tissue in the periphery of the

tumor (Fernández Coello, Moritz-Gasser, Martino, Martinoni, Matsuda, & Duffau, 2013; Roux & Trémoulet, 2002; Rofes & Miceli, 2014). To support the relevance of this issue, in one of the few studies that used verbs intraoperatively, Lubrano et al. (2014) indicated that nouns and verbs may be represented in partially segregated networks. They compared the production of verbs in sentence context with that of object names, and found specific object naming sites in the temporal cortex and inferior frontal gyrus (IFG), and specific action naming sites in the IFG and posterior middle frontal gyrus (MFG). Similar data emerged from intraoperative studies that used non-finite verbs (e.g., to play, playing) (i.e., Bello et al., 2007; Corina, Gibson, Martin, Poliakov, Brinkley, & Ojemann 2005, Havas et al., 2015).

Knowledge gathered from studies on the cognitive neuroscience of nouns and verbs can be applied to construct or to refine assessment and treatment tools for patients with neurological damage, and to draw hypotheses on the sensitivity and specificity of brain regions involved in noun and verb processing (e.g., de Aguiar, Paolazzi, & Miceli, 2015; Vigliocco, Vinson, Druks, Barber, & Cappa, 2011). Neurosurgical studies are particularly useful to address these issues. In fact, the intraoperative sensitivity and specificity of DES in detecting functionally-relevant tissue are difficult to achieve preoperatively with non-invasive neuroimaging techniques, such as fMRI or TMS, due to aspects of the so-called brain shift such as drainage of cerebrospinal fluid, tissue swelling/damage, patient position, etc. (e.g., Nabavi, Stark, Dörner, & Mehdorn, 2012; Picht et al., 2013).

The neurosurgical approach to the intraoperative study of finite verb production (and of language mapping in general) may benefit from rigorous methodological approaches, in order to overcome some current limitations. A first step in this direction could be the systematic use of standardized tasks, developed specifically to meet the constraints of intraoperative mapping. So

far, many potentially relevant results were obtained with home-made tasks, that are exposed to the influence of nuisance factors and therefore pose obvious problems of duplicability of results. Another resource could be provided by careful quantitative and qualitative analyses of the responses produced by patients during DES.

In the operating room, patients' responses are typically scored dichotomously, and only general qualitative information is provided (but, see e.g. Hamberger & Tamny, 1999; Lubrano et al., 2014, for exceptions). A response produced correctly within 4 seconds (i.e., the maximum duration allowed for DES to prevent seizures Duffau, 2004; Kayama, 2012) is taken to indicate that the stimulated structure has no role in the function tapped by the task. By contrast, any response that differs from the target or is produced later than the allowed response time counts as an error, and is taken to show that the stimulated site plays a relevant role in the cognitive function under exam. More detailed error analyses could lead to a better understanding of brain-behavior relations and therefore to preventing functional damage during surgery.

Different types of incorrect responses can occur during intraoperative language mapping with both object and action naming, pointing to different underlying neurofunctional processes. For example, the fact that DES in the temporal lobe and/or along ventral associative pathways (e.g., the inferior fronto-occipital fasciculus, IFOF) may trigger whole-word errors (e.g., semantic paraphasias and failures to respond), has been taken to suggest that these brain structures are involved in lexical-semantic processes (e.g., Caplan, Vanier, & Baker, 1986; Caramazza, Papagno, & Ruml, 2000; Sarubbo et al., 2015). At segment-level errors (e.g., phonemic paraphasias) during DES of the prefrontal cortex and/or dorsal associative pathways (e.g., arcuate fasciculus, AF) may indicate that these latter structures play a role in retrieving and

producing segmental information (e.g., Binder, Desai, Graves, & Conant, 2009; Hickok & Poeppel, 2007; Lubrano et al., 2014).

### **6.1.1. Aims and predictions**

The present study evaluates the relevance of tasks that assess language in sentence context. We sought to identify cortical regions involved in the production of nouns and in the production of verbs in sentence context, using similar methods to recent studies (Corina et al., 2005; Herbet, Lafargue, Moritz-Gasser, Bonnetblanc, & Duffau, 2014; Lubrano et al., 2014). To provide sensitive and specific surgical mapping results, we administered tasks designed and standardized for awake surgery (Rofes, de Aguiar, & Miceli, 2015), performed a rigorous preoperative examination, looked for significant DES sites on individual subjects, and contrasted the results of each individual in a single-case series fashion. To complement the current literature, we also provided data on subcortical DES and a careful examination of error types.

Based on the current literature, we expected the DES condition to yield more errors as compared to the no-DES condition. DES may interfere more with finite verb production than with noun production, when delivered to the prefrontal cortex, as suggested by some DES, TMS and lesion studies (e.g., Cappelletti, Fregni, Shapiro, Pascual-Leone, & Caramazza, 2008; Lubrano et al., 2014, cf. Crepaldi et al., 2011; Hernandez-Pavon et al., 2014 for more distributed representations in neuroimaging studies). As regards error types, DES of cortical and subcortical regions corresponding to dorsal pathways (e.g., frontal and parietal cortices; AF) should elicit mainly phonological paraphasias, and DES of cortical and subcortical regions corresponding to ventral pathways (e.g., infero-mesial and infero-lateral temporo-occipital cortices; inferior-frontal occipital fasciculus, IFOF) should elicit mainly semantic paraphasias (Sarubbo et al.,

2015, cf. Caplan et al., 1986; Caramazza et al., 2000, Lubrano et al., 2014). No specific predictions were formulated as regards anomias, since failure to name has been associated with damage to a largely distributed network including MFG, STG, IPL and corresponding subcortical sites (e.g., Lubrano et al., 2014; Rofes & Miceli, 2014; Sarubbo et al., 2015).

## 6.2. Methods

### 6.2.2. Participants

Six Italian-speaking, right-handed subjects with supratentorial tumors in the left hemisphere participated in this study. Subject characteristics and tumor variables are summarized in Table 6.1.

Table 6.1

*Subject characteristics and tumor variables*

	<u>Age</u>	<u>Gender</u>	<u>Edu.</u>	<u>Job</u>	<u>Hand.</u>	<u>Grade</u>	<u>Type</u>	<u>Site</u>
CRA	39	F	8	Homemaker	Right	II	Oligodendroglioma	Parietal (IPL)
CRO	70	F	13	Retired	Right	IV	Glioblastoma	Prefrontal (MFG)
MG*	63	M	17	Stock trader	Right	III	Oligodendroglioma	Prefrontal (SFG,MFG,IFG)
PR	41	M	13	Builder	Right	III	Astrocytoma	Prefrontal (FP, MFG)
SO*	45	M	8	Carpenter	Right	II	Oligoastrocytoma	Prefrontal (IFG, MFG)
TT	37	M	8	Builder	Right	III	An.astrocytoma	Prefrontal (SFG, MFG)

*Notes.* An. = anaplastic.; Hand.=handedness; Edu. = Education in years; FP = Fronto polar; IFG = Inferior frontal gyrus; IPL = Inferior parietal lobe; MFG = Middle frontal gyrus; STG = Superior temporal gyrus; \*Subject had been previously operated at another institution.

### 6.2. Materials

A three-task intraoperative protocol was administered. It included an automated speech task (counting from 1 to 10) and two oral naming tests: an object naming task and a task that requires the production of finite verbs in a simple sentence context. Both tests are standardized and have

been specifically designed for awake surgery procedures. In the object naming task, patients are asked to produce a noun and a determiner phrase (*Ecco la mela*; “Here [is] the apple”<sup>7</sup>). In the production of finite verbs, patients produce a pronoun and an action verb in the correct inflected form (*Lei abbraccia*; “She hugs”). In the finite verb production task, 74% of the agents are masculine (e.g. “He irons”) and 26% are feminine (e.g., “She sings”). For a detailed description, see Rofes et al., 2015a. To minimize errors not triggered by DES, intraoperative tests were constructed with items and item-categories to which patients responded flawlessly before surgery. The following psycholinguistic contrasts were implemented in the experimental lists administered pre-operatively: frequent v non-frequent words; long v short words (in phonemes); biological v non-biological nouns (“apple” v “table”); instrumental v non instrumental verbs (“to sew” v “to jump”); manipulable v non-manipulable verbs (“to type” v “to jump”); verbs name-related to a noun v verbs not-name-related to a noun (“to brush” v “to sew”); regular v irregular verbs (“to play” v “to sing”); and transitive v intransitive verbs (“to cook” v “to walk”). These dimensions were considered in constructing the finite verb production task based on the evidence reported in the literature (for review, see Rofes & Miceli, 2014; Rofes et al., 2015a; Rofes et al., 2015b). All stimuli were selected based on the controls’ ability to respond correctly in 4 seconds. This is standard practice in awake surgery, as it indicates a safe DES time that triggers language errors reliably and with minimal risk of intraoperative seizures (e.g., Duffau, 2004; Kayama, 2012). A bipolar electrode delivering 4-second trains of DES (50~60Hz, 0.2msec) was used. The amperage of the stimulator ranged from 2.5-10mA.

---

<sup>7</sup> This utterance is grammatical in Italian.



#### **6.2.4. Procedure**

An awake craniotomy was performed to expose cortical structures. Sterilized tags were placed in the periphery of the tumor. The counting task was administered to detect areas relevant to oral movements and to set the amperage of the stimulator. Patients were asked to respond to the two picture-naming tasks to reveal areas relevant for language production. The naming tasks were administered intraoperatively at least twice: for cortical and for subcortical DES. Each stimulus was presented for 4 seconds on a computer laptop. Areas where at least 2 out of 3 instances of DES triggered errors were considered relevant for language production and were not removed.

All surgical procedures (with the exception of subject PR's, see below) were video recorded to perform post-hoc quantitative and qualitative error analyses. Responses were counted as correct if the target item was produced within 4 seconds. Barnard's tests were used to compare the number of correct/incorrect responses in the different conditions: DES v no-DES, object naming v finite verb production, by test and by area. This test is particularly powerful for data analysis in 2x2 contingency tables. We used one-tailed statistics because we expected that DES would interfere with performance, and therefore that the number of correct responses in the DES condition would be lower. The types of errors observed during DES were also noted (e.g., anomia, phonological paraphasia, fragment, semantic paraphasia, latency, morpho-syntactic error). Stimulated areas were discussed with the neurosurgeon on the basis of sulcal and gyral anatomy (e.g., SFG, MFG, IFG, etc.). A similar methodology has been reported in the literature (e.g., Lubrano et al., 2014; Herbet et al., 2014).

## 6.3. Results

### 6.3.1. Quantitative analyses

In pre-operative assessments, none of the potentially relevant psycholinguistic dimensions (e.g., frequency, word length, semantic category, instrumentality, transitivity, etc.) significantly constrained performance in any participant (Table 6.2). Therefore, all categories were used for intraoperative testing, and for each category only the items to which patients had responded incorrectly in the preoperative assessment were eliminated from intraoperative mapping.


Table 6.2

*Preoperative differences in psycholinguistic dimensions (subsets)*

		<u>CRA</u>	<u>CRO</u>	<u>MG</u>	<u>PR</u>	<u>SO</u>	<u>TT</u>
Nouns	Bio_Sub	FE=0,476	FE=0,737	FE=1,000	FE=1,000	FE=0,604	FE=1,000
	Frequency	FE=1,000	FE=0,741	FE=0,342	FE=1,000	FE=0,342	FE=0,605
	Length	FE=0,738	FE=0,065	FE=0,346	FE=1,000	FE=1,000	FE=1,000
Verbs	Tr_Sub	FE=0,459	FE=1,000	FE=0,233	FE=0,489	FE=1,000	FE=1,000
	Reg_Sub	FE=1,000	FE=1,000	FE=1,000	FE=1,000	FE=0,484	FE=0,226
	NRN_Sub	FE=0,576	FE=0,153	FE=1,000	FE=0,471	FE=1,000	FE=1,000
	MN_Sub	FE=1,000	FE=0,088	FE=0,490	FE=1,000	FE=1,000	FE=0,189
	Length	FE=0,349	FE=0,258	FE=1,000	FE=1,000	FE=1,000	FE=1,000
	Frequency	FE=1,000	FE=1,000	FE=0,485	FE=1,000	FE=0,485	FE=1,000
	Inst_Sub	FE=0,666	FE=1,000	FE=0,234	FE=1,000	FE=0,489	FE=1,000

*Notes.* FE = Fisher's Exact Test for count data, p-value. Biol = biological v non-biological entities ("apple" v "table"); Frequency = frequent v non-frequent words; Length = long v short words (in phonemes); Inst = Instrumental v non instrumental verbs ("to sew" v "to jump"); MN= Manipulable v non-manipulable verbs ("to type" v "to jump"); NRN = Name related to a noun v non-name related to a noun ("to brush" v "to sew"); Reg. = Regular v non-regular verb ("to sing" v "to play"); Tran. = Transitive v intransitive verb ("to cook" v "to walk"). Each of the subsets is explained in Rofes et al. 2015b.

Intraoperative results for the different conditions are reported in Table 6.3. Figure 6.1 shows the brain areas where a statistical difference was found between the number of correct and incorrect responses during DES with object naming, production of finite verbs, or both. DES-positive cortical sites are reported in Figure 6.1A, subcortical DES sites in Figure 6.1B.



*Figure 6.1.* Significant DES sites per subject. A = cortical, B = subcortical. The number of stimulations is higher during cortical than subcortical DES in subjects CRA (68 v 17), SO (56 v 25), and TT (44 v 25). Cortical and subcortical stimulations were equal for CRO (14). Subcortical DES was not performed to MG. The number of stimulations could not be counted in PR.

Subject CRA produced no errors in 32 stimulations during object naming (henceforth, 0/32 errors, where the numerator indicates the total number of errors and the denominator the total number of stimulations per test) and 9/36 errors in the production of finite verbs. The difference

between the DES condition for objects and finite verbs was significant (Wald=2.752; Nuisance=0.414;  $p=0.003^*$ ). CRA produced a significantly greater number of errors during the DES v no-DES condition at the subcortical level (7/10 v 3/31, Wald=2.646; Nuisance=0.112;  $p=0.006^*$ ). Object naming triggered a significant number of errors during subcortical DES, possibly in the posterior part of the arcuate fasciculus (3/7 errors, Wald=1.922; Nuisance=0.223;  $p=0.037^*$ ). Mapping with the finite verb task fell just short of significance, at the subcortical level (4/10 errors; Wald=1.922; Nuisance=0.111;  $p=0.051$ ).

In subject CRO, mapping with the finite verb task was not possible due to the large number of spontaneous errors (5/12) produced in the absence of DES. In object naming, this subject produced more object naming errors during DES as compared to no-DES (6/8 v 1/20 errors, Wald=2.556, Nuisance=0.223;  $p=0.006^*$ ). This was particularly obvious following DES in the middle part of the MFG (3/4 errors; Wald=3.122; Nuisance=0.131;  $p=0.001^*$ ) and in the posterior part of the MFG (3/3 errors; Wald=3.521, Nuisance=0.182;  $p=0.001^*$ ), which also corresponded to a face motor area, as indicated by intraoperative motor mapping. No significant differences were observed during subcortical mapping.

Subject MG produced 0/11 errors during DES with finite verbs, and 7/33 errors during object naming. Error rate was greater in the DES than in the no-DES condition (Wald=4.062; Nuisance=0.061;  $p=0.001^*$ ), specifically for object naming (Wald=3.485; Nuisance=0.091;  $p=0.001^*$ ), and in the middle part of the IFG (3/11 errors; Wald=3.766; Nuisance=0.021;  $p=0.001^*$ ), the middle and posterior parts of the MFG (2/8 errors; Wald=3.503; Nuisance=0.041;  $p=0.003^*$ ; and 2/5 errors; Wald=4.311; Nuisance=0.031;  $p=0.002^*$ , respectively). DES was applied three times to 2 subcortical areas corresponding to the middle and posterior MFG. The patient produced errors in 2 of 3 stimulations for a total number of 10 verb picture stimuli.

However, MG failed to respond to 3 stimuli in the no-DES condition. Therefore, the mapping procedure was interrupted.

Subject PR produced errors to both objects and finite verbs when DES was delivered to the dorsal premotor cortex, anterior to an area in the superior motor cortex where motor hand movements were affected. Object naming errors were only detected when DES was delivered to an area of prefrontal cortex anterior to the mouth area in the middle part of the premotor cortex. Errors with finite verbs were only detected when DES was delivered to a posterior part of the SFG. During subcortical DES, errors were elicited with both tasks, possibly when stimulating subcortical areas corresponding to the posterior part of the MFG and SFG. These areas were deemed significant for language processing, as errors were elicited 2/3 times during DES. It was not possible to run statistical analyses on the mapping data of this participant, as a full video of the surgical procedure could not be acquired.

Subject SO produced a significant number of errors when DES was applied to the IFG (pars opercularis), during the production of finite verbs in sentence context (2/9 errors; Wald=3.263, Nuisance=0.122;  $p=0.004^*$ ). In the same area, results of DES during object naming did not reach significance (2/4 errors; Wald=1.511, Nuisance=0.131;  $p=0.091$ ). A significant number of errors was obtained during DES, particularly with finite verbs, in a subcortical area corresponding to the most anterior-medial part of the insula, possibly corresponding to the IFOF (2/3 errors; Wald=2.653, Nuisance=0.152;  $p=0.014^*$ ).

Subject TT produced a significant number of errors during object naming when subcortical DES was applied (4/13; Wald=2.267; Nuisance=0.0454;  $p=0.013^*$ ), particularly in two subcortical areas corresponding to the MFG, close to the premotor cortex: one posterior (3/3;

Wald=4.141; Nuisance=0.161; p=0.001\*) and one inferior-posterior (1/3 errors; Wald=2.216; Nuisance=0.1212; p=0.033\*). No significant differences were observed at the cortical level.

Table 6.3


*Individual comparisons per condition.*

		<u>CRA</u>	<u>CRO</u>	<u>MG</u>	<u>SO</u>	<u>TT</u>
Cortical	S v NS	W=0.449	W=1.288	W=4.062	W=1.569	W=0.101
		Np=0.021	Np=0.04	Np=0.061	Np=0.969	Np=9.969
		p=0.374	p=0.167	p=0.001*	p=0.078	p=0.494
	S Obj v NS Obj	W=0.386	W=2.556	W=3.485	W=0.846	W=0.208
		Np=0.949	Np=0.223	Np=0.091	Np=0.939	Np=0.929
		p=0.397	p=0.006*	p=0.001*	p=0.218	p=0.454
	S Fv v NS Fv	W=0.657	NA	W=1.191	W=0.902	W=0.267
		Np=0.031*		Np=0.031	Np=0.02	Np=9.929
		p=0.321		p=0.178	p=0.272	p=0.449
	S Obj v S Fv	W=2.75	NA	W=1.103	W=0.687	W=0.716
		Np=0.414		Np=0.888	Np=0.08	Np=0.04
		p=0.004*		p=0.167	p=0.317	p=0.307
Subcortical	S v NS	W=2.646	W=0.522	NA	W=3.821	W=1.094
		Np=0.112	Np=0.07		Np=0.141	Np=0.021
		p=0.006*	p=0.392		p=0.001*	p=0.218
	S Obj v NS Obj	W=1.922	W=0.522	NA	W=1.654	W=2.267
		Np=0.223	Np=0.07		Np=0.394	Np=0.454
		p=0.037*	p=0.392		p=0.091	p=0.013*
	S Fv v NS Fv	W=1.771	NA	NA	W=3.442	W=0.716
		Np=0.112			Np=0.192	Np=0.939
		p=0.051			p=0.002*	p=0.277
	S Obj v S Fv	W=0.127	NA	NA	W=0.422	W=0.716
		Np=0.787			Np=0.323	Np=0.939
		p=0.492			p=0.418	p=0.277

*Notes.* Statistics per individual area are reported in the results. S = DES, NS = no-DES; Obj = object naming; FV = production of finite verbs in sentence context. Wald = Wald statistic, Np= Nuisance parameter, p= p-value (two-tailed).SFG = Superior frontal gyrus, MFG = Middle frontal gyrus; IFG = Inferior frontal gyrus; STG =Superior temporal gyrus; MTG = Middle temporal gyrus; ITG = Inferior temporal gyrus; IPL = Inferior parietal lobe. NA = Not applicable because DES in one of the conditions was not possible. Statistical analyses were not possible for patient PR. Significant results per area are indicated in text.

### 6.3.2. Error analyses

The cortical (6.2A) and subcortical (6.2B) location of the various types of error produced by each subject are indicated in Figure 6.2.



*Figure 6.2.* Error types during DES per subject. A = cortical, B = subcortical. The number of stimulations is higher during cortical than subcortical DES in subjects CRA (68 v 17), SO (56 v 25), and TT (44 v 25). Cortical and subcortical stimulations were equal for CRO (14). Subcortical DES was not performed to MG. The number of stimulations could not be counted in PR.

Subject CRA asked for a break when responding to the test with finite verbs, while this did not happen while responding to object naming. She produced 5 semantic paraphasias in 32 stimulations (henceforth, 5/32 semantic paraphasias, where the numerator indicates the number of specific errors and the denominator indicates the total number of stimulations per test) and 3/32 anomias when the finite verb production task was administered during cortical DES. Subcortical DES elicited 2/7 semantic paraphasias and 1/7 unrelated word for object naming, and 2/10 semantic paraphasias and 4/10 anomias during the production of finite verbs.

Subject CRO produced 4/7 latencies (i.e., responses produced beyond the allowed, 4-second limit), 1/7 fragment, 1/7 anomia, and 1/7 semantic error (“*hammer*” for “*cherry*”), when stimulated in the MFG during object naming. No stimulation was applied during the finite verb production task, due to the high number of spontaneous errors (i.e., 7/12 errors), most of which were lexical-semantic: 3 anomias, 2 semantic paraphasias, 1 change of category (“*He lion*” for “*He roars*”), and 1 noun-centered circumlocution in response to a lexical verb (i.e., “*She does gymnastics*” for “*She jumps*”).

Subject MG produced phonological errors during cortical DES with object naming: 4/33 fragments (e.g., “*Ecco il pf peperone*” “*Ecco il po peperone*” for “*Here [is] the bell pepper*”), 2/33 phonological paraphasias (e.g., “*Ecco il cia caccivite* [target: *cacciavite*] for “*Here [is] the screwdriver*”), and 1/33 part/whole error (“*Here [is] the foot*, for “*Here [is] the leg*”). DES in the subcortical areas corresponding to the middle premotor cortex induced 1/3 anomia and 1/3 fragment (“*Lui stra...*” for “*Lui strappa*”, he tears).

As for subject PR, we cannot provide error counts, as only a partial video of the surgery was available. This subject produced latencies and semantic paraphasias (e.g., “*onion*” for “*tomato*”) when DES was delivered to the middle premotor cortex, during object naming.



Semantic errors and anomias were observed in object naming during subcortical DES. During cortical DES of the SFG, production of finite verbs yielded responses in which the correctly produced introductory pronoun was followed by an anomia (i.e., “*He...*” instead of “*He jumps*”). During subcortical DES of the same region, perseverative errors on the verb emerged, along with difficulty on pronoun gender (e.g., “*He..., he she combs, he combs, he she combs, he combs*” for “*She combs*”).

Subject SO produced 1/6 latencies to finite verbs, and 2/6 latencies and 1/6 semantic paraphasias (i.e., guitar for violin) to object names, when DES was delivered to the IFG. During subcortical DES to an area corresponding to the IFG (possibly involving the IFOF, before it turns medially to the caudate nucleus), he produced 1/1 fragment (i.e., *ca...* for *carciofo*, artichoke) during object naming. During finite verb production, errors resulted in 1/3 change of category (“*He hair.noun*” for “*He listens.verb*”), and 1/3 error resulting in incorrect pronoun gender followed by a semantic paraphasia (i.e., “*She writes*” for “*He licks*”).

Subject TT complained of problems with his eyes, arm/hand, and face during DES in the posterior part of the IFG and MFG, close to the premotor cortex (i.e., “I cannot see”, “I cannot keep my eyes still”; “I cannot move the arm/hand”, “I did not feel it”; “I cannot feel my face”). He produced 2 errors during cortical DES (1/25 anomia during object naming and 1/19 latency during the production of finite verbs), and 5 anomic errors during subcortical DES in an area corresponding to the posterior part of the MFG, close to the premotor cortex (3/3 during object naming and 2/7 during the production of finite verbs).

## **6.4. Discussion**

We carried out quantitative and qualitative analyses of the performance on spoken production tasks in patients undergoing awake surgery for the removal of supratentorial tumors in the left, language-dominant hemisphere. We administered an automated speech task (i.e., counting from one to ten) and two overt picture-naming tasks (i.e., object naming and production of finite verbs in sentence context). Test items for intraoperative use were selected for each individual so as to minimize spontaneous errors during surgery. We assessed whether significant differences emerged when DES was applied as compared to when it was not applied, when DES was used during object naming as opposed to finite verb production (regardless of DES site), and when DES was directed to specific perisylvian areas. In all subjects, brain sites relevant for the production of object names, of finite verbs, or of both were detected. This is in agreement with studies in which DES allowed the mapping of language processes (e.g., Ojemann & Mateer, 1979), and also with more recent investigations, specifically focused on noun and verb production (e.g., Bello et al., 2007; Corina et al., 2005; Havas et al., 2015; Lubrano et al., 2014). Critical DES sites and errors partially agreed with the current literature.

### **6.4.1. DES v no-DES (collapsing across nouns and verbs)**

Three of five subjects showed significant differences between the DES and the no-DES condition. Subject MG showed differences when DES was delivered to cortical sites, and subjects CRA and SO during subcortical DES. The figures entered in this contrast correspond to the sum total of correct and incorrect responses, collapsed across word types, tasks and brain sites. Therefore, the fact that 2/5 subjects (CRO and TT) did not show significant differences between the two conditions does not indicate that DES failed to map language areas in these

patients. Rather, it stresses that significant differences between correct and incorrect responses are found only when DES is delivered to a specific brain site, and during a specific cognitive task – hence, that language mapping in the brain requires that an appropriate task be selected to map the brain site under investigation (e.g., Fernández-Coello et al. 2013, Rofes & Miceli, 2014). In our sample, this is indicated by subjects CRA and CRO (Table 6.3.). In these two subjects, no differences were observed between the cortical DES and no-DES conditions when data were collapsed across tasks, but a significant difference appeared during finite verb production (CRA) and during object naming (CRO), respectively.

From a methodological standpoint, the mere figures of the incidence of correct and incorrect responses across cumulated tasks in the DES v no-DES condition may be misleading. Results may be biased, as performance may be affected by several factors, such as task difficulty (i.e., one task may be intrinsically more difficult than the other and trigger errors that are not related to DES), order of task administration (i.e., the second task may trigger more errors than the first, because by the time it is administered the patient is more tired), brain site stimulated (i.e., one of the two tasks may trigger more errors than the other in a specific brain area), etc. Therefore, regardless of the results of this comparison, looking at differences between tasks and across brain sites provides more relevant information than collapsing across nouns and verbs.

#### **6.4.2. Object naming and finite verbs (independently of stimulation site)**

Data confirm that DES resulted in more incorrect responses than the no-DES condition, at the cortical and/or at the subcortical level. Subjects CRO and MG produced more errors in object naming during cortical DES. Subjects CRA and TT showed the same effects during subcortical DES. Subject CRA produced significantly more errors to finite verbs than to nouns when DES

was applied at the cortical level, even though no differences between specific brain sites were observed. Subject SO showed the same tendency during subcortical DES, but the number of stimulations was too small to allow strong conclusions.

These results show that neither the object naming task nor the finite verb production task is by itself more or less appropriate for language mapping, as in different subjects one or the other can be disproportionately impaired, regardless of DES site. However, the same observation shows that a finite verb production task does allow the detection of language areas that are not responsive during object naming. Therefore, the combination of the two tasks during awake surgery provides a more adequate language map during DES than object naming alone. In our study, as in others (e.g., Havas et al., 2015; Lubrano et al., 2014), using two intraoperative tasks did not noticeably increase fatigue in the patient, nor the time needed to complete surgical procedures, as compared to object naming alone.

Differences between naming objects and producing finite verbs may be difficult to document in patients who are operated awake, as these subjects typically suffer from mild deficits and the items administered during surgery are those they named correctly in the preoperative assessment. This latter point raises the question of whether patients who score below the norm before surgery already should be considered for awake surgery. This issue is relevant, because functional language areas cannot be detected during fully asleep surgeries, and because subjects who currently are not operated awake may still benefit from an awake procedure, if intraoperative items are selected carefully. In the present study, the problem arose with CRO, who performed below the norm on both finite verbs (53% correct) and objects (70% correct) before surgery. Since her tumor affected the posterior part of the MFG, the awake surgery protocol was applied in order to spare as much critical tissue as possible. The

intraoperative use of finite verbs to map language in the brain was ruled out, because during surgery CRO produced 7/12 errors on verbs in the no-DES condition. However, language was successfully mapped by administering the object naming task. Two areas relevant for object naming were mapped in MFG, which would not have been detected if the patient had undergone a fully asleep surgery. One week after surgery, CRO's scores in object naming were similar to those obtained in preoperative assessments object naming (70% v 76% correct, Barnard's test  $z=0.527$ ,  $p=0.263$  one tailed), whereas a significant decline was observed on finite verbs (53% v 29% correct, Barnard's test  $z=2.924$ ,  $p=0.001$  one-tailed), which had not been presented during surgery. In our view, observations like this invite to increase efforts aimed at finessing the interpretation and exploitation of preoperative results, so as to set up the best possible intraoperative mapping (i.e., one that allows to map language in a minimal time even in the presence of abnormal baseline performance on language tasks).

#### **6.4.3. Object naming and finite verbs (by stimulation site)**

The cortical sites where object naming or finite verb production triggered a significant number of errors were highly variable, and in partial agreement with available lesion data (e.g., Kemmerer et al., 2012; Lubrano et al., 2014; Mätzig et al., 2009; Pillon & d'Honichtun, 2011; Rofes & Miceli, 2014; Rofes et al., 2015b). For example, patients in Lubrano et al. (2014) produced more errors when DES was applied in posterior areas of the MFG and SFG during finite verb production than during noun production. In subjects CRO and MG, significant areas were found only during object naming. They were distributed across different sites in the posterior part of the inferior, middle, and superior frontal gyri. In subject SO, a significant language site was found only in the pars opercularis of the IFG during DES with finite verbs. In subjects CRA and TT, no

significant DES sites were found during cortical DES with either language task. In subject PR, an area in the SFG could only be mapped during finite verb production, and a posterior area of the MFG elicited errors to both object names and finite verbs. Even though no specific area could be systematically tied to object naming or to finite verb production, our results confirm that nouns and verbs may be represented in partly separable neural substrates. In two subjects, only one task was affected during DES – object naming in CRA, finite verb production in MG. In subject PR, cortical DES to the dorsal premotor cortex yielded comparable results in the two tasks, but DES anterior to the middle motor cortex only interfered with object naming, whereas DES to the posterior part of SFG yielded errors in verb naming only.

Observations collected during subcortical DES are variable and do not show a specificity for object naming or finite verb tasks. Two subjects could not be mapped at all; CRA and TT could be mapped only during object naming; SO only during finite verb production, and PR during both tasks. One of the reasons for these less-than-satisfactory results is that subcortical DES is typically less exhaustive than cortical DES. When it is started, surgery has been underway for a long time already – the patient has undergone cortical DES with language monitoring during tumor resection, and is likely to be tired. Consequently, fewer brain sites can be stimulated and fewer data points can be obtained. In our sample, more sites were stimulated during cortical than subcortical DES in subjects CRA (68 cortical v 17 subcortical), SO (56 v 25), and TT (44 v 25) and the same number was stimulated in CRO (14). Only 3 subcortical stimulations were applied to MG, due to a sudden cluster of anomias in the no-DES condition, possibly due to tiredness; and stimulated sites could not be counted in PR.

At a more theoretical level, it is worth asking to what extent the fact that we used picture naming paradigms to elicit nouns and finite verbs may have influenced results. In an object

naming task, the stimulus picture typically represents the corresponding object (e.g., an apple). In verb naming tasks, however, representing the to-be-named action often imposes constraints that may influence results. For example, representing the stimulus “to roar” involves a picture that includes an agent (e.g., a lion) who is performing that action. Responding “roars” requires not only selecting the target action word but also inhibiting the production of the name “lion”. In other words, executive control mechanisms must inhibit noun selection and focus on verb retrieval, to allow producing the target verb. When these mechanisms are interfered with by DES, patients may produce errors such as those of subject CRO who said “*He lion*” for “*He roars*”, or of subject SO who produced “*He hair*” for “*He listens*”. To circumvent these problems, one may choose to only use pictures of actions that do not include objects. This could be accomplished by selecting only stimuli that represent non-manipulable nouns and verbs (e.g., Saccuman et al., 2006), or by deleting objects from the action drawings that may require them (e.g. deleting or blurring the ball in the image corresponding to the verb “to kick”). Another possibility is to use paradigms based on naming to oral description (e.g., Hamberger & Tamny, 1999). In this latter case, pictorial constraints are bypassed and the sentence describing the action may still include an agent/object (e.g., the stimulus sentence for the verb “to kick” could be “To strike with one’s foot”).

#### **6.4.4. Error analyses**

Anomias, fragments, and latencies were detected across subjects, during DES of the IFG and MFG. DES of the IFG also elicited pronoun gender errors (e.g., “*She combs*” for “*He combs*”), while DES of the MFG triggered phonological and semantic paraphasias. DES of the SFG triggered perseverations and change of pronoun gender, and DES of the IPL yielded anomic and

semantic errors. Subcortical DES elicited anomias in all cases. These resulted in failures to respond altogether, or in responses that included only the determiner in object naming (i.e., “The...”) or only the pronoun in finite verb production (i.e., “She...”). In subcortical areas corresponding to the IFG and MFG, phonological errors and fragments were elicited. Semantic paraphasias occurred during DES of subcortical areas corresponding to the IPL. DES of segments of the arcuate fasciculus elicited anomic errors in CRA and MG, and TT.

The current literature hypothesizes that dorsal networks are putatively more critical for phonological processing and ventral networks for semantic processing (e.g., Binder et al., 2009, Hickok & Poeppel, 2007; Sarubbo et al., 2015). In agreement with these claims, we found phonological errors when DES was applied to dorsal structures (e.g., to the posterior part of the MFG in subject MG), and semantic errors when DES was delivered to ventral structures (e.g., to the anterior part of the IFOF in subject SO,). Based on our overall data set and on the current literature, however, such a distinction is either too sharp or very difficult to replicate in neurosurgical studies. For example, stimulation to the prefrontal cortex elicited semantic errors in subjects CRO, PR, and SO. Similar results were found in a recent meta-analysis Sarubbo et al., 2015). Furthermore, although posterior temporal areas were not stimulated in our study, DES in the posterior part of the MTG elicited more phonemic than semantic paraphasias (Lubrano et al., 2014), and diffuse temporal lobe damage has already been reported to trigger phonological errors in the absence of semantic errors (subject DM, Caramazza et al., 2000).

The occurrence of semantic paraphasias in CRA during DES of the IPL is also at variance with some current observations. For example, DES of dorsal pathways elicited mainly phonological paraphasias (e.g., Sarubbo et al., 2015), and lesions extending to the posterior part of the arcuate fasciculus can yield phonological errors in the absence of semantic errors (e.g.



Caplan et al., 1986, Caramazza et al., 2000). In addition, Lubrano et al. (2014) reported 3/3 phonological paraphasias but no semantic paraphasias, during DES of the supramarginal gyrus in the course of object naming and action naming. Having said that, semantic errors during IPL stimulation were described in other surgical mapping studies (for review, Rofes & Miceli, 2014). These errors may be due to the involvement of the IPL in semantic processing, as suggested by neuroimaging studies (Binder et al., 2009).

Anomic errors were ubiquitous. They occurred following cortical and subcortical DES regardless of task and of brain site, in accordance to other studies (e.g., Rofes & Miceli, 2014; Sarubbo et al., 2015). This was probably because failure to name a stimulus may result from an impairment of lexical and/or semantic knowledge, and therefore result from damage to a widespread neural substrate (Gainotti et al., 1986). Finally, pronoun gender errors are difficult to interpret. They could be artifactual, and result from a list bias, since 52/70 targets in our verb task are masculine. This could be the case in subject PR, who produced a masculine pronoun instead of the feminine target (i.e., "*He combs*" for "*She combs*"). However, this account is unlikely in subject SO, who produced the reverse error type (a feminine pronoun instead of a masculine pronoun), perhaps due to a genuine difficulty with pronoun selection. Clearly, more observations are needed to reliably interpret this observation.

## **6.5. Conclusion**

An approach based on an object naming task and a finite verb production task allows the assessment of language production, within a safe time frame. Furthermore, it allows a more thorough mapping of the brain in patients with perisylvian gliomas than an object naming task alone. In subject SO, finite verb production was successfully used to map language function at

the cortical and subcortical levels, when object naming provided null results. In subject PR, it allowed the identification of language function in the SFG and MFG, and during subcortical DES. Errors varied across patients and were in partial agreement with the current literature. Further work should be directed at carrying out exhaustive neurocognitive diagnoses in subjects who are being considered for awake procedures, and at achieving a fine-grained understanding of preoperative results – patients who present with preoperative language deficits may still benefit from awake surgery. Current standards in language mapping during awake surgery can be steadily improved by interactions between neurosurgery and the neuroscience of language, and by the development of new, standardized tools.

# Chapter 7

## General discussion

This dissertation aimed to contribute to the effort to include current knowledge from aphasiology, neurolinguistics, and cognitive neuroscience in the practice of awake surgery. Awake surgery is a surgical procedure applied to people with brain tumors or pharmacologically intractable epilepsy (Berger, 1996; Duffau et al., 1999; Hunke, Van de Wiele, Itzhak, & Rubinstein, 1998; Ojemann & Mateer, 1979). As the term implies, patients are kept awake during surgery, so that they can perform language tasks during electrical stimulation of cortical and subcortical brain structures (Bello et al., 2007; Benzagmout, Gatignol, & Duffau, 2007; Duffau et al. 2003; Ojemann, Ojemann, Lettich, & Berger, 1989; Skirboll, Ojemann, Berger, Lettich, & Winn, 1996). If the patient can respond correctly to a language stimulus when an electrical pulse is applied to the brain, it is concluded that the stimulated area is not relevant for that language function, and therefore, can be safely removed without inducing postoperative language deficits (cf. Borchers, Hummelbach Logothetis, & Karnath, 2011; Desmurguet, Bonnetblanc, & Duffau, 2013). By contrast, when the application of electrical pulses to the brain repeatedly triggers errors, this is taken as an indication that the area is critically involved in the language skills probed by the task, and therefore should not be removed, as this would result in postoperative language deficits (e.g., Penfield & Boldrey, 1937; Ojemann & Mateer, 1979, Duffau, 2004).

One impetus for this dissertation was the lack of tasks and standardized materials available for and used in the assessment of people with brain tumors (e.g., Finch & Copland, 2014; Połczyńska 2009; Rofes & Miceli, 2014). Object naming is in many cases the only task used for intraoperative language mapping, and one of the few tasks regularly used to assess

language deficits before or after surgery (for a review see De Witte & Mariën, 2013). However, object naming tasks are not sufficient for thorough intraoperative testing. This is for several reasons: despite object naming being intact other language abilities can be impaired (Santini et al., 2012; Satoer et al., 2014; e.g., verb production can be impaired, Caramazza & Hillis, 1991); object naming tasks do not assess some everyday language processes such as sentence formation, reference to action, or to moments in time (e.g., Vigliocco et al., 2011); and nouns and other word categories, such as verbs may be partially segregated in the brain (e.g., Mätzig et al., 2009, Kemmerer et al., 2012).

In addition, language skills are often evaluated using informal assessments. However, failure to control for critical psycholinguistic variables that may constrain performance may yield results that are difficult to interpret. Consequently, we developed a new set of standardized tasks for awake surgery. Such tasks may contribute to improving the sensitivity of intraoperative language mapping, as well as that of preoperative and postoperative assessments in this population. The novelty of the tasks we proposed is that they are standardized; controlled for linguistic variables that affect the performance of people with aphasia; validated in people with aphasia; and have been compared to current tests used in awake surgery.

We argued that tasks thus constructed allow the examiner to obtain better control over the patient's performance, and ultimately help the neurosurgeon to reach their goals – more extensive tumor removal whilst ensuring preservation of language skills (e.g., De Witt Hammer et al., 2012; Sanai & Berger, 2009). Furthermore, information obtained with these tasks, and for example with the administration of object naming and a new task before, can contribute to more accurate language assessments before and after surgery. Moreover, it may lead to a better understanding of the neural underpinnings of language: through the use of controlled language

tasks in people with brain tumors, it may be possible to replicate results obtained in other lesion studies, as well as in work from aphasiology and neurolinguistics (e.g., Anderson, Damasio, & Tranel, 1990; Havas et al., 2015; Lubrano, Filleron, Démonet, & Roux, 2012; Moritz-Gasser & Duffau, 2013).

In this thesis we focused on the role of production tasks using finite verbs and the evaluation of their ability to assess language processing in comparison with the current gold standard (i.e., object naming). As a preliminary step in the development of novel, controlled object and verb production tasks, we critically reviewed the literature twice. First, we discussed published studies that proposed intraoperative tasks requiring the production of verbs in isolation and in sentences, and we graphically represented the stimulation sites and error types that were reported in each study (Chapter 2). Subsequently, we reviewed the tasks reported as used intraoperatively, and discussed the advantages and disadvantages of each from a cognitive and neurolinguistic perspective (Chapter 5). With this information in mind, we constructed and validated two naming tasks for native speakers of Italian undergoing awake surgery – an object naming task and a finite verb production task (Chapter 3). We administered the finite verb production task and other picture naming tasks to subjects with post-stroke aphasia, to establish which tasks showed the strongest correlation with communicative abilities in daily life (Chapter 4). Also, we introduced and compared our standardized finite verb task in the operating theatre and compared it with an object naming task (Chapter 6).

In this final section we discuss the results obtained in this dissertation with a focus on theoretical and clinical issues of importance in awake surgery.

### **8.1. Applying language knowledge to functional neurosurgery**

Chapters 2 and 5 included a review of the seminal work by Penfield in the 1930-40s, Ojemann, and Mateer in the 1970-80s, and more contemporary work of figures such as Berger, Bello, Duffau and Démonet. With few exceptions (e.g., Fernández Coello, Moritz-Gasser, Martino, Martinoni, Matsuda, & Duffau, 2013; Sanai, Mirzadeh, & Berger, 2008; Zemmoura, Herbet, Moritz-Gasser, & Duffau, 2015), contemporary practice in awake surgery focuses on production tasks, rather than comprehension or verification tasks. This approach is understandable from a neurosurgical perspective. In the most widely used production task (i.e., object naming), the patient is asked to produce the name of an object, thus providing the neurosurgeon and the neuropsychologist/clinical linguist with direct feedback on language abilities while surgery is carried out. Long response latencies, failures to respond, semantic or phonemic paraphasias, etc., all provide clear indication that linguistic skills are being interfered with.

Comprehension and verification tasks are less demanding than overt production tasks, as the patient is not required to speak. For example, in the Pyramids and Palm Trees Test (Howard & Patterson, 1992), patients see a stimulus picture/word and are asked to point to one of two pictures/words (a semantically related and a semantically unrelated alternative). In addition, in such comprehension tasks a correct response can be produced even if knowledge of the item under scrutiny is not entirely spared. For example, for an image of two hands, given the choice between a pair of gloves and a pair of shoes, the subject may respond correctly either because he knows what gloves are, or because he does not know what gloves are, but knows that shoes are not worn on hands. Furthermore, due to the intraoperative setup, only the person assessing language can see the response of the patient, whereas the neurosurgeon cannot. This may render

intraoperative comprehension tasks more cumbersome to administer and interpret, while also being less sensitive to language impairments.

As discussed in Chapters 2 and 5, and noted above, of the tasks proposed for intraoperative use, object naming is the most frequently implemented in regular practice, and the one regarded as best practice (i.e., the gold standard). The central role assigned to object naming accrues from the fact that producing the word that corresponds to a visually-presented object engages a number of cognitive and linguistic processes (e.g., Coltheart 2001; Whitworth, Webster, & Howard, 2005). It requires the ability to obtain an abstract structural representation of the stimulus, to retrieve its meaning representation and the corresponding phonological lexical information, and to program and produce the corresponding string of sounds. Since damage to any of these stages is very likely to have a disruptive effect on language production and/or comprehension (e.g., Goodglass & Wingfield, 1997), identifying the brain structures involved in these processes is necessary in order to prevent permanent post-operative damage. However, in this context an important contribution of chapters 2 and 3 to the current literature is the argument that object naming may not provide a sufficiently comprehensive evaluation of language skills, and therefore may not be sufficient to avoid language impairment during surgery.

Applying current knowledge from neurolinguistics, aphasiology, and cognitive neuroscience to tasks used during awake surgery entails spelling out the language processes engaged by each task and their neural correlates. This preliminary step should allow a principled, pre-operative evaluation of the benefits accruing from the use of a specific task in opposition (or in addition) to others. In this context, asking a patient to produce a noun when seeing a picture of an object, or a verb when seeing a picture of an action may seem to require very similar processes. This is because, *prima facie*, and regardless of the specific functional architecture of

the language system (for contrasting views see, for example, Caramazza, 1997 and Levelt, 1989), both tasks involve understanding the pictorial stimuli, accessing their meaning, finding the corresponding lexical form, and articulating it correctly. However, when the two tasks are considered more carefully from a neurolinguistic perspective, clear distinctions emerge.

Verbs, and particularly verbs in sentence context, have certain characteristics that nouns lack. For example, they entail a predicate-argument and a thematic structure (Carlson & Tanenhaus, 1989), require agreement relations with their subject (Hale & Keyser, 1998), and allow reference to time and events within a specific time frame (Bastiaanse, Bamyaci, Hsu, Lee, Yarbay Duman, & Thompson, 2011). Nouns and verbs also differ from the neural perspective. Verb processing may mostly involve the frontal and inferior-parietal lobe, while noun processing may mainly recruit neural structures in the temporal lobe (Damasio & Tranel, 1993; Lubrano, Filleron, Démonet, & Roux, 2014; Miceli, Silveri, Nocentini, & Caramazza, 1988). Therefore, unsurprisingly, verbs and nouns may be differentially affected in people with aphasia (Whitworth, Webster, & Howard, 2005; Vigliocco, Vinson, Druks, Barber, & Cappa, 2011). In light of these contrasts, we developed two standardized tasks, specific to intraoperative language mapping (Chapter 3), in order to establish whether the neurofunctional distinctions between the two word categories resulted in dissociated performance during direct cortical stimulation procedures (Chapter 6).

## **8.2. Developing new and standardized tasks in awake surgery**

At the end of Chapter 2, we argued that neurosurgery teams should carefully consider the tasks they implement. Tasks should be specifically designed for awake surgery, should be standardized for a specific language, and should be developed so as to allow an in-depth assessment of language – one that goes beyond the dichotomy of correct/incorrect performance on a specific



task or item. Some of these issues were considered and put to work in Chapter 3 where we provided new materials for language mapping in Italian: an object naming test similar to the current gold standard (e.g., DO-80; Metz-Lutz, Cremin, Deloche, Hannequin, Ferrand, Perrier, & Blavier, 1991), and a task for the production of finite verbs in sentence context.

One of the purposes of Chapter 3 was to show that standardized protocols can be set up with relatively modest resources. Hopefully, this will encourage neurosurgical teams to apply the same or similar approaches. Amongst other things, the duration of picture presentation must be considered in test preparation. This is because naming must occur during electrical stimulation, which cannot be applied for more than 4 seconds in order to prevent epileptic discharges (Duffau, 2004; Kayama, 2012). Therefore, only stimuli that healthy volunteers name in less than 4 seconds should be selected, even though from a neurolinguistic point of view a more prolonged presentation might be preferable (in fact, some studies indicate that in people with aphasia performance accuracy may peak when items are shown for 5 seconds or more; e.g., Brookshire, 1971).

In item selection, several psycholinguistic variables were considered. This was to make sure that items were easy to produce (i.e., that there was a high agreement between the expected name of the picture and the response provided by healthy volunteers), and to permit identification of critical dimensions that could be affected in each patient (e.g., low and high frequency words, transitive and intransitive verbs, objects and actions, etc.). A first objective of this procedure is to decrease the number of false positives by ensuring that not only the *items* that patients are unable to produce before surgery, but also *item categories* that patients have difficulty with preoperatively are not used intraoperatively. A second objective is to establish, before and after surgery, whether deficits in the tasks used during surgery (in this case, object

naming and finite verb production) are due to semantic, lexical, or grammatical deficits, while at the same time ruling out that differences between noun and verb production are due to a mismatch between the psycholinguistic characteristics of the two sets of stimuli. For example, in Chapter 4, we constructed four stimulus sets that were psycholinguistically matched in all respects, except that they were used for distinct tasks (i.e., producing a noun, an infinitive, a finite verb, or a noun-related infinitive). Keeping frequency, imageability, age of acquisition and other critical variables under control in the four experimental sets licensed the conclusion that finite verbs are more accurate predictors of language abilities in daily living than nouns.

In Chapter 6, the patients included in our sample did not show preoperative deficits for specific psycholinguistic variables. Therefore, a personal intraoperative test could be prepared by removing the items that a particular patient was unable to respond to, without needing to control for specific psycholinguistic variables (frequency, imageability, etc.). This lack of psycholinguistic effects - which contrasts with the effects often reported in other brain-damaged populations (e.g., Jonkers & Bastiaanse, 2007; Kemmerer & Tranel, 2000; Kittredge, Dell, Verkuilen & Schwartz, 2008; Lambon Ralph & Howard, 2000; Luzzatti, Raggi, Piastrini, Contardi, & Pinna 2002) - could be due to the fact that preoperative deficits in our subjects were very mild. This is generally true of patients undergoing awake surgery, as the lack of severe deficits is one of the inclusion criteria for this procedure (e.g., Bello et al., 2007). Replication of the results presented in Chapter 6 and further work in people with brain tumors will help to decide whether or not controlling the items for different variables is essential in the preoperative assessments of people with brain tumors. Nevertheless, the control of the variables described in Chapter 3 and used in Chapters 4 and 6 may allow development of stimuli for intraoperative assessment also in patients with more marked naming deficits, particularly if these can be

ascribed to specific items or item variables. For these patients, it should still be possible to select a sufficient number of correct items for intraoperative use (i.e., at least 3 items per stimulation site).

While we did not consider this aspect, finer-grained results could be obtained from the same intraoperative materials by considering spoken response times (e.g., (Arévalo, 2002; Székely et al., 2005; cf. Tsigka, Papadelis, Braun, & Miceli, 2014). Reaction time measures may be a more sensitive index than mere error rates, and allow detection of mild language deficits in patients with brain tumors. If this is the case, items to which patients respond after longer delays before surgery can be removed from the intraoperative assessment, as a measure to minimize false positives. However, reaction times may also be more difficult to interpret, as they will be susceptible both to practice effects (leading to faster responses) and to potential interference from effects of anaesthesia (general slowing) or the surgical environment more generally. Consequently, our choice to use accuracy rather than reaction times is likely to lead to results that are both more straightforward to interpret and more reliable.

### **8.3. The production of finite verbs (in awake surgery)**

In Chapter 4, we contrasted the impairments displayed on four different naming tasks (i.e., object naming, producing finite verbs, action naming with infinites and verb generation) with those of a role-playing test (CADL2, Italian version; Carlomagno et al., 2013). All picture-naming tasks were found to be sensitive probes of language abilities in everyday life, as they all correlated significantly with the CADL2. Importantly, however, the finite verb production task correlated more strongly with the CADL2 than object naming. This result might reflect the intrinsic differences between verbs and nouns discussed above and in chapters 2, 3, and 5. However, it

does not necessarily entail that finite verbs tasks are more sensitive *per se* than object naming tasks. In fact, the analysis of individual results showed that some patients scored worse on object naming than in verb tasks. Therefore, it is advisable to use object naming and the production of finite verbs to assess patients with aphasia (regardless of etiology), rather than to risk missing relevant data by restricting the evaluation to object naming or finite verb production only.

Based on theoretical and empirical knowledge of the mechanisms underlying the production of verbs and nouns, in Chapter 6 we studied the neurofunctional correlates of object naming and finite verb production during surgery - in Chapter 5 we piloted some of the analyses. The novelty of this work is that we tackled three issues that typically are not addressed in the literature: the subcortical mapping of finite verbs, an intraoperative analysis of error types, and a discussion of the effects of electrical stimulation. Similar to other studies, the investigation attests to the feasibility and usefulness of an approach to language mapping that uses both nouns and finite verbs (see also Lubrano et al., 2014; Havas et al. 2015): patients produced more errors during electrical stimulation than when naming in the absence of stimulation. Furthermore, some patients produced errors only in object naming, while for others the verb tests were more sensitive.

From a neurofunctional perspective, the distinction between more fronto-parietal involvement in grammatical processes (and hence, in the production of finite verbs) and more temporal involvement in lexical processes (and hence, in the production of nouns) has been inconsistently documented (e.g., Damasio & Tranel, 1993; Mätzig et al., 2009; Vigliocco et al., 2011, also Chapters 2 and 3). We worked with six people with brain tumors during surgery. Four of these participants produced more errors during stimulation mapping for object naming. In two cases this happened when stimulation was delivered to cortical frontal areas, and in two other

cases when stimulation was delivered to subcortical areas in the inferior parietal lobe, and in the posterior part of the middle frontal gyrus. In a fifth participant with a frontal tumor, stimulation yielded errors during both object naming and finite verb production at the cortical and subcortical levels. And in a sixth participant with a lesion in the inferior and middle frontal gyri, cortico-subcortical stimulation only yielded errors during finite verb production.

Error types were inconsistent with the hypothesis that ventral networks are more involved in semantic processing and dorsal networks are more involved in phonological processing (e.g., Binder et al., 2009; Caplan, Vanier, & Baker, 1986; Caramazza, Papagno, & Rumel, 2000; Hickok & Poeppel, 2007; Sarubbo et al., 2015). Semantic paraphasias and phonological paraphasias were observed when stimulating dorsal networks (e.g., a subcortical area corresponding to the inferior parietal lobe, possibly a posterior segment of the arcuate fasciculus), and when stimulating cortical areas connected by ventral pathways, for example, as indicated by stimulation of a surgical marker of the inferior fronto-occipital fasciculus in the insula.

Similarly to Chapter 4, the results of Chapter 6 lead us to conclude that an approach to intraoperative language mapping based on at least two tasks (finite verb production and object naming) is preferable to an intraoperative assessment that only includes object naming. Neither task alone provides a satisfactory evaluation of language skills. Since both tasks probe partially distinct types of knowledge and recruit partially distinct neural substrates, their joint use during surgery guarantees a more reliable assessment of language and a greater likelihood of language sparing. Hence, in order to improve the quality and the sensitivity of language assessment during awake surgery, we believe that at least these two standardized tasks are required. Surgical teams should be aware of the theoretical implications and intraoperative advantages of each task (e.g.,

brain areas/networks where tasks may be more sensitive and specific, types of language errors that may emerge and what do they mean, what level of language is being tackled with the task and which other levels can be tested, etc.), based on current knowledge in aphasiology, neurolinguistics, and cognitive neuroscience.

#### **8.4. Conclusion**

New and more refined techniques are being applied to meet surgical needs such as removing more tumor tissue, improving survival rates, and preserving the patients' language processes and quality of life. This dissertation contributed to further the implementation of the production of finite verbs, which is a task that until now had not been routinely considered. Other than that, we strengthened the need of administering standardized tasks and comparing new tasks with gold standards.

We are currently considering pairing naming tasks with other imaging methods during surgery and in the different follow-ups (e.g., electrocorticography, encephalography). We are also considering the introduction of tasks to assess writing, comprehension, and working memory processes. Simultaneously, we are working on a fast protocol of narrative speech that could be in part analyzed semi-automatically, and used in follow-up assessments and even during surgery. This is because formal testing cannot be administered too often, as it may result in test-retest effects and time constraints in clinical practice, and also because narrative speech integrates different language levels and comes closest to language use in daily life. Above all, efforts may be directed at understanding the attributes and neural underpinnings of each task, and the benefit that it may bring to each patient, possibly directing us towards clearer answers to questions posed in aphasiology, neurolinguistics, and cognitive neuroscience.

# References

- Adhikary, S., Thiruvengkatarajan, V., Babu, K., & Tharyan, P. (2011). The effects of anaesthetic agents on cortical mapping during neurosurgical procedures involving eloquent areas of the brain. *Cochrane Database of Systematic Reviews*, 11, 1-12. doi:10.1002/14651858.CD006679.pub2
- Aftonomos, L.B, Steele, R.D., Appelbau, J.S., & Harris, V.M. (2001). Relationships between impairment-level assessments and functional-level assessments in aphasia: Findings from LCC treatment programmes. *Aphasiology*, 15, 951-964.
- Aggujaro, S., Crepaldi, D., Pistarini, C., Taricco, M., & Luzzatti, C. (2006). Neuro-anatomical correlates of impaired retrieval of verbs and nouns: Interaction of grammatical class, imageability and actionality. *Journal of Neurolinguistics*, 19, 175-194. doi:10.1016/j.jneuroling.2005.07.004
- Anderson, S.W., Damasio, H., & Tranel, D. (1990). Neuropsychological impairments associated with lesions caused by tumor or stroke. *Archives of Neurology*, 47, 397-405. doi:10.1001/archneur.1990.00530040039017
- Andersson, M., & Sandström, T. (2014). [Swedish: Intraoperative language testing through the elicitation of meaning construction: Further development of the language test procedure in neurosurgery awake based on current research, practical observations and intraoperative pilot testing]. Faculty of Medicine, Department of Neuroscience, Uppsala University, Sweden. Retrieved from: <http://www.diva-portal.org/smash/record.jsf?pid=diva2:783621>
- Arévalo, A. (2002). Teasing apart actions and objects: a picture-naming study. *Center for Research in Language Newsletter*, 14, 3-15. Retrieved from <http://crl.ucsd.edu/newsletter/14-2/article.html>

- Aten, J. L., Caligiuri, M. P., & Holland, A. L. (1982). The efficacy of functional communication therapy for chronic aphasic patients. *Journal of Speech and Hearing Disorders*, 47, 93-96. doi:10.1044/jshd.4701.93
- Bakheit, A. M., Carrington, S., Griffiths, S., & Searle, K. (2005). High scores on the Western Aphasia Battery correlate with good functional communication skills (as measured with the Communicative Effectiveness Index) in aphasic stroke patients. *Disability and Rehabilitation*, 27, 287-291. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/16040530>
- Barsalou, L. W. (1983). Ad hoc categories. *Memory & Cognition*, 11, 211-227. doi:10.3758/BF03196968
- Barton, M.I., Maruszewski, M., & Urea, D. (1969). Variation of stimulus context and its effect on word-finding ability in aphasics. *Cortex*, 5, 351-365. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/5374220>
- Bastiaanse, R., & Jonkers, R. (1998). Verb retrieval in action naming and spontaneous speech in agrammatic and anomic aphasia. *Aphasiology*, 12, 951-969. doi:10.1080/02687039808249463
- Bastiaanse, R., & Van Zonneveld, R. (2004). Broca's aphasia, verbs and the mental lexicon. *Brain and Language*, 90, 198-202. doi:10.1016/S0093-934X(03)00432-2
- Bastiaanse, R., Bamyaci, E., Hsu, C.-J., Lee, J., Yarbey Duman, T., & Thompson, C. K. (2011). Time reference in agrammatic aphasia: a cross-linguistic study. *Journal of Neurolinguistics*, 24, 652-673. doi:10.1016/j.jneuroling.2011.07.001



- Bastiaanse, R., Edwards, S., Maas, E., & Rispens, J. (2003). Assessing comprehension and production of verbs and sentences: The Verb and Sentence Test (VAST). *Aphasiology*, 17, 49-73. doi:10.1080/02687030244000455
- Bedny, M., & Caramazza, A. (2011). Perception, action, and word meanings in the human brain: The case from action verbs. *Annals of the New York Academy of Sciences*, 1224, 81-95. doi:10.1111/j.1749-6632.2011.06013.x
- Bell, B. D., Davies, K. G., Hermann, B. P., & Walters, G. (2000). Confrontation naming after anterior temporal lobectomy is related to age of acquisition of the object names. *Neuropsychologia*, 38, 83-92. doi:10.1016/S0028-3932(99)00047-0
- Bello, L., Acerbi, F., Giussani, C., Baratta, P., Taccone, P., & Songa, V. (2006). Intraoperative language localization in multilingual patients with gliomas. *Neurosurgery*, 58, 115-125. doi:00-0000/00.NEU.0000000000.00000.ac
- Bello, L., Galluci, M., Fava, M., Carraba, G., Giussani, C., Acerbi, F., et al. (2007). Intraoperative subcortical language tract mapping guides surgical removal of gliomas involving speech areas. *Neurosurgery*, 60, 67-82. doi:10.1227/01.NEU.0000249206.58601.DE
- Bello, L., Gambini, A., Castellano, A., Carraba, G., Acerbi, F., Fava, E., et al. (2008). Motor and language DTI fiber tracking combined with intraoperative subcortical mapping for surgical removal of gliomas. *Neuroimage*, 39, 369-382. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/17911032>
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B (Methodological)*, 289-300. Retrieved from: <http://www.jstor.org/stable/2346101>

- Benzagmout, M., Gatignol, P., & Duffau, H. (2007). Resection of world health organization grade II gliomas involving Broca's area: Methodological and functional considerations. *Neurosurgery*, *61*, 741-753. doi:10.1227/01.NEU.0000280069.13468.B2
- Berger, M.S. (1996). Minimalism through intraoperative functional mapping. *Clinical Neurosurgery*, *43*, 324-337. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/9247814>
- Berndt R.S., Mitchum C.C., Haendiges A.N., & Sandson J. (1997). Verb retrieval in aphasia. 1. Characterizing single word impairments. *Brain and Language*, *56*, 68-106. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/>
- Berndt, R. S., Haendiges, A. N., Burton, M. W., & Mitchum, C. C. (2002). Grammatical class and imageability in aphasic word production: Their effects are independent. *Journal of Neurolinguistics*, *15*, 353-371. doi:10.1016/S0911-6044(01)00030-6
- Bertinetto, P. M., Burani, C., Laudanna, A., Marconi, L., Ratti, D., Rolando, C., & Thornton, A. M. (2005). CoLFIS. Corpus and Frequency Lexicon of Written Italian. Retrieved from: <http://www.ge.ilc.cnr.it/strumenti.php>
- Best, W., Greenwood, A., Grassly, J., & Hickin, J. (2008). Bridging the gap: Can impairment-based therapy for anomia have an impact at the psycho-social level? *International Journal of Language and Communication Disorders*, *43*, 390-407. doi:10.1080/13682820701608001.
- Binder J.R., Swanson S.J., Hammeke T.A., & Sabsevitz D.S. (2008). A comparison of five fMRI protocols for mapping speech comprehension systems. *Epilepsia*, *49*, 1980-1997. doi:10.1111/j.1528-1167.2008.01683.x.

- Binder, J. R., Desai, R. H., Graves, W. W., & Conant, L. L. (2009). Where is the semantic system? A critical review and meta-analysis of 120 functional neuroimaging studies. *Cerebral Cortex*, 19(12), 2767-2796. doi:10.1093/cercor/bhp055
- Bird, H., Howard, D., & Franklin, S. (2000). Why is a verb like an inanimate object? Grammatical category and semantic category deficits. *Brain and Language*, 72, 246-309. doi:10.1006/brln.2000.2292
- Bizzi, A., Blasi, V., Falini, A., Ferroli, P., Cadioli, M., Danesi, U., et al. (2008). Presurgical functional MR Imaging of language and motor functions: Validation with intraoperative electrocortical mapping. *Radiology*, 248, 579-589. doi:10.1148/radiol.2482071214
- Black, M., & Chiat, S. (2003). Noun-verb dissociations: A multi-faceted phenomenon. *Journal of Neurolinguistics*, 16, 231-250. doi:10.1016/S0911-6044(02)00017-9
- Bogka N., Masterson J., Druks J., Fragkioudaki M., Chatziprokopiou E.S., & Economou, K. (2003). Object and action picture naming in English and Greek. *European Journal of Cognitive Psychology*, 15, 371-403. doi:10.1080/09541440303607
- Bookheimer, S. Y., Zeffiro, T. A., Blaxton, T. A., Gaillard, W., & Theodore, W. H. (2000). Activation of language cortex with automatic speech tasks. *Neurology*, 55, 1151-1157. doi:10.1212/WNL.55.8.1151
- Borchers, S., Himmelbach, M., Logethetis, N., & Karnath, H.O. (2011). Direct electrical stimulation of human cortex – the gold standard for mapping brain functions? *Nature Reviews Neuroscience*, 13, 63-70. doi:10.1038/nrn3140
- Borius, P. Y., Giussani, C., Draper, L., & Roux, F.E. (2012). Sentence translation in proficient bilinguals: A direct electrostimulation brain mapping. *Cortex*, 48(5), 614-622. doi:0.1016/j.cortex.2011.01.011

- Brookshire, R.H. (1971). Effects of trial time and inter-trial interval on naming by aphasic subjects. *Journal of Communication Disorders*, 3, 289-301. doi:10.1016/0021-9924(71)90035-9
- Bub, D. N., & Masson, M. E. (2012). On the dynamics of action representations evoked by names of manipulable objects. *Journal of Experimental Psychology: General*, 141, 502-517. doi:10.1037/a0026748
- Caplan D., Gow D., & Makris N. (1995). Analysis of lesions by MRI in stroke patients with acoustic-phonetic processing deficits. *Neurology*, 45, 293-8. doi:10.1212/WNL.45.2.293
- Caplan, D., Vanier, M., & Baker, C. (1986). A case study of reproduction conduction aphasia I: Word production. *Cognitive Neuropsychology*, 3, 99-128. doi:10.1080/02643298608252671
- Cappa, S. F., & Perani, D. (2003). The neural correlates of noun and verb processing. *Journal of Neurolinguistics*, 16, 183-189. doi:S0911604402000131
- Cappelletti, M., Fregni, F., Shapiro, K., Pascual-Leone, A., & Caramazza, A. (2008). Processing nouns and verbs in the left frontal cortex: A transcranial magnetic stimulation study. *Journal of Cognitive Neuroscience*, 20, 707-720. doi:10.1162/jocn.2008.20045
- Caramazza, A. (1997). How many levels of processing are there in lexical access? *Cognitive Neuropsychology*, 14, 177-208. doi:10.1080/026432997381664
- Caramazza, A., & Hillis, A. E. (1991). Lexical organization of nouns and verbs in the brain. *Nature*, 349, 788-790. doi:10.1038/349788a0
- Caramazza, A., Berndt, R. S., & Basili, A. G. (1983). The selective impairment of phonological processing: A case study. *Brain and Language*, 18, 128-174. doi:10.1006/brln.2000.2379

- Caramazza, A., Papagno, C., & Rumel, W. (2000). The selective impairment of phonological processing in speech production. *Brain and Language*, 75, 428-450. doi:10.1006/brln.2000.2379
- Carlomagno, S., Nassimbeni, S., Zannino, G., Trifelli, E., Grattarola, C., & Razzano, C. (2013). Il Communicative Activities in Daily Life (CADL-2). Dati normativi della versione in lingua italiana. *Logopedia eCcomunicazione*, 3, 285-304. Retrieved from <http://www.erickson.it/Riviste/Pagine/Scheda-Numero-Rivista.aspx?ItemId=40582>
- Carlomagno, S., Pandolfi, M., Labruna, L., Colombo, A., & Razzano, C. (2001). Recovery from moderate aphasia in the first year poststroke: Effect of type of therapy. *Archives of physical medicine and rehabilitation*, 82, 1073-1080. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11494187>
- Carlson, G. N., & Tanenhaus, M. K. (1989). Thematic roles and language comprehension. In W. Wilkins (Ed.), *Syntax and semantics: Thematic relations*. (pp. 263-300). San Diego: Academic Press
- Carroll, J. B., & White, M. N. (1973). Word frequency and age of acquisition as determiners of picture-naming latency. *Quarterly Journal of Experimental Psychology*, 25, 85-95. doi:10.1080/14640747308400325
- Catricalà, E., Della Rosa, P., Ginex, V., Mussetti, Z., Plebani, V., & Cappa, S.F. (2013). An Italian battery for the assessment of semantic memory disorders. *Neurological Sciences*, 34, 985-993. doi:10.1007/s10072-012-1181-z
- Chacko A.G., Thomas S.G., Babu K.S., Daniel R.T., Chacko G., Prabhu K., Cherian V., & Korula G. (2013). Awake craniotomy and electrophysiological mapping for eloquent area

- tumours. *Clinical Neurology and Neurosurgery*, 115, 329-34.  
doi:10.1016/j.clineuro.2012.10.022
- Chang E.F., Raygor K.P., & Berger M.S. (2015). Contemporary model of language organization: an overview for neurosurgeons. *Journal of Neurosurgery*, 122, 250-61.  
doi:10.3171/2014.10.JNS132647.
- Cicchetti, D.V. (1994). Guidelines, criteria, and rules of thumb for evaluating normed and standardized assessment instruments in psychology. *Psychological Assessment*, 6, 284-290.  
doi:10.1037/1040-3590.6.4.284
- Cipolotti, L., Healy, C., Chan, E., Bolsover, F., Lecce, F., White, M., et al. (2015). The impact of different aetiologies on the cognitive performance of frontal patients. *Neuropsychologia*, 68, 21-30. doi:10.1016/j.neuropsychologia.2014.12.025
- Coltheart, M. (2001). Assumptions and methods in cognitive neuropsychology. In B. Rapp (Ed.), *The Handbook of Cognitive Neuropsychology* (pp. 3-21). Philadelphia, PA: Psychology Press.
- Conner, C. R., Chen, G., Pieters, T. A., & Tandon, N. (2013). Category specific spatial dissociations of parallel processes underlying visual naming. *Cerebral Cortex*, 24, 2741-2750. doi:10.1093/cercor/bht130
- Conroy, P., Sage, K., & Lambon Ralph, M. A. (2009). Errorless and errorful therapy for verb and noun naming in aphasia. *Aphasiology*, 23, 1311-1337. doi:10.1080/02687030902756439
- Corina, D. P., Gibson, E. K., Martin, R., Poliakov, A., Brinkley, J., & Ojemann, G. (2005). Dissociation of action and object-naming: Evidence from cortical stimulation mapping. *Human Brain Mapping*, 24, 1-10. doi:10.1002/hbm.20063

- Craig, H. K., Hinckley, J. J., Winkelseth, M., Carry, L., Walley, J., Bardach, L., et al. (1993). Quantifying connected speech samples of adults with chronic aphasia. *Aphasiology*, 7, 155-163. doi:10.1080/02687039308249503
- Crawford, J. R., Garthwaite, P. H., & Porter, S. (2010). Point and interval estimates of effect sizes for the case-controls design in neuropsychology: Rationale, methods, implementations, and proposed reporting standards. *Cognitive Neuropsychology*, 27, 245-260. doi:10.1080/02643294.2010.513967
- Crepaldi, D., Berlingeri, M., Paulesu, E., & Luzzatti, C. (2011). A place for nouns and a place for verbs? A critical review of neurocognitive data on grammatical-class effects. *Brain and Language*, 116, 33-49. doi:10.1016/j.bandl.2010.09.005
- Crepaldi, D., Inghinoli, C., Verga, R., Contardi, A., Semenza, C., & Luzzatti, C. (2011). On nouns, verbs, lexemes, and lemmas: Evidence from the spontaneous speech of seven aphasic patients. *Aphasiology*, 25, 71-92. doi:10.1080/02687031003637114
- Damasio, A. R., & Tranel, D. (1993). Nouns and verbs are retrieved with differently distributed neural systems. *Proceedings of the National Academy of Sciences of the United States of America*, 90, 4957–4960. doi:10.1073/pnas.90.11.4957
- Damasio, H., Grabowski, T. J., Tranel, D., Hichwa, R. D., & Damasio, A. R. (1996). A neural basis for lexical retrieval. *Nature*, 380, 499-505. doi:10.1038/380499a0
- Daniele, A., Giustolisi, L., Silveri, M.C., Colosimo, C., & Gainotti, G. (1994). Evidence for a possible neuroanatomical basis for lexical processing of nouns and verbs. *Neuropsychologia*, 32, 1325-1341. doi:10.1016/0028-3932(94)00066-2

- Davis, M. H. Meunier, F., & Marslen-Wilson, W.D. (2004). Neural responses to morphological, syntactic and semantic properties of single words: An fMRI study. *Brain and Language*, 89, 439-449. doi:10.1016/S0093-934X(03)00471-1
- de Aguiar, V., Bastiaanse, R., Capasso, R., Gandolfi, M., Smania, N., Rossi, G., & Miceli, G. (2015). Can tDCS enhance item-specific effects and generalization after linguistically motivated aphasia therapy for verbs? *Frontiers in Behavioral Neuroscience*, 9, 1-17. doi:10.3389/fnbeh.2015.00190
- de Aguiar, V., Paolazzi, C., & Miceli, G. (2015). tDCS in post-stroke aphasia: The role of stimulation parameters, behavioral treatment and patient characteristics, *Cortex*, 63, 296-316. doi:10.1016/j.cortex.2014.08.015.
- De Benedictis, A., & Duffau, H. (2011). Brain Homotopy : From esoteric concept to practical surgical applications. *Neurosurgery*, 68, 1709-1723. doi:10.1227/NEU.0b013e3182124690
- De Bleser, R., & Kauschke, C. (2003). Acquisition and loss of nouns and verbs: Parallel or divergent patterns? *Journal of Neurolinguistics*, 16, 213-229. doi:10.1016/S0911-6044(02)00015-5
- De Witt Hamer, P. C., Gil Robles, S., Zwinderman, A. H., Duffau, H., & Berger, M.S. (2012). Impact of intraoperative stimulation brain mapping on glioma surgery outcome: A meta-analysis. *Journal of Clinical Oncology*, 30, 2559-2565. doi:10.1200/JCO.2011.38.4818
- De Witt Hamer, P., Moritz-Gasser, S., Gatignol, P., & Duffau, H. (2011). Is the human left middle longitudinal fascicle essential for language? A brain electrostimulation study. *Human Brain Mapping*, 32, 962-973. doi:10.1002/hbm.21082



- De Witte E., Satoer D., Robert E., Colle H., Verheyen S., Visch-Brink E., & Mariën P. (2015). The Dutch Linguistic Intraoperative Protocol: A valid linguistic approach to awake brain surgery. *Brain and Language*, 140, 35-48. doi:10.1016/j.bandl.2014.10.011.
- De Witte, E., & Mariën, P. (2013). The neurolinguistic approach to awake surgery reviewed. *Clinical Neurology and Neurosurgery*, 115, 127-145. doi:10.1016/j.clineuro.2012.09.015
- Den Ouden, D. B., Fix, S., Parrish, T. B., & Thompson, C.K. (2009). Argument structure effects in action verb naming in static and dynamic conditions. *Journal of Neurolinguistics*, 22, 196-215. doi:10.1016/j.neuroling.2008.10.004
- Den Ouden, D. B., Hoogduin, H., Stowe, L., & Bastiaanse, R. (2008). Neural correlates of Dutch Verb Second in speech production. *Brain and language*, 104(2), 122-31. doi:10.1016/j.bandl.2007.05.001
- Desmurget, M., Song, Z., Mottolese, C., & Sirigu, A. (2013). Re-establishing the merits of electrical brain stimulation. *Trends in Cognitive Neuroscience*, 17, 422-449. doi:10.1016/j.tics.2013.07.002
- doi:10.1080/02687040143000311
- Duffau, H. (2004). Cartographie fonctionnelle per-opératoire per stimulations électriques directes: Aspects méthodologiques. *Neurochirurgie*, 50(4), 474-483. doi:NCHIR-09-2004-50-4-0028-3770-101019-ART05
- Duffau, H. (2013). A new philosophy for diffuse-low grade glioma (DLGG): Oncological and functional outcomes. *Neurochirurgie*, 59, 2-8. doi:10.1016/j.neuchi.2012.11.001
- Duffau, H., Capelle, L., Denvil, D., Sichez, N., Gatignol, P., Taillandier, L., et al. (2003). Usefulness of intraoperative electrical subcortical mapping during for low-grade gliomas located within eloquent brain regions functional results in a consecutive series of 103

- patients. *Journal of Neurosurgery*, 98, 764-778. Retrieved from [www.ncbi.nlm.nih.gov/pubmed/12691401](http://www.ncbi.nlm.nih.gov/pubmed/12691401)
- Duffau, H., Capelle, L., Sichez, J., Faillot, T., Abdenmour, L., Law Koune, J. D., et al. (1999). Intra-operative direct electrical stimulations of the central nervous system: The Salpêtrière experience with 60 patients. *Acta Neurochirurgica*, 141, 1157-67. doi:10.1007/s007010050413
- Duffau, H., Lopes, M., Arthuis, F., Bitar, A., Sichez, J-P., Van Effenterre, R., et al. (2005). Contribution of intraoperative electrical stimulations in surgery of low grade gliomas: A comparative study between two series without (1985-96) and with (1996-2003) functional mapping in the same institution. *Journal of Neurology, Neurosurgery, & Psychiatry*, 76, 845-851. doi:10.1136/jnnp.2004.048520
- Duffau, H., Velut, S., Mitchell, M.C., Gatignol, P., & Capelle, L. (2004). Intra-operative mapping of the subcortical visual pathways using direct electrical stimulations. *Acta Neurochirurgica*, 146, 265-270. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/>
- Federmeier, K. D., Segal, J. B., Lombrozo, T., & Kutas, M. (2000). Brain responses to nouns, verbs and class-ambiguous words in context. *Brain*, 123, 2252–2566. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11099456>
- Fedio, P., & Van Buren, J. M. (1975). Memory and perceptual deficits during electrical stimulation in the left and right thalamus and parietal subcortex. *Brain and Language*, 2, 78-100. doi:10.1016/S0093-934X(75)80056-3
- Fernández Coello, A., Moritz-Gasser, S., Martino, J., Martinoni, M., Matsuda, R., & Duffau, H. (2013). Selection of intraoperative tasks for awake mapping based on relationships between

- tumor location and functional networks. *Journal of Neurosurgery*, 119, 1380-1394. doi:10.3171/2013.6.JNS122470.
- Finch, E., & Copland, D. A. (2014). Language outcomes following neurosurgery for brain tumours: A systematic review. *NeuroRehabilitation*, 34, 499-514. doi:10.3233/NRE-141053
- Finocchiaro, C., & Miceli, G. (2002). Verb actionality in aphasia: Data from two aphasic subjects. *Folia Linguistica*, 36, 335–357. doi:10.1515/flin.2002.36.3-4.335.
- Finocchiaro, C., Basso, G., Giovenzana, A., & Caramazza, A. (2010). Morphological complexity reveals verb-specific prefrontal engagement. *Journal of Neurolinguistics*, 23, 553-563. doi:0.1016/j.jneuroling.2010.04.004
- FitzGerald, D.B., Cosgrove, G.R., Ronner, S., Jiang, H., Buchbinder, B.R., Belliveau, J. W., et al. (1997). Location of language in cortex: A comparison between functional MR imaging and electrocortical stimulation. *American Journal of Neuroradiology*, 18, 1529-39. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/9296196>
- Foki, T., Gartus, A., Geissler, A., & Beisteiner, R. (2008) Probing overtly spoken language at sentential level – A comprehensive high-field BOLD-fMRI protocol reflecting everyday demands. *Neuroimage*, 39, 1613-1624. doi:10.1016/j.neuroimage.2007.10.020
- Forsythe, A., Mulhern, G., & Sawey, M. (2008). Confounds in pictorial sets: The role of complexity and familiarity in basic-level picture processing. *Behavior Research Methods*, 40, 116-129. doi:10.3758/BRAI.40.1.116
- Fromm, D., & Holland, A. L. (1989). Functional communication in Alzheimer's disease. *Journal of Speech and Hearing Disorders*, 54, 535-540. doi:10.1044/jshd.5404.535

- Fucetola, R., Tabor Connor, L., Perry, J., Leo, P., Tucker, F., & Corbetta, M. (2006). Aphasia severity, semantic, and depression predict functional communication in acquired aphasia. *Aphasiology*, 20, 449-461. doi:10.1080/02687030500390177
- Gainotti G., Caltragione C., & Miceli G. (1978). Immediate visual-spatial memory in hemisphere-damage patients impairment of verbal coding and of perceptual processing. *Neuropsychologia*, 16, 501-507. doi:10.1016/0028-3932(78)90074-X.
- Gainotti, G., Silveri, M. C., Villa, G., & Miceli, G. (1986). Anomia with and without lexical comprehension disorders. *Brain and Language*, 29, 18-33. doi:10.1016/0093-934X(86)90031-3
- Garrett, M. F. (1976). Syntactic processes in sentence production. In R. J. Wales & E. C. T. Walker (Eds.), *New approaches to language mechanisms* (pp. 231-256). Amsterdam: North-Holland.
- Gentner, D. (1982). Why nouns are learned before verbs: Linguistic relativity versus natural partitioning. In S. Kuczaj II (Ed.), *Language development, Volume 2: Language, thought and culture* (pp. 301-334). Hillsdale, N.J.: Lawrence Erlbaum.
- Gibson, J.J. (2013). *The ecological approach to visual perception: Classic Edition*. Hove, East Sussex: Psychology Press; 2013
- Gil-Robles, S., & Duffau, H. (2010). Surgical management of World Health Organization Grade II gliomas in eloquent areas: The necessity of preserving a margin around functional structures. *Neurosurgical Focus*, 28, 1-9. doi:10.3171/2009.12.FOCUS09236.
- Giussani, C., Roux, F.E., Bello, L., Lauwers-Cances, V., Papagno, C., Gaini, S. M., et al. (2009). Who is who: Areas of the brain associated with recognizing and naming famous faces: Clinical article. *Journal of Neurosurgery*, 110, 289-299. doi:10.3171/2007.8.17566

- Giussani, C., Roux, F.E., Ojemann, J., Sganzerla, E.P., Pirillo, D., & Papagno, C. (2010). Is preoperative functional magnetic resonance imaging reliable for language areas mapping in brain tumor surgery? Review of language functional imaging and direct cortical stimulation correlation studies. *Neurosurgery*, 66, 113-120. doi:10.1227/01.NEU.0000360392.15450.C9
- Goldberg, E., & Goldfarb, R. (2005). Grammatical category ambiguity in aphasia. *Brain and Language*, 95, 293-303. doi:10.1016/j.bandl.2005.01.01
- Goldstein, B., Obrzut, J.E., John, C., Hunter, J.V., & Armstrong, C.L. (2004). The impact of low-grade brain tumors on verbal fluency performance. *Journal of Clinical and Experimental Neuropsychology*, 26, 750-758. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/>
- Goodglass, H., & Kaplan, E. (1972). *The assessment of aphasia and related disorders*. Philadelphia: Lea & Febiger.
- Goodglass, H., & Kaplan, E. (1983). *Boston diagnostic aphasia examination booklet*. Lea & Febiger.
- Goodglass, H., & Stuss, D.T. (1979). Naming to picture versus description in three aphasic subgroups. *Cortex*, 15, 199-211. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/477337>
- Goodglass, H., & Wingfield, A. (1997). Word-finding deficits in aphasia: Brain-behavior relations and clinical symptomatology. In H. Goodglass, & A. Wingfield (Eds.), *Anomia: Neuroanatomical and Cognitive Correlates* (pp. 3-27). San Diego, CA: Academic Press.
- Goodglass, H., Klein, B., Carey, P., & Jones, K. (1966). Specific semantic word categories in aphasia. *Cortex*, 2, 74-89. doi:10.1016/S0010-9452(66)80029-1

- Goral, M., & Kempler, D. (2009). Training verb production in communicative context: Evidence from a person with chronic non-fluent aphasia. *Aphasiology*, 23(12), 1383-1397. doi:10.1080/02687030802235203
- Hale, K., & Keyser, S. J. (1998). The basic elements of argument structure. *MIT Working Papers in Linguistics*, 32, 73-118. Retrieved from [http://web.mit.edu/~linguistics/events/tributes/hale/papers/Ch.1\\_Basic\\_Elements.3.pdf](http://web.mit.edu/~linguistics/events/tributes/hale/papers/Ch.1_Basic_Elements.3.pdf)
- Hamberger, M. J. & Tamny, T. R. (1999). Auditory naming and temporal lobe epilepsy. *Epilepsy Research*, 35, 229-243. doi:10.1016/S0920-1211(99)00023-6
- Hamberger, M., McClelland, S., McKhann, G. M., Williams, A. C., & Goodman, R. R. (2007). Distribution of auditory and visual naming sites in nonlesional temporal lobe epilepsy patients and patients with space-occupying temporal lobe lesions. *Epilepsia*, 48, 531-538. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/17326797>
- Hamberger, M., Seidel, W., Goodman, R. R., Perrine, K., & McKhann, G. M. (2003). Temporal lobe stimulation reveals anatomic distinction between auditory naming processes. *Neurology*, 60, 1478-1483. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12743235>
- Hamberger, M., Seidel, W., McKhann, G. M., Perrine, K., & Goodman, R. R. (2005). Brain stimulation reveals critical auditory naming cortex. *Brain*, 128, 2742-2749. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/16109751>
- Hamberger, M.J. (2015). Object naming in epilepsy and epilepsy surgery. *Epilepsy and Behavior*, 15, 27-33. doi:10.1016/j.yebeh.2014.12.019.
- Hart J.Jr., Berndt, R.S., & Caramazza, A. (1985) Category-specific naming deficit following cerebral infarction. *Nature*, 316, 439-440. doi:10.1038/316439a0

- Hartsuiker, R. J., & Barkhuysen, P. N. (2006). Language production and working memory: The case of subject-verb agreement. *Language and Cognitive Processes*, 21, 181-204. doi:10.1080/01690960400002117
- Hauk, O., Johnsrude, I., & Pulvermüller, F. (2004). Somatotopic representation of action words in human motor and premotor cortex. *Neuron*, 41, 301-307. doi:10.1016/S0896-6273(03)00838-9
- Havas, V., Gabarrós, A., Juncadella, M., Rifa-Ros, X., Plans, G., Acebes, J. J., et al. (2015). Electrical stimulation mapping of nouns and verbs in Broca's area. *Brain and Language*, 145, 53-63. doi:10.1016/j.bandl.2015.04.005
- Herbert, R., Hickin, J., Howard, D., Osborne, F., & Best, W. (2008). Do picture-naming tasks provide a valid assessment of lexical retrieval in conversation in aphasia? *Aphasiology*, 22, 184-203. doi:10.1080/02687030701262613
- Herbet, G., Lafargue, G., Moritz-Gasser, S., Bonnetblanc, F., & Duffau, H. (2014). Interfering with the neural activity of mirror-related frontal areas impairs mentalistic inferences. *Brain Structure and Function*, 220, 2159-2169. doi:10.1007/s00429-014-0777-x
- Herholz, K., Reulen, H. J., von Stockhausen, H. M., Thiel, A., Ilmberger, J., Kessler J., et al. (1997). Preoperative activation and intraoperative stimulation of language-related areas in patients with glioma. *Neurosurgery*, 41, 1253-1262. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/9402576>
- Hernandez-Pavon, J. C., Mäkelä, N., Lehtinen, H., Lioumis, P., & Mäkelä, J. P. (2014). Effects of navigated TMS on object and action naming. *Frontiers in Human Neuroscience*, 8, 1-9. doi:10.3389/fnhum.2014.00660

- Hickok, G., & Poeppel, D. (2007). The cortical organization of speech processing. *Nature Reviews Neuroscience*, 8, 393-402. doi:10.1038/nrn2113
- Hillis, A., & Caramazza, A. (1995). Representation of grammatical knowledge in the brain. *Journal of Cognitive Neuroscience*, 7, 369–407. doi:10.1162/jocn.1995.7.3.396
- Hillis, A.E., Tuffiash, E., & Caramazza, A. (2002). Modality-specific deterioration on naming verbs in nonfluent, primary progressive aphasia. *Journal of Cognitive Neuroscience*, 14, 1099-1108. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12419132>
- Holland, A. L. (1980). *Communicative Abilities in Daily Living* (CADL). Baltimore: University Park Press.
- Holland, A., Frattali, C., & Fromm, D. (1998). *Communication Activities of Daily Living* (CADL-2). Austin, TX: Pro-Ed.
- Howard, D., Patterson, K., Wise, R., Brown, W.D., Firston, K., Weiller, C., & Frackowiak, R. (1992). The cortical localization of the lexicons – positron emission tomography evidence. *Brain*, 115, 1769-1782, doi:10.1093/brain/115.6.1769
- Howard, D., Swinburn, K., & Porter, G. (2010). Putting the CAT out: what the Comprehensive Aphasia Test has to offer. *Aphasiology*, 24, 56-74. doi:10.1080/02687030802453202
- Huber, W., Poeck, K., & Willmes, K. (1983). The Aachen Aphasia Test. *Advances in Neurology*, 42, 291-303. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/6209953>
- Huncke, K., Van de Wiele, B., Itzhak, F., & Rubinstein, E. (1998). The asleep-awake-asleep anesthetic technique for intraoperative language mapping. *Neurosurgery*, 42, 1-9. doi:10.1097/00006123-199806000-0006
- Hurkmans J., Jonkers R., Boonstra A.M., Stewart R.E., & Reinders-Messelink H.A. (2012). Assessing the treatment effects in apraxia of speech: Introduction and evaluation of the



- Modified Diadochokinesis Test. *International Journal of Language and Communication Disorders*, 47, 427-36. doi:10.1111/j.1460-6984.2012.00155.x.
- Jacobs, B.J. (2001). Social validity of changes in informativeness and efficiency of aphasic discourse following linguistic specific treatment (LST). *Brain and Language*, 78, 115-127. doi:10.1006/brln.2001.2452
- Jonkers, R., & Bastiaanse, R. (2007). Action naming in anomic aphasic speakers: Effects of instrumentality and name relation. *Brain and language*, 102(3), 262-72. doi:10.1016/j.bandl.2007.01.002
- Katz, R.C., Hallowell, B., Code, C., Armstrong, E., Roberst, P., & Pound, C. et al. (2000). A multinational comparison of aphasia management practices. *International Journal of Language & Communication Disorders*, 35, 303–314. doi:10.1080/136828200247205
- Kayama, T. (2012). The guidelines for awake craniotomy: Guidelines committee of the Japan awake surgery conference. *Neurologia Medico-Chirurgica*, 52, 119-141. Retrieved from [https://www.jstage.jst.go.jp/article/nmc/52/3/52\\_3\\_119/\\_pdf](https://www.jstage.jst.go.jp/article/nmc/52/3/52_3_119/_pdf)
- Kemmerer, D., & Tranel, D. (2000). Verb retrieval in brain-damaged subjects : 1. analysis of stimulus, lexical, and conceptual factors. *Brain and Language*, 73, 347-392. doi:10.1006/brln.2000.2311
- Kemmerer, D., Rudrauf, D., Manzel, K., & Tranel, D. (2012). Behavioral patterns and lesion sites associated with impaired processing of lexical and conceptual knowledge of actions. *Cortex*, 48, 826–848. doi:10.1016/j.cortex.2010.11.001.
- Kertesz, A. (1982). *The Western Aphasia Battery*. New York: Grune & Stratton.
- Kertesz, A., & McCabe, P. (1977). Recovery patterns and prognosis in aphasia. *Brain*, 100, 1-18. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/861709>

- Kiefer, M., & Pulvermüller, F. (2012). Conceptual representations in mind and brain: Theoretical developments, current evidence and future directions. *Cortex*, 48, 805-825. doi:10.1016/j.cortex.2011.04.006
- Kilbride, R.D. (2013). Intraoperative Functional Cortical Mapping of Language. *Journal of Clinical Neurophysiology*, 30, 591-596. doi:10.1097/01.wnp.0000436900.48243.9f
- Kristo G., Raemaekers M., Rutten G.J., de Gelder B., & Ramsey N.F. (2015). Inter-hemispheric language functional reorganization in low-grade glioma patients after tumour surgery. *Cortex*, 64, 235-48. doi:10.1016/j.cortex.2014.11.002
- Leclercq, D., Duffau, H., Delmaire, C., Capelle, L., Gatignol, P., Ducros, et al. (2010). Comparison of diffusion tensor imaging tractography of language tracts and intraoperative subcortical stimulations. *Journal of Neurosurgery*, 112, 503-511. doi:10.3171/2009.8.JNS09558
- Levelt, W. J. M. (1989). *Speaking: From intention to articulation*. Cambridge, MA: MIT Press.
- Lomas, J., Pickard, L., Bester, S., Elbard, H., Finlayson A., & Zoghaib, C. (1989). The communicative effectiveness index: Development and psychometric evaluation of a functional communication measure for adult aphasia. *Journal of Speech and Hearing Disorders*, 54, 113-124. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/2464719>
- Lotto, M., & Schubert, A. (2008). Anesthesia for epilepsy surgery. In H. Lüders (Ed.), *Textbook of Epilepsy Surgery* (pp. 921-930). London: Informa UK Ltd.
- Lubrano, V., Filleron, T., Démonet, J.F., & Roux, F.E. (2014). Anatomical correlates for category-specific naming of objects and actions: A brain stimulation mapping study. *Human Brain Mapping*, 35, 429-433. doi:10.1002/hbm.22189.

- Lubrano, V., Roux, F.E., & Démonet, J.F. (2004). Writing-specific sites in frontal areas: A cortical stimulation study. *Journal of Neurosurgery*, 101, 787-798. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/15540917>
- Luzzatti, C., Raggi, R., Zonca, G., Pistarini, C., Contardi, A., & Pinna G.D. (2002). Verb-noun double dissociation in aphasic lexical impairments: The role of word frequency and imageability. *Brain and Language*, 444, 432-444. doi:10.1006/brln.2001.2536
- Magrassi L., Bongetta D., Bianchini S., Berardesca M., & Arienta C. (2010). Central and peripheral components of writing critically depend on a defined area of the dominant superior parietal gyrus. *Brain Research*, 30, 145-54. doi:10.1016/j.brainres.2010.05.046.
- Marshall, J.C., & Newcombe, F. (1996). Syntactic and semantic errors in paralexia. *Neuropsychologia*, 4(2), 169–176. doi:10.1016/0028-3932(66)90045-5
- Mätzig, S., Druks, J., Masterson, J., & Vigliocco, G. (2009). Noun and verb differences in picture naming: Past studies and new evidence. *Cortex*, 45, 738–758. doi:10.1016/j.cortex.2008.10.003
- Measso, G., Cavarzeran, F., Zappalà, G., Lebowitz, B.D., Crook, T.H., Pirozzolo, F.J., et al. (1993). The mini-mental state examination: Normative study of an Italian random sample. *Developmental Neuropsychology*, 9, 77-85. doi:10.1080/87565649109540545
- Mehdorn, H.M., Goebel, S., & Nabavi, A. (2013). Direct Cortical Stimulation and fMRI. In S. Ulmer, & O. Jansen (Eds.), *fMRI – Basics and Clinical Applications* (pp. 169-175). Berlin Heidelberg: Springer-Verlag.
- Mesulam, M.M. (1990). Large-scale neurocognitive networks and distributed processing for attention, language, and memory. *Annals of Neurology*, 28, 597-613. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/2260847>

- Metz-Lutz, M.N., Kremin, G., Deloche, G., Hannequin, D., Ferrand, I., Perrier, et al. (1991). Standardisation d'un test de dénomination orale: Contrôle des effets de l'âge, du sexe et du niveau de scolarité chez les sujets adultes normaux. *Revue de Neuropsychologie*, 1, 73-95. Retrieved from [http://rnp.resodys.org/IMG/pdf/Standardisation\\_d.pdf](http://rnp.resodys.org/IMG/pdf/Standardisation_d.pdf)
- Miceli, G., Capasso, R., Laudanna, A., & Burani, C. (1994). *Batteria per l'analisi dei deficit afasici* (Battery for the Evaluation of Aphasic Disorders). CEPSAG, Roma.
- Miceli, G., Capasso, R., Monti, A., Santini, B., & Talacchi, A. (2012). Language testing in brain tumor patients. *Journal of Neurooncology*, 108, 247-252. doi:10.1007/s11060-012-0810-y
- Miceli, G., Laudanna, A., & Capasso, R. (2006) *Batteria per l'analisi dei deficit afasici, Revised*. (Battery for the Evaluation of Aphasic Disorders). EMS, Bologna.
- Miceli, G., Mazzucchi, A., Menn, L., & Goodglass, H. (1983). Contrasting cases of Italian agrammatic aphasia without comprehension disorder. *Brain and Language*, 19, 65-97. doi:10.1016/0093-934X(83)90056-1
- Miceli, G., Silveri, C., Nocentini, U., & Caramazza, A. (1988). Patterns of dissociation in comprehension and production of nouns and verbs. *Aphasiology*, 2(3/4), 351-358. doi:10.1080/02687038808248937
- Miceli, G., Silveri, C., Villa, G., & Caramazza, A. (1984). On the basis for the agrammatic's difficulty in producing main verbs. *Cortex*, 20, 207-220. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/6204813>
- Miceli, G., Silveri, M. C., Romani, C., & Caramazza, A. (1989). Variation in the pattern of omissions and substitutions of grammatical morphemes in the spontaneous speech of so-called agrammatic patients. *Brain and Language*, 36, 447-492. doi:10.1016/0093-934X(89)90079-5

- Miceli, G., Silvieri, C., Nocentini, U., & Caramazza, A. (1988). Patterns of dissociation in comprehension and production of nouns and verbs. *Aphasiology*, 2(3/4), 351-358. doi:10.1080/02687038808248937
- Miceli, G., Silvieri, C., Villa, G., & Caramazza, A. (1984). On the basis for the agrammatic's difficulty in producing main verbs. *Cortex*, 20, 207-220. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/6204813>
- Moret-Tatay, C., & Perea, M. (2011). Do serifs provide an advantage in the recognition of written words? *Journal of Cognitive Psychology*, 23, 619-624. doi:10.1080/20445911.2011.546781
- Moritz-Gasser, S, & Duffau, H. (2013). The anatomo-functional connectivity of word repetition: Insights provided by awake brain tumor surgery. *Frontiers in Human Neuroscience*, 7, 1-4. doi:10.3389/fnhum.2013.00405.
- Moritz-Gasser, S., Herbet, G., Maldonado, I.L., & Duffau, H. (2012). Lexical access speed is significantly correlated with the return to professional activities after awake surgery for low-grade gliomas. *Journal of Neurooncology*, 107, 633-641. doi:10.1007/s11060-011-0789-9
- Moritz-Gasser, S., Herbet, G., & Duffau, H. (2013). Mapping the connectivity underlying multimodal (verbal and non-verbal) semantic processing: A brain electrostimulation study. *Neuropsychologia*, 51, 1814-1822. doi:10.1016/j.neuropsychologia.2013.06.007.
- Morrison, C.M., Ellis, A.W., & Quinlan, P.T. (1992). Age of acquisition, not word frequency, affects object naming, not object recognition. *Memory & Cognition*, 20(6), 705-714. doi:10.1016/j.actpsy.2008.05.005

- Motomura, K., Fujii, M., Maesawa, S., Kuramitsu, S., Natsume A., & Wakabayashi, T. (2014). Association of dorsal inferior frontooccipital fasciculus fibers in the deep parietal lobe with both reading and writing processes: A brain mapping study. *Journal of Neurosurgery*, 121, 142-148, doi:10.3171/2014.2.JNS131234
- Nabavi, A., Stark, A.M, Dörner, L., & Mehdorn, H.M. (2012). Surgical navigation with intraoperative imaging: Special operating room concepts. In A. Quiñones-Hinojosa (Ed.), *Schmidek and Sweet: Operative neurosurgical techniques: Indications, Methods, and Results* (Expert Consult-Online and Print). Elsevier Health Sciences.
- Nicholas, L.E., & Brookshire, R.H. (1993). A system for quantifying the informativeness and efficiency of the connected speech of adults with aphasia. *Journal of Speech, Language, and Hearing Research*, 36, 338-350. doi:10.1044/jshr.3602.338
- Nickels, L., & Howard, D. (1995). Aphasic naming: What matters? *Neuropsychologia*, 33, 1281-1303. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/8552229>
- Nickels, L., & Howard, D. (2004). Dissociating effects of number of phonemes, number of syllables, and syllabic complexity on word production in aphasia: It's the number of phonemes that counts. *Cognitive Neuropsychology*, 21, 57-78. doi:10.1080/02643290342000122
- Ojemaann, G., & Mateer, C. (1979). Human language cortex: Localization of memory, syntax, and sequential motor-phoneme identification systems. *Science*, 205(4413), 1401-1403. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/472757>
- Ojemann, G., Ojemann, J., Lettich, E., & Berger, M. (1989). Cortical language localization in left, dominant hemisphere. *Journal of Neurosurgery*, 108, 411-421. doi:10.3171/JNS/2008/108/2/0411

- Ojemann, J., Ojemann, G., & Lettich, E. (2002). Cortical stimulation mapping of language cortex by using a verb generation task: Effects of learning and comparison to mapping based on object naming. *Journal of Neurosurgery*, 97(1), 33-38. doi:10.3171/jns.2002.97.1.0033
- Ojemann, G.A. (1978). Organization of short-term verbal memory in language areas of human cortex: Evidence from electrical stimulation. *Brain and Language*, 5, 331-340. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/656902>
- Ojemann, G.A., & Dodrill, C.B. (1985). Verbal memory deficits after left temporal lobectomy for epilepsy. Mechanism and intraoperative prediction. *Journal of Neurosurgery*, 62, 101-7. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/3964840>
- Oliveri, M., Finocchiaro, C., Shapiro, K., Gangitano, M., Caramazza, A., & Pascual-Leone, A. (2004). All talk and no action: A transcranial magnetic DES study of motor cortex activation during action word production. *Journal of Cognitive Neuroscience*, 16, 374-381. doi:10.1162/089892904322926719
- Orsini, A., Grossi, D., Capitani, E., Laiacona, M., Papagno, C., & Vallar, G. (1987). Verbal and spatial immediate memory span: Normative data from 1355 adults and 1112 children. *Italian Journal of Neurological Sciences*, 8, 539-548. doi:10.1007/BF02333660
- Papagno, C. (2011). Naming and the role of the uncinate fasciculus in language function. *Current Neurology and Neuroscience Reports*, 11, 553-559. doi:10.1007/s11910-011-0219-6
- Papagno, C., Casarotti, A., Comi, A., Gallucci, M., Riva, M., & Bello, L. (2012). Measuring clinical outcomes in neuro-oncology. A battery to evaluate low-grade gliomas (LGG). *Journal of Neurooncology*, 108, 269-275. doi:10.1007/s11060-012-0824-5

- Papagno, C., Galluci, M., Casarotti, A., Castellano, A., Falini, A., Fava, E., et al. (2011). Connectivity constraints on cortical reorganization of neural circuits involved in object naming. *Neuroimage*, 55, 1306-1313. doi:10.1016/j.neuroimage.2011.01.005
- Patterson, K. E., & Shewell, C. (1987). Speak and spell: Dissociations and word-class effects. In M. Coltheart, R. Job, & G. Sartori (Eds). *The Cognitive Neuropsychology of language*. (pp.273-294). Hillsdale: Lawrence Erlbaum.
- Penfield, W., & Boldrey, E. (1937). Somatic motor and sensory representation in the cerebral cortex of man as studied by electrical stimulation. *Brain: A Journal of Neurology*, 60, 389-433. doi:10.1093/brain/60.4.389
- Picht, T., Krieg, S.M., Sollmann, N., Rösler, J., Niraula, B., Neuvonen, T., et al. (2013). A comparison of language mapping by preoperative navigated transcranial magnetic stimulation and direct cortical stimulation during surgery. *Neurosurgery*, 72, 808-819. doi:10.1227/NEU.0b013e3182889e01
- Pillon, A., & d'Honinckun, P. (2011). The organization of the conceptual system. The case of the “object versus action” dimension. *Cognitive Neuropsychology*, 27, 587-613. doi:10.1080/02643294.2011.609652
- Pinker, S. & Prince, A. (1994). Regular and irregular morphology and the psychological status of rules of grammar. In S. D. Lima, R. L., Corrigan, & G. K. Iverson (Eds.), *The reality of linguistic rules*. Philadelphia: John Benjamins.
- Połączyńska, M. (2009). New tests for language mapping with intraoperative electrical stimulation of the brain to preserve language in individuals with tumors and epilepsy: A preliminary follow-up study. *Poznań Studies in Contemporary Linguistics*, 45, 261-279. doi:10.2478/v10010-009-0015-5



- Porter, G., & Howard, D. (2004). *CAT: comprehensive aphasia test*. Psychology Press
- Prins, R., & Bastiaanse, R. (2004). Analysing the spontaneous speech of aphasic speakers. *Aphasiology*, 18, 1075-1091. doi:10.1080/02687030444000534
- Pulvermüller, F. (2005). Brain mechanisms linking language and action. *Nature Reviews Neuroscience*, 6, 576-582. doi:1038/nrn1706
- Pulvermüller, F., & Shtyrov, Y. (2009). Spatiotemporal signatures for large-scale synfire chains for speech processing as revealed by MEG. *Cerebral Cortex*, 19, 79-88. doi:10.1093/cercor/bhn060.
- Pulvermüller, F., Neninger, B., Elbert, T., Mohr, B., Rockstroh, R., Koebbel, P., et al. (2001). Constraint-induced therapy of chronic aphasia after stroke. *Stroke*, 32, 1621-1626. doi:10.1161/01.STR.32.7.1621
- Riddoch, M. J., & Humphreys, G. W. (1993). *BORB: Birmingham object recognition battery*. LEA.
- Rochon, E., Saffran, E. M., Berndt, R. S., & Schwartz, M. F. (2000). Quantitative analysis of aphasic sentence production: Further development and new data. *Brain and Language*, 72, 193-218. doi:10.1006/brln.1999.2285
- Rofes, A., & Miceli, G. (2014). Language mapping with verbs and sentences in awake surgery: A Review. *Neuropsychology Review*, 24, 185-99. doi:10.1007/s11065-014-9258-5
- Rofes, A., Bastiaanse, R., & Martínez-Ferreiro, S. (2014). Conditional and future tense impairment in non-fluent aphasia. *Aphasiology*, 28, 99-115. doi:10.1080/02687038.2013.850650

- Rofes, A., Capasso, R., & Miceli, G. (2015). Verb production tasks in the measurement of communicative abilities in aphasia. *Journal of Clinical and Experimental Neuropsychology*, 37, 483-502. doi:10.1080/13803395.2015.1025709
- Rofes, A., De Witte, E., Mariën, P., & Bastiaanse, R. (2012). The Verb In Sentence Context test (VISC). Rijksuniversiteit Groningen, Groningen, The Netherlands.
- Rofes, A., De Witte, E., Mariën, P., & Bastiaanse, R. (2013). Object naming may overestimate patients' language performance after neuro-oncological surgery: A case study. *Stem-Spraak- en Taalpathologie*, 18, 59-63. doi:32.8310/S01/1813-59
- Rofes, A., Spina, G., Miozzo, A., Fontanella, M., & Miceli, G. (in press). Advantages and disadvantages of intraoperative language tasks in awake surgery: A three-task approach for prefrontal tumors. *Journal of Neurosurgical Sciences*. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/26159550> [article in press]
- Rofes, A., de Aguiar, V., & Miceli, G. (2015). A minimal standardization setting for language mapping tests: An Italian example. *Neurological Sciences*, 36, 1113-9. doi:10.1007/s10072-015 2192-3
- Roux, F.E., & Trémoulet, M. (2002). Organization of language areas in bilingual patients: A cortical stimulation study. *Journal of Neurosurgery*, 97, 857-864. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12405374>
- Roux, F.E., Boukhatem, L., Draper, L., Sacko, O., & Démonet, J.F. (2009). Cortical calculation localization using electrostimulation: Clinical article. *Journal of Neurosurgery*, 110, 1291-1299. doi:10.3171/2008.8.JNS17649.
- Roux, F.E., Boulanouar, K., Lotterie, J.A., Mejdoubi, M., LeSage, J.P., & Berry, I. (2003). Language functional magnetic resonance imaging in preoperative assessment of language

- areas: Correlation with direct cortical stimulation. *Neurosurgery*, 52, 1335-1347. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12762879>
- Roux, F.E., Lubrano, V., Lauwers-Cances, V., Trémoulet, M., Mascott, C.R., & Démonet, J. F. (2004). Intra-operative mapping of cortical areas involved in reading in mono- and bilingual patients. *Brain*, 127, 1796-1810. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15240432>
- Roux, F.E., Boetto, S., Sacko, O., Chollet, F., & Trémoulet, M. (2003). Writing, calculating, and finger recognition in the region of the angular gyrus: A cortical stimulation study of Gerstmann syndrome. *Journal of Neurosurgery*, 99, 716-27. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/>
- Roux, F.E., Durand, J.B., Réhault, E., Planton S., Draper L., & Démonet, J.F. (2014). The neural basis for writing from dictation in the temporoparietal cortex. *Cortex*, 50, 64-75. doi:10.1016/j.cortex.2013.09.012
- Rueschemeyer, S.A., van Rooij, D., Lindemann, O., Willems, R.M., & Bekkering, H. (2010). The function of words: Distinct neural correlates for words denoting differently manipulable objects. *Journal of Cognitive Neuroscience*, 22, 1844-1851. doi:10.1162/jocn.2009.21310
- Ruis, C., Wajer, I.M.H, Robe, P.A., & van Zandvoort, M.J. (2014). Awake Craniotomy and Coaching. *Open Journal of Medical Psychology*, 3, 382-389. doi:10.4236/ojmp.2014.35040
- Sabsevitz, D.S., Swanson, S.J., Hammeke, T.A., Spanaki, M.V., Possing, E.T., Morris, G.L. et al. (2003). Use of preoperative functional neuroimaging to predict language deficits from epilepsy surgery. *Neurology*, 60, 1788-1792. doi:10.1212/01.WNL.0000068022.05644.01

- Saccuman, M.C., Cappa, S.F., Bates, E. A., Arévalo, A., Della Rosa, P., Danna, M., & Perani, D. (2006). The impact of semantic reference on word class: An fMRI study of action and object-naming. *Neuroimage*, 32, 1865-1878. doi:10.1016/j.neuroimage.2006.04.179
- Sach, M., Seitz, R.J., & Indefrey, P. (2004). Unified inflectional processing of regular and irregular verbs: A PET study. *Neuroreport*, 15, 533-537. doi:0.1097/01.wnr.0000113529.32218.92
- Sacko, O., Lawers-Cances, V., Brauge, D., Sesay, M., Brenner, A., & Roux, F.E. (2011). Awake craniotomy vs surgery under general anesthesia for resection of supratentorial lesions. *Neurosurgery*, 68, 1192-1199. doi:10.1227/NEU.0b013e31820c02a3
- Saffran, E.M., Berndt, R.S., & Schwartz, M.F. (1989). The quantitative analysis of agrammatic production: Procedure and data. *Brain and Language*, 37, 440–479. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/2804622>
- Sanai, N., & Berger, M. (2009). Operative techniques and the value of extent of resection. *Neurotherapeutics*, 6, 478-486. doi:10.1016/j.nurt.2009.04.005
- Sanai, N., Mirzadeh, Z., & Berger, M. (2008). Functional outcome after language mapping for glioma resection. *The New England Journal of Medicine*, 358, 18-27. doi:10.1056/NEJMoa06781
- Santini, B., Talacchi, A., Squintani, G., Casagrande, F., Capasso, R., & Miceli, G. (2012). Cognitive outcome after awake surgery for tumors in language areas. *Journal of Neurooncology*, 108, 319-326. doi:10.1007/s11060-012-0817-4
- Sarubbo, S., De Benedictis, A., Merler, S., Mandonnet, E., Balbi, S., Granieri, E., & Duffau, H. (2015). Towards a functional atlas of human white matter. *Human Brain Mapping*, 36, 3117-3116. doi:10.1002/hbm.22832

- Satoer, D, Vincent A., Dirven, C., Smits, M., Visch-Brink, E. (2011). Spontaneous speech in patients with presumed low-grade gliomas in eloquent areas before and after awake craniotomy. *Neuro-oncology*, 13, 121-126. doi:10.1093/neuonc/nor159
- Satoer, D., Work.J., Visch-Brink, E., Smits, M., Dirven, C., & Vincent, A. (2012). Cognitive functioning early after surgery of gliomas in eloquent areas. *Journal of Neurosurgery*, 117, 831-838. doi:10.3171/2012.7.JNS12263
- Shapiro, K., & Caramazza, A. (2003). Grammatical processing of nouns and verbs in left frontal cortex? *Neuropsychologia*, 41, 1189-1198. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12753958>
- Shapiro, K., & Caramazza, A. (2003). The representation of grammatical categories in the brain. *Trends in Cognitive Sciences*, 7, 201-206. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/12757821>
- Shapiro, K., Shelton, J., & Caramazza, A. (2000). Grammatical class in lexical production and morphological processing: Evidence from a case of fluent aphasia. *Cognitive Neuropsychology*, 17, 665-682. doi:10.1080/026432900750038281.
- Silveri, M. C., & Di Betta, A. M. (1997). Noun-verb dissociations in brain-damaged patients: Further evidence. *Neurocase*, 3, 477-488. doi:10.1080/13554799708405023
- Skirboll, S.S., Ojemann, G.A., Berger, M.S., Lettich, E., & Winn, H.R. (1996). Functional cortex and subcortical white matter located within gliomas. *Neurosurgery*, 38, 678-684. Retrieved from <http://ovidsp.tx.ovid.com.proxy-ub.rug.nl/sp-3.9.1a/ovidweb.cgi>
- Snodgrass, J. G., & Vanderwart, M. (1980). A standardized set of 260 pictures: Norms for name agreement, image agreement, familiarity, and visual complexity. *Journal of Experimental Psychology: Human Learning and Memory*, 6, 174-215. doi:10.1037/0278-7393.6.2.174

- Spinnler, H., & Tognoni, G. (1987). *Standardizzazione e taratura italiana di test neuropsicologici*. Masson Italia Periodici.
- Surbeck, W., Hildebrandt, G., & Duffau, H. (2015). The evolution of brain surgery on awake patients. *Acta neurochirurgica*, 157, 77-84. doi:10.1007/s00701-014-2249-8
- Székely, A., D'Amico, S., Devescovi, A., Federmeier, K., Herron, D., Iyer, G., et al. (2005). Timed action and object naming. *Cortex*, 41, 7-25. doi:10.1016/S0010-9452(08)70174-6
- Talacchi, A., Avella, D., Denaro, L., Santini, B., Meneghelli, P., Savazzi, S., et al. (2012). Cognitive outcome as part and parcel of clinical outcome in brain tumor surgery. *Journal of Neurooncology*, 108(2), 327-332. doi:10.1007/s11060-012-0818-3
- Talacchi, A., Taphoorn, M., & Miceli, G. (2012). Reconsidering outcomes that matter to patients. *Journal of Neurooncology*, 108, 219-20. doi:10.1007/s11060-012-0830-7
- Teixidor, P., Gatignol, P., Leroy, M., Masuet-Aumatell, C., Capelle, L., & Duffau, H. (2007). Assessment of verbal working memory before and after surgery for low-grade glioma. *Journal of Neurooncology*, 81, 305-313. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/16944311>
- Thompson, C. K. (2003). Unaccusative verb production in agrammatic aphasia: The argument structure complexity hypothesis. *Journal of Neurolinguistics*, 16, 151-167. doi:10.1016/S0911-6044(02)00014-3
- Thompson, C., Lange, K. L., Schneider, S. L., & Shapiro, L. P. (1997) Agrammatic and non-brain damaged subject's verb and verb argument structure production. *Aphasiology*, 11, 473-490. doi:10.1080/02687039708248485.

- Thompson-Schill, S.L., D'Esposito, M., Aguirre, G.K., & Farah, M.J. (1997). Role of left inferior prefrontal cortex in retrieval of semantic knowledge: A reevaluation. *Proceedings of the National Academy of Sciences*, 94, 14792-14797. doi:027-8424y97y9414792-6
- Tomasino, B., Marin, D., Maieron, M., D'Agostini, S., Medeossi, I., Fabbro, F., Skrap, M., & Luzzatti, C.A. (2015). A multimodal mapping study of conduction aphasia with impaired repetition and spared reading aloud. *Neuropsychologia*, 18, 214-226. doi:10.1016/j.neuropsychologia.2015.02.023
- Tranel, D., Adolphs, R., Damasio, H., & Damasio, A.R. (2001). A neural basis for the retrieval of words for actions. *Cognitive Neuropsychology*, 18, 655-674. doi:10.1080/02643290143000015
- Tranel, D., Kemmerer, D., Adolphs, R., Damasio, H., & Damasio, A.R. (2003). Neural correlates of conceptual knowledge for actions. *Cognitive Neuropsychology*, 20, 409-432. doi:10.1080/02643290244000248
- Tsapkini, K., Jarema, G. & Kehayia, E. (2002). A morphological processing deficit in verbs but not in nouns: A case study in a highly inflected language. *Journal of Neurolinguistics*, 15, 265-288. doi:10.1016/S0911-6044(01)00039-2
- Tsigka, S., Papadelis, C., Braun, C., & Miceli, G. (2014). Distinguishable neural correlates of verbs and nouns: A MEG study on homonyms. *Neuropsychologia*, 54, 87-97. doi:10.1016/j.neuropsychologia. 2013.12.018.
- Tyler, L., Bright, P., Fletcher, P., & Stamatakis, E. A. (2004). Neural processing of nouns and verbs: The role of inflectional morphology. *Neuropsychologia*, 42, 512-523. doi:10.1016/j.neuropsychologia.2003.10.001

- VanVoorhis, C.R.W., & Morgan, B.L. (2007). Understanding power and rules of thumb for determining sample sizes. *Tutorials in Quantitative Methods for Psychology*, 3, 43-50. Retrieved from <http://www.tqmp.org/Content/vol03-2/p043/p043.pdf>
- Vigliocco, G., Vinson, D., Druks, J., Barber, H., & Cappa, S. (2011). Nouns and verbs in the brain: A review of behavioural, electrophysiological, neuropsychological and imaging studies. *Neuroscience and Biobehavioral Reviews*, 35, 407-426. doi:10.1016/j.neubiorev.2010.04.007
- Wager M., Du Boisgueheneuc F., Pluchon C., Bouyer C., Stal V., Bataille B., Guillevin C.M., & Gil R. (2013). Intraoperative monitoring of an aspect of executive functions: Administration of the Stroop test in 9 adult patients during awake surgery for resection of frontal glioma. *Neurosurgery*, 72, 169-181. doi:10.1227/NEU.0b013e31827bf1d6.
- Wampold, B.E., & Freund, R.D. (1987). Use of multiple regression in counseling psychology research: A flexible data-analytic strategy. *Journal of Counseling Psychology*, 34, 372-372. doi:10.1037/0022-0167.34.4.372
- Warrington, E.K. & McCarthy, R. (1983). Category specific access dysphasia. *Brain*, 106, 859-878. doi:10.1093/brain/106.4.859
- Warrington, E.K. & Shallice, T. (1984). Category specific semantic impairments. *Brain*, 107, 829-854. doi:10.1093/brain/107.3.829
- Weitzner M.A., Meyers C.A., & Byrne K. (1996). Psychosocial functioning and quality of life in patients with primary brain tumors. *Journal of Neurosurgery*, 84, 29-34. Retrieved from: <http://www.ncbi.nlm.nih.gov/pubmed/8613832>
- Whitworth, A., Webster, J., & Howard, D. (2005). Identifying and characterizing impairments: Principles and evidence. In A. Whitworth, J. Webster, D. Howard (Eds.), *A Cognitive*



*neuropsychological approach to assessment and intervention in aphasia*. (pp. 11-22). Hove, East Sussex: Taylor & Francis.

Williams, S.E. (1983). Factors influencing naming performance in aphasia: A review of the literature. *Journal of Communication Disorders*, 16, 357-372. doi:10.1016/0021-9924(83)90018-7

Williams, S.E., & Canter, G.J. (1982). The influence of situational context on naming performance in aphasic syndromes. *Brain and Language*, 17, 92-106. doi:0093-934X/82/05009

Williams, S.E., & Canter, G.J. (1987). Action-naming performance in four syndromes of aphasia. *Brain and Language*, 32, 124-136. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/3651804>

Wilson VanVoorhis, C.R., & Morgan, B.L. (2007). Understanding power and rules of thumb for determining sample sizes. *Tutorials in Quantitative Methods for Psychology*, 3, 43-50. Retrieved from <http://www.tqmp.org/Content/vol03-2/p043/p043.pdf>

Woods, S. P., Carey, C. L., Tröster, A. I., & Grant, I. (2005). Action (verb) generation in HIV-1 infection. *Neuropsychologia*, 43, 1144-1151. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15817172>

Yokohama, S., Miyamoto, T., Riera, J., Kim, J., Akitsuki, Y., Iwata, K., et al. (2006). Cortical mechanisms involved in the processing of verbs: An fMRI study. *Journal of Cognitive Neuroscience*, 18, 1304-1313. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/16859416>

Yorkston, K. M., & Beukelman, D. R. (1980). An analysis of connected speech samples of aphasic and normal speakers. *Journal of Speech and Hearing Disorders*, 45, 27-36. doi:10.1044/jshd.4501.27

- Zacà D., Jarso S., & Pillai J.J. (2013). Role of semantic paradigms for optimization of language mapping in clinical fMRI studies. *American Journal of Neuroradiology*, 34, 1966-1971. 10.3174/ajnr.A3628.
- Zacà D., Nickerson J.P., Deib G., & Pillai J.J. (2012). Effectiveness of four different clinical fMRI paradigms for preoperative regional determination of language lateralization in patients with brain tumors. *Neuroradiology*, 54, 1015-1025. doi:10.1007/s00234-012-1056-2.
- Zemmoura, I., Herbet, G., Moritz-Gasser, S., & Duffau, H. (2015). New insights into the neural network mediating reading processes provided by cortico-subcortical electrical mapping. *Human Brain Mapping*, 36, 2215-2230. doi:10.1002/hbm.22766
- Zingeser, L. B., & Berndt, R. S. (1990). Retrieval of nouns and verbs in agrammatism and anomia. *Brain and Language*, 39, 14-32. doi:10.1016/0093-934X(90)90002-X.
- Zingeser, L.B., & Berndt, R.S. (1988). Grammatical class and context effects in a case of pure anomia: Implications of models of language production. *Cognitive Neuropsychology*, 5, 473-516. doi:10.1080/02643298808253270

## APPENDICES

### APPENDIX A: Cortico-subcortical areas where language processing was detected and error types during electrical stimulation mapping (Chapter 2)

Table A2.1

*Cortical areas in language dominant hemisphere where language processing was detected with each task and error types per area and per test*

	<u>ACTN</u>	<u>VSC</u>	<u>NOOD</u>	<u>RSEN</u>	<u>SCOMPL</u>	<u>SCOMPR</u>	<u>TRANSLP</u>	<u>VGEN</u>	<u>Error type</u>
<b>SFG</b>	Bello et al. 2006, 2007 a, c	Lubrano et al. 2014					Borius et al. 2012 g	Ojemann et al. 2002	Anomia and paraphasias
<b>MFG</b>	<b>Bello et al. 2006, 2007</b>	Lubrano et al. 2014 b, e		Roux et al. 2004 e	Ojemann & Mateer, 1979			<b>Ojemann et al. 2002 ; Roux et al. 2003</b>	Anomia and paraphasias
<b>IFG</b>	<b>Bello et al. 2006, 2007</b>	Lubrano et al. 2014 b, h		Roux et al. 2004 e	Ojemann & Mateer, 1979		Borius et al. 2012 h	<b>Ojemann et al. 2002 ; Roux et al. 2003</b>	Anomia and semantic paraphasias
<b>PrG</b>				Roux et al. 2004 f	Ojemann & Mateer, 1979			<b>Ojemann et al. 2002 ; Roux et al. 2003</b>	+
<b>PoG</b>				Roux et al. 2004 f	Ojemann & Mateer, 1979			Ojemann et al. 2002	+
<b>SMG</b>		Lubrano et al. 2014 a		Roux et al. 2004	Ojemann & Mateer, 1979	Bello et al. 2007	Borius et al. 2012	Ojemann et al. 2002	Sentence comprehension error ; Phonemic and semantic paraphasias

<b>AnG</b>				Roux et al. 2004	Ojemann & Mateer, 1979	Bello et al. 2007		Ojemann et al. 2002	Sentence comprehension error ; Phonemic and semantic paraphasias
<b>STG</b>	Bello et al. 2007	Lubrano et al. 2014 a	<b>Hamberger et al. 2003, 2007</b>		Ojemann & Mateer, 1979	Bello et al. 2007		Ojemann et al. 2002	Anomia; sentence comprehension error; semantic and phonemic paraphasias
<b>MTG</b>	Bello et al. 2007	Lubrano et al. 2014 a	<b>Hamberger et al. 2003, 2007</b>	Roux et al. 2004 e	Ojemann & Mateer, 1979			Ojemann et al. 2002	Anomia and phonemic and semantic paraphasias
<b>ITG</b>			Hamberger et al. 2003, 2007 i						Anomia
<b>Error type</b>	Anomia and semantic paraphasias	Anomia; Phonemic and semantic paraphasias, neologism, hesitation	Anomia Phonemic and semantic paraphasia	Anomia. Hesitation, phonemic and semantic paraphasias	Anomia, paraphasias, grammatical errors	Sentence comprehension error	Anomia	Anomia, hesitation, paraphasias	

*Notes.* ACTN = Action naming; VSC = Producing a finite verb in sentence context; NOOD = Naming objects to oral description; RSEN = Reading sentence aloud; SCOMPL = Sentence completion; SCOMPR = Sentence Comprehension; TRANSLP = Translating Paragraphs (No areas where detected where translation was the only language procedure affected); VGEN = Verb Generation. Cortical areas: SFG = Superior Frontal Gyrus, MFG = Middle Frontal Gyrus, IFG = Inferior Frontal Gyrus, PrG = Precentral Gyrus, PoG = Postcentr gyrus, SMG =Supramarginal gyrus, AnG = Angular Gyrus, STG = Superior temporal gyrus, MTG = Middle temporal gyrus, ITG = Inferior temporal gyrus. References in boldface indicate that language processing was frequently detected with the test or that the language process was specific to that area (e.g., Naming to Oral description was specific for STG and MTG). The sign (+) indicates that the error types were not clearly specified per brain area. a = area common to object naming, b = area specific to verbs, c = area specific to the periphery of the tumor, e = area specific to the posterior part of the gyrus, f = area specific to the inferior part of the gyrus, g = area specific to the superior part of the gyrus, h = area specific to Broca's area, i = area not frequent

Table A2.2

*Subcortical areas in the language-dominant hemisphere where language processing was detected by action naming and error types produced by patients during stimulation of that area*

	<u>ACTN</u>	<u>Error type</u>
<b>AF</b>	Bello et al. (2006,2007)	Anomia; Phonemic paraphasias
<b>IFOF</b>	Bello et al. (2007, 2008)	Semantic paraphasias
<b>ILF</b>	Bello et al. (2007)	Anomia; Reduction of spontaneous speech Semantic paraphasias.
<b>PV</b>	Bello et al. (2006)	
<b>SLF</b>	Bello et al. (2008)	Phonemic paraphasias
<b>SuF</b>	Bello et al. (2006, 2007)	Anomia; Reduction of spontaneous speech; Phonemic paraphasias
<b>UNC</b>	Bello et al. (2007, 2008)	Semantic paraphasia

*Notes.* ACTN = Action Naming; Subcortical areas: AF = Arcuate fasciculus, IFOF = Inferior fronto-occipital-fasciculus, ILF = Inferior longitudinal fasciculus, PV = Periventricular white matter, SLF = Superior longitudinal fasciculus, SuF = Subcallosal fasciculus and UNC = Uncinate fasciculus.

## **1. VISC subsets: description, number of items per condition, matching**

### **FAL\_Sub**

*Action-related verbs* describe actions performed with face/mouth, arm/hand and leg/foot. Hauk, Johnsrude and Pulvermüller (2004) argue that this type of actions (e.g., ‘to lick’, ‘to pick’ and ‘to kick’) activate the motor strip in a somatotopic manner (Leyton & Sherrington, 1917; Penfield & Boldrey, 1937). These verbs are similar to manipulation verbs, but “manipulation” refers to hand actions that require an object (e.g., ‘to type’ needs a keyboard) whereas “arm/hand action” does not entail object use (e.g., ‘to point’ is not a manipulation action as it does not require an object, even though it is an action, typically performed with the hand). *FAL\_Sub* includes 8 items per condition: face action (1), arm action (2), leg action (3). Items were matched for Picture Name Agreement (PNA<75), H-statistic (H-STAT), Visual complexity (Vcomplex), Age of Acquisition (AOA), Imageability (Imag), Frequency (Freq, Log10+1Freq, HighLowFq), Length in phonemes (LPh, LongShortPh), Instrumentality (Inst), Name-relatedness to a noun (NRN), Manipulation (MN) and Regularity (Reg\_Sub). Items could not be matched for Transitivity (TR) and Number of Internal arguments (NIA) as no transitive verbs could be added in face actions (1).

### **HighLowFqV\_Sub**

Effects of frequency were described in the VISC subsets. HighLowFqV\_Sub includes 17 items per condition: high frequency (1) and low frequency (0) based on the median of the sample (median=6.28). Conditions were matched for Picture Name Agreement (PNA<75), H-statistic (H-STAT), Visual complexity (Vcomplex), Age of Acquisition (AOA), Imageability (Imag), Instrumentality (Inst)m Action relatedness (FAL), Manipulation (MN).

No matching was possible for Transitivity (Tr), Name-relatedness to a noun (NRN) and Regularity (Reg\_Sub).

### **Inst\_Sub/NRN\_Sub**

The term *instrumental verbs* refers to actions requiring the use of an instrument that is not a body part. Instrumental verbs are divided in two categories: *name-related* (i.e., verbs that are phonologically identical to the name of the instrument, for example ‘to brush’) and verbs that are *not name-related* to the instrument (e.g., ‘to sew’). Instrumental verbs are easier than non-instrumental verbs for patients with aphasia (Jonkers & Bastiaanse, 2007), possibly because they activate two lemmas: one corresponding to the verb, one to the instrument. This coactivation may facilitate retrieval of the word form (the lexeme) but not necessarily of the lemma (Bastiaanse & Van Zonneveld, 2004). *Inst\_Sub* contains 24 items per condition: instrumental verb (1), non-instrumental verb (0). Items were matched for picture name agreement (PNA<75), H-statistic (H-STAT), Visual complexity (Vcomplex), Age of Acquisition (AOA), Imageability (Imag), Frequency (Freq, Log10+1, HighLowFq), Length in phonemes (LPh, LongShortPh), Manipulation (MN) and Regularity (Reg\_Sub). Items are not matched for Transitivity (Tr), Name-relatedness (NRN), Action-related verbs (FAL). *NRN\_Sub* contains 9 items per condition: verb that is related to a noun (1), verb that is not related to a noun (0). Items were matched for picture name agreement (PNA<75), H-statistic (H-STAT), Visual complexity (Vcomplex), Age of Acquisition (AOA), Imageability (Imag), Frequency (Freq, HighLowFq), Transitivity (Tr), Length in phonemes (LPh, LongShortPh), Action relatedness (FAL), Manipulation (MN) and Regularity (Reg\_Sub). Matching for Picture Name Agreement was low ( $p=0.070$ ). No matching was possible for Instrumentality ( $p=0.31$ ), as all noun-related items were also instrumental.

### **LongShortPhV\_Sub**

Effects of word length in naming were described in the ECCO subsets. *LongShortPhV\_Sub* includes 25 items per category: short verbs (0) and long verbs (1) based on the median of the infinitive in the sample (median=6). Items are matched for Picture Name Agreement (PNA<75), H-statistic (H-STAT), Visual complexity (Vcomplex), Age of Acquisition (AOA), Imageability (Imag), Frequency (Freq, Log10+1Freq, HighLowFq), Instrumentality (Inst), Name-relatedness to a noun (NRN), Action relatedness (FAL), Manipulation (MN) and Regularity (Reg\_Sub).

### **MN\_Sub**

The term *manipulation* refers to objects that are used with the hand (e.g., cellphone) and to actions that are performed by manipulating an object (e.g., to type). It is used to explain relations between body and mind. Some authors argue that words related to manipulation are represented (at least partially) in the motor strip, as suggested by studies using neuroimaging and motor evoked potentials (Willems & Hagoort, 2007, for review) as well as by lesion data (Saygin, Wilson, Dronkers, & Bates, 2004; Tranel et al., 2003). *MN\_Sub* includes 25 items per condition: non-manipulable verb (0), and manipulable verb (1). Items are matched for Picture Name Agreement (PNA<75), H-statistic (H-STAT), Visual complexity (Vcomplex), Age of Acquisition (AOA), Imageability (Imag), Frequency (Freq, Log10+1Freq, HighLowFq), Length in phonemes (LPh, LongShortPh), Name-relatedness to a noun (NRN), and Regularity (Reg\_Sub). Items were not matched for Transitivity (Tr), Instrumentality (Inst), and Action relatedness (FAL) as these variables correlate with Manipulation (MN). For example non-manipulable verbs (0) typically are non-instrumental, and vice-versa.



## **Reg\_Sub**

*Regular* verbs follow a typical conjugation paradigm and do not change their stem. *Irregular* verbs, by contrast, either do not follow the typical conjugation paradigm or change their stem form. English examples of irregular verbs are ‘sing’ and ‘take’, that change their root form into ‘sang’ and ‘took’ in the past, and ‘sung’ and ‘taken’ in the past participle. The debate on regular and irregular verbs focuses on whether or not they are processed by the same cognitive mechanisms and represented in the same neural network. Connectionist models assume no differences in the processing and representation of regular and irregular verbs (Sach, Seitz, & Indefrey, 2003), whereas dual-process models argue for different cognitive mechanisms and localization (Pinker & Prince 1988; Ullman et al., 1997). On the latter view, irregular verbs are stored as whole lexical forms and hence engage more heavily lexical mechanisms located in the temporal lobe and right cerebellum. Forms of regular verbs, on the other hand, are not stored as lexical entries, and their inflectional processes are triggered by morphosyntactic operations located in the left frontal lobe and basal ganglia (Howard et al., 1992; Ullman, 2004). According to dual-process models, a selective difficulty producing regular verbs is consistent with a disruption of grammatical processes, whereas deficits in retrieving irregular verb forms occur in the context of lexical damage. *Reg\_Sub* contains 16 items per condition. These refer to verbs that are regular in their whole conjugation (0) and verbs that have an irregular form in their conjugation (1). This form does not need to correspond to the third person singular. Items are matched for picture name agreement (PNA<75), H-statistic (H-STAT), Visual complexity (Vcomplex), Age of Acquisition (AOA), Imageability (Imag), Frequency (Freq, Log10+1Freq, HighLowFq), Transitivity (Tr), Length in phonemes (LPh, LongShortPh), Instrumentality (Inst), Action relatedness (FAL), Manipulation (MN) and Regularity (*Reg\_Sub*). All items are not name-related to a noun (NRN)

## **Tr\_Sub**

*Transitive* verbs are accompanied by a direct object (e.g., to cook, to kiss); *intransitive* verbs (e.g., to walk, to swim) cannot be accompanied by a direct object. *Intransitive* verbs are further divided in two categories: unergative verbs have one external argument that functions as an agent (e.g., to pray in ‘the nun prays’) and unaccusatives have one internal argument that functions as a theme and raises to subject position (e.g., to fall in ‘the kid<sub>i</sub> falls t<sub>i</sub>’). Therefore, a specific deficit for unaccusative verbs may denote difficulty with syntactic movement operations. People with aphasia typically fare worse with unaccusative and transitive than unergative verbs (e.g., Luzzatti et al., 2002); however, the reverse pattern is also on record (De Bleser & Kauschke, 2003). *Tr\_Sub* contains 23 items per condition: transitive verbs (1), intransitive verbs (0). Items are matched for picture name agreement (PNA<75), H-statistic (H-STAT), Visual complexity (Vcomplex), Age of Acquisition (AOA), Imageability (Imag), Frequency (Freq, Log10+1Freq, HighLowFq), Length in phonemes (LPh, LongShortPh), Instrumentality (Inst), Name-relatedness to a noun (NRN), Action relatedness (FAL), Manipulation (MN) and Regularity (Reg\_Sub).

## **Ac\_Balanced**

Differences between nouns/objects and verbs/actions were described in the ECCO subsets. *Ac\_Balanced* includes 20 items matched with a subset of 20 items extracted from the ECCO (*Ob\_Balanced*). The items are matched for picture name agreement (PNA<75, PNA+75), H-statistic (H-STAT), Age of Acquisition (AOA), Imageability (Imag), Frequency (Freq, Log10+1Freq, HighLowFq), biological/artifactual entities (Bio), Length in phonemes (LPh, LongShortPh), and semantic category (SemCat). Items could not be balanced for objective Visual Complexity (Forsythe et al. 2008) as indicated in the *Ob\_Balanced* section.

## **2. ECCO subsets: description, number of items per condition, matching**

### **Bio\_Sub**

The term *biological* entities refers to conspecifics, animals, fruits, and vegetables: whereas *non-biological* entities correspond to inanimate objects (musical instruments, tools, means of transportation, furniture, etc.). Within biological entities, a further distinction is drawn between those provided (animals) and those not provided (plants) with self-generated movement. Selective impairment and sparing of these categories have been reported, and typically signify semantic damage (see Capitani, Laiacona, Mahon, & Caramazza, 2003, for a review). *Bio\_Sub* includes 17 items per condition: biological (1) entities such as conspecifics, animals, fruits, vegetables and non-biological (0) or artifactual entities such as instruments, means of transportation, and furniture. Conditions are matched for Picture Name Agreement (PNA<75), H-statistic (H-STAT), Visual complexity (VcomplexGIF), Age of Acquisition (AOA), Imageability (Imag), Frequency (Freq, Log10+1Freq, HighLowFq), Length in Phonemes (LPh, LongshortPh), and Semantic category (SemCat).

### **HighLowFqN\_Sub**

*Written frequency* estimates can be obtained by counting the number of times that a word appears in a corpus (e.g., the CELEX database). Effects of frequency on response times and accuracy are reported in healthy populations (Oldfield & Wingfield, 1965) and in aphasia (Kemmerer & Tranel, 2000; Kittredge, Dell, Verkuilen, & Schwartz, 2008). Difficulty naming low-frequency items is generally attributed to deficits in lexical processing (Lesser & Milroy, 1993), but may be due also to semantic impairment or to a difficulty mapping semantic representations into lexical entries (Whitworth, Webster, & Howard, 2005). *HighLowFqN\_Sub* includes 18 items per condition: high (1) and low (0) frequencies calculated based on the median of the sample (median=6.28). Conditions are matched for

picture name agreement (PNA<75), H-statistic (H-STAT), Visual complexity (VcomplexGIF), Age of Acquisition (AOA), Imageability (Imag), Length in Phonemes (LPh, LongshortPh), Biological/artifactual entities (Bio), and Semantic category (SemCat).

### **LongShortPhN\_Sub/LPhN\_Sub**

Effects of *word length* in naming speed were found in healthy young individuals (Barry, Morrison, & Ellis, 1997), in elderly people (Hodgson & Ellis, 1998) and in aphasic speakers. Nickels and Howard (2004) disentangled ‘number of phonemes’ from ‘number of syllables’ and ‘syllabic complexity’, finding that number of phonemes (but not number of syllables) predicted word repetition accuracy in nine aphasic speakers showing effects of length. The typical length effects result in longer words being more difficult to process, usually as a consequence of post-lexical damage. The opposite pattern has been also occasionally reported (Lambon Ralph & Howard, 2000), and has been attributed to a difficulty accessing output phonology (Whitworth, Webster, & Howard, 2005) - longer words have fewer phonological neighbors and thus fewer competitors for activation. *LongShortPhN\_Sub* includes 21 items per condition: long words (1) and short words (0) based on the median of the sample (median = 6). *LPhN\_Sub* includes 10 items per condition: 0 = 4 phonemes; 1 = 5 phonemes; 2 = 6 phonemes; 3 = 7 phonemes. The conditions in both subsets are matched for Picture Name Agreement (PNA<75), H-statistic (H-STAT), Visual complexity (VcomplexGIF), Age of Acquisition (AOA), Imageability (Imag), Frequency (Freq, Log10+1Freq, HighLowFq), Biological/artifactual entities (Bio) and Semantic category (SemCat).

## **Ob\_Balanced**

Differences between nouns and verbs exist along semantic and morphosyntactic dimensions. Verbs typically refer to actions while nouns to objects. Verbs entail thematic role assignment, subject-verb agreement, complex argument structure, time reference, while nouns function as arguments of the predicate (e.g., Rofes & Miceli, 2014). Accuracy, reaction times and electrophysiological measures may be different between verbs and nouns, particularly when verbs are used in sentence context (Mätzig et al., 2009; Vigliocco et al., 2011, for reviews). *Ob\_Balanced* includes 20 items. These items are matched with a subset of 20 items extracted from the VISC (*Ac\_Balanced*). Items are matched for Picture Name Agreement (PNA<75, PNA+75), H-statistic (H-STAT), Age of Acquisition (AOA), Imageability (Imag), Frequency (Freq, Log10+1Freq, HighLowFq), Biological/artifactual entities (Bio), Length in phonemes (LPh, LongShortPh), and Semantic category (SemCat). Items could not be balanced for objective Visual Complexity (Forsythe et al. 2008), as the lossless GIF compression measures we proposed for both tests were not comparable, as the drawings had been made by different authors, probably using different softwares and different compression measures.

## **3. VISC: materials and procedure, imageability and age-of-acquisition ratings**

### *Materials and procedure*

Stimuli consist of 85 black-and-white drawings, similar to those of other tests (Bastiaanse, Edwards, Maas, & Rispens, 2003; Hammelrath, 2000; Metz-Lutz et al., 1991; Snodgrass & Vanderwart, 1980). Stimuli were presented by using Psychtoolbox for MATLAB. A beep was presented 500ms prior to stimulus presentation, and each stimulus was shown for 4000ms. The Psychsound commands of Psychtoolbox were used to record the responses produced by each subject to each item. Responses that corresponded to the expected target or

were rated as plausible given the ratings of healthy participants in the first study, and that were produced within the 4000ms time frame, were rated as correct.

#### *Imageability and age-of-acquisition ratings*

Imageability and age-of-acquisition ratings were collected through an on-line questionnaire to obtain ratings for all test items. We assessed the AoA of the target objects and that of the synonyms produced by healthy participants in the first step of the picture-name agreement study. Participants were asked to estimate on a 5-point Likert scale at which age they thought they had learned the word in either its spoken or written form (0-3 years, 4-6 years, 7-10 years, 10-12 years, older than 12 years). The procedure is similar to Carroll and White (1973). To assess Imageability, we asked participants to indicate on a 5-point Likert scale how difficult it was to imagine the object (very difficult, difficult, average, easy, very easy). A series of low-imageability action verbs was included in the list of to-be-rated words (e.g., categorize, deviate, define, tolerate). To the best of our knowledge, there is no exhaustive list of imageability of Italian verbs. Therefore, 65 low-imageability verbs from the list in Bird, Franklin and Howard (2001) were translated into Italian. Below we provided the instructions for each questionnaire, in English and Italian.

### **3. ECCO: materials and procedure, imageability and age-of-acquisition ratings**

#### *Materials and procedure*

Sixty-two object drawings from the Snodgrass and Vanderwart set (1980) were used. Stimuli were presented individually on a computer screen, and the starter word *Ecco...* (This is...) was shown above each stimulus. Participants were asked to produce the name of the object preceded by its determiner (e.g., *Ecco la pera*, 'Here [is] the pear').

### *Imageability and age-of-acquisition ratings*

The same procedure as for the VISC was followed. In addition to the 57 target items, all corresponding to concrete objects, the imageability questionnaire included 65 low-imageability nouns (e.g., philosophy, consciousness, reputation, merit), that had been rated between 2 and 3 on a 7-point scale in a study by Della Rosa, Catricalà, Vigliocco, and Cappa (2010). In order to prevent list biases, the names of low-imageability items were pseudo-randomly interspersed with those of the high-imageability, target objects. This ensured that stimuli in this task covered a wide imageability range, and prevented participants from inappropriately rating test items (all concrete objects) as being of low imageability. We followed the procedure of Paivio, Yuille and Madigan (1968).

## 5. Instructions for imageability and age of acquisition in English and in Italian

### *Age-of-acquisition*

For an experiment regarding some properties of the memory of words, we need your opinion on the age-of-acquisition of some words. We are interested to know when you learned for the first time the meaning of a series of words, in the spoken or in the written form. In the following questionnaire, you will be asked to read some words. You are asked to answer on a scale from 1 to 5 if you estimate to have learned the word, in its spoken form or written form, when you were '0 to 5 years', '4 to 6 years', '7 to 9 years', '10 to 12 years' or when you were 'older than 12'.

For example you will read the word 'pasta' (specific to ECCO)/ 'to snow' (specific to VISC), and will estimate when you learned that word. If you think you learned it when you were around 0 to 3 years, please tick the first box of the scale.

ITALIAN. *Per un esperimento che riguarda alcune proprietà della memoria di parole, abbiamo bisogno del tuo giudizio sull'età di acquisizione di alcune parole. Ci interessa sapere quando, nel corso della tua vita, hai imparato per la prima volta il significato di una serie di parole, in forma parlata o scritta. Nel seguente questionario ti viene chiesto di leggere alcune parole. Dovrai rispondere su una scala da 1 a 5 se ritieni di aver appreso la parola, verbalmente o in forma scritta, quando avevi dagli '0 ai 3 anni', dai '4 ai 6 anni', dai '7 ai 9 anni', tra i '10 e i 12 anni' o 'dopo i 12 anni'.*

*Per esempio vedrai la parola 'pasta' (specific to ECCO)/ 'nevicare' (specific to VISC). Dovrai stimare quando l'hai imparata per la prima volta. Se pensi di averla imparata nella fascia d'età compresa tra 0 e 3 anni, dovrai segnare la prima casella della scala.*



## Imageability

Words differ in their capacity to be imageable. Some words can be imagined very easily, others need more effort. The goal of this experiment is to estimate the difficulty to construct a visual image for each noun (specific to ECCO)/ verb (specific to VISC) on a list. You will answer on a scale from 1 to 5, indicating if constructing a mental image for each word is 'very easy', 'easy', 'average', 'difficult' or 'very difficult'.

(Specific to ECCO). For example, think about the nouns 'house' and 'love' (in Italian, the latter is indicated by an unambiguous noun). Probably it is easy to construct a mental image of 'house': you could think of the image of a building divided in different floors and rooms. Instead, constructing the mental image for 'love' is probably more difficult. Think of the feeling and of the instinct that unites two people.

(Specific to VISC). For example, think about the verbs 'to eat' and 'to know'. Probably, 'to eat' is easy to imagine: you could think of a man eating a piece of cake. Instead, constructing the mental image for 'to know' is probably more difficult. Think about what do you do when you ask yourself 'Do I know that?'. You could touch your head and look surprised. Therefore, you could think of the image of a woman who touches her head in surprise.

ITALIAN. *Le parole differiscono per la loro immaginabilità. Alcune parole possono essere immaginate facilmente, altre richiedono un maggior impegno. Lo scopo di questo esperimento è di stimare la difficoltà di costruire un'immagine mentale per ogni nome (specific to ECCO) / verbo (specific to VISC) in una lista. Dovrai rispondere su una scala con valori da 1 a 5; dovrai stabilire se per ogni parola è 'molto facile', 'facile', 'medio', 'difficile', 'molto difficile' costruire un'immagine mentale.*

*(Specific to ECCO) Per esempio, pensa ai nomi 'casa' e 'amore'. Probabilmente è facile costruire un'immagine mentale per 'casa': basta pensare all'immagine di un edificio suddiviso in piani e vani, adibito ad abitazione. Invece per 'amore' è probabilmente più difficile. Pensa al sentimento ed all'istinto che lega due persone. (Specific to VISC) Per esempio, pensa ai verbi 'mangiare' e 'conoscere'. 'Mangiare' probabilmente è facile da immaginare: basta pensare all'immagine di un uomo che sta mangiando un pezzo di torta. Invece 'conoscere' probabilmente è più difficile da immaginare. Pensa a cosa fai quando ti chiedi 'questo lo conosco?'. Potresti toccarti la testa e sembrare sorpreso. Quindi potresti pensare all'immagine di una donna sorpresa che si tocca la testa.*

## 6. MATLAB script to calculate the H-STATISTIC

```
function H = StatisticH(p)

% STATISTICH
% It computes the statistic H for a given set of proportions.
% It is based on the formula provided by Snodgrass & Vanderwart (1980)
%
% SCORING INSTRUCTIONS
% a) Eliminate: do not know responses
% b) Include with correctly name word: mispronunciations, diminutives
% c) Include with unabbreviated forms: uncommon abbreviations (b carriage
for
% baby carriage)
% d) Count as separate words: not target word (vehicle for airplane, potato
% for carrot); commonly accepted abbreviations (TV for television),
% elaborations (bell pepper for pepper)
% e) Count the first response only: when two correct names are used
% f) Count the correct response: when two names are used within the given
% time
%
% EXAMPLES OF USAGE
% I) For the picture "plane", 30 people said "plane" 30 times. The result
is
% 0 which indicates perfect agreement.
% StatisticH ([30/30])
%
% II) For the picture "plane", 15 people said "plane", and 15 people said
% "airplane".
% The result is 1.
% StatisticH ([15/30, 15/30])
%
% III) For the picture "plane", 10 people said "plane", 10 people
"airplane",
% and 10 people said "I do not know"
% The result is 1
% StatisticH ([10/20, 10/20])
%
% IV) For the picture "plane", 10 people said "plane", 10 people "airplane",
5
% people "I do not know", and 5 people "vehicle".
% The result is 1.5219
% StatisticH ([10/25, 10/25, 5/25])
%
% ----- Error handling -----
if(sum(p < 0) > 0) % We don't want negative p's
    error('Input error: at least one p coefficient is negative');
end
if(sum(p > 1) > 0) % We don't want p's greater than 1
    error('Input error: at least one p coefficient is greater than one');
end
pSum = sum(p);
if(pSum ~= 1) % The sum is expected to be 1
    disp('Warning: the p coefficients are not summing up to 1');
end
% ----- Calculation -----
elementsToBeSummed = p.*log2(1./p);
%
% When some p is zero, the function will throw a NaN ...
positionsOfZeros = find(p == 0); % ... so we find them ...
```

```
elementsToBeSummed(positionsOfZeros) = 0; % ... and use the limit when p
tends to zero instead of direct evaluation
%
H = sum(elementsToBeSummed);
%
End
```

APPENDIX C: Matching for linguistic variables, post-hoc tests, and individual results  
(Chapter 4)

Table C4.1

*Matching of the tests for relevant linguistic variables including imageability*

	<u>FinVerb</u>	<u>ActNam</u>	<u>Vgen</u>	<u>ObNam</u>	<u>NVgen</u>	<u>Significance</u>
Frequency	32.32(8.5)	44.01(19.2)	42.9(12)	34.73(13.26)	45(14.4)	$\chi^2(4,100)=1.203$ , p=1.616
Imageability	1.19(0.02)	1.2(0.02)	1.18(0.02)	1.14(0.01)		$\chi^2(3,80)=5.358$ , p= 0.147
Age of Acquisition	1.91(0.11)	1.84(0.09)	1.78(0.12)	1.98(0.12)		$\chi^2(3,80)=1.883$ , p= 0.597
Length in Phonemes	5.65(0.25)	6.3(0.3)	5.75(0.3)	6.2(0.4)	6.3(0.48)	$\chi^2(4,100)=$ 1.616, p=0.806
Manipulable	14	12	11	9	10	
Non-Manipulable	6	8	9	11	10	
Transitive	12	14	13			
Intransitive	8	6	7			
Internal Arguments	12	14	13			
Instrumental	11	11	8			
Non-Instrumental	9	9	12			
Irregular	7	3	4			
Regular	13	17	16			
Name Related	4	3	4			
Non-Name Related	16	17	16			
Face	3	1	0			
Arm	12	13	12			
Leg	3	3	3			
Face/Arm	1	2	3			
Arm/Leg	1	0	0			
Non-Face/Arm/Leg	0	1	2			
Biological				8	4	
Non-Biological				12	16	

*Notes.* ActNam: Action Naming; ObNam: Object Naming; FinVerb: Producing a finite verb in sentence context; Vgen: Verb generation; NVgen: Name of the picture for which the verb is produced in the Verb Generation task (e.g., ball > to play). Standard deviation is reported in parentheses. N=20 items per task.

Table C4.2

*Individual behavioral data (percentage correct).*

	<u>CADL-2</u>	<u>CETI</u>	<u>CAL</u>	<u>ObNam</u>	<u>Vgen</u>	<u>FinVerb</u>	<u>ActNam</u>
AL	100	93.8	96.7	95	80	80	100
BL	83	73.8	81.1	90	75	80	75
BS	59	75	83.3	0	25	0	5
CB	93	78.8	73.3	90	55	60	0
CC	82	92.5	91.1	85	90	60	55
CK	56	42.5	53.3	35	5	5	25
ES	90	68.8	80	55	55	65	40
FT	95	68.8	41.1	35	45	70	60
GC	93	83.8	78.9	5	25	40	40
GM	53	65	52	0	0	0	0
LF	89	57.6	65.6	10	0	35	10
OS	84	81.3	92.2	85	80	80	65
OT	88	82.5	83.3	50	30	75	10
PG	70	51.3	41.1	0	0	0	5
PI	90	73.8	67.8	85	80	90	75
PS	49	70	72	65	15	20	10
RA	97	82.5	83.3	90	90	90	70
RG	80	53.8	52.2	0.0	15.0	25.0	30.0
RL	89	78.8	77.8	30	45	50	25
TG	67	57.5	55.6	65	30	30	5
UA	87	83.8	62.2	80	15	60	40
m(sd)	81 (15)	72 (14)	71 (16)	50 (36)	41 (32)	48 (31)	37 (30)

*Notes.* ActNam: Action Naming; ObNam: Object Naming; FinVerb: Producing a finite verb in sentence context; Vgen: Verb generation. m(sd): mean and standard deviation.

Table C4.3

*Post-hoc tests of picture-naming tasks (Z and FDR corrected p-values).*

	<u>VGen-</u> <u>ObNam</u>	<u>FinVerb-</u> <u>ObNam</u>	<u>ActNam-</u> <u>ObNam</u>	<u>FinVerb-</u> <u>VGen</u>	<u>ActNam-</u> <u>VGen</u>	<u>ActNam-</u> <u>FinVerb</u>
	Z= -1.542 p=0.1845	Z=-0.308 p=0.7577	Z=-2.015 p=0.0878	Z=-2.017 p=0.0878	Z=1.073 p=0.3399	Z=-2.546 p=0.0653

*Notes.* ActNam: Action Naming; ObNam: Object Naming; FinVerb: Producing a finite verb in sentence context; Vgen: Verb generation.

Table C4.4

*Spearman correlations of picture-naming tasks and functional-measures (rho and p-value).*

	<u>CADL-2</u>	<u>CETI</u>	<u>CAL</u>	<u>ObNam</u>	<u>VGen</u>	<u>FinVerb</u>	<u>ActNam</u>
CADL-2	1	0.511*	0.324	0.441*	0.561*	0.727*	0.538*
	p=0	p=0.0215	p=0.1516	p=0.0492	p=0.0129	p=0	p=0.0147
CETI	0.511*	1	0.763*	0.541*	0.611*	0.564*	0.452*
	p=0.021	p=0	p=0	p=0.0147	p=0.006	p=0.0129	p=0.0454
CAL	0.324	0.763*	1	0.554*	0.701*	0.542*	0.437*
	p=0.1516	p=0.	p=0	p=0.0136	p=0	p=0.0148	p=0.0494
ObNam	0.441*	0.541*	0.544*	1	0.779*	0.776*	0.577*
	p=0.0494	p=0.0148	p=0.0136	p=0	p=0	p=0	p=0.0113
VGen	0.561*	0.611*	0.701*	0.780*	1	0.830*	0.683*
	p=0.0129	p=0.0064	p=0.0009	p=0	p=0	p=0	p=0.0014
FinVerb	0.727*	0.554*	0.542*	0.776*	0.830*	1	0.793*
	p=0	0.0129	0.0148	p=0	p=0	p=0	p=0
ActNam	0.538*	0.462*	0.437*	0.578*	0.683*	0.792*	1
	p=0.0148	p=0.0454	p=0.0495	p=0.0113	p=0.0013	p=0	p=0

*Notes.* ActNam: Action Naming; ObNam: Object Naming; FinVerb: Producing a finite verb in sentence context; Vgen: Verb generation. We calculated false discovery rate (FDR) adjusted p-values to correct for multiple comparisons. \*Significant at  $p \leq 0.05$

Table C4.5

*Comparisons of overlapping correlations (Steiger's Z)*

	<u>ObNam-Vgen</u>	<u>ObNam-FinVerb</u>	<u>ObNam-ActNam</u>
CADL-2	Z=-0.92509 p=0.2662	Z=-2.34918 p=0.0282*	Z=-0.53177 p=0.2975
CETI	Z=-1.02434 p=0.4290	Z=-0.17940 p=0.4290	Z=0.49075 p=0.4290
CAL	Z=-1.25967 p=0.3120	Z=0.09320 p=0.4629	Z=0.64510 p=0.3891

*Notes.* ActNam: Action Naming; ObNam: Object Naming; FinVerb: Producing a finite verb in sentence context; Vgen: Verb generation. We calculated false discovery rate (FDR)-adjusted p-values to correct for multiple comparisons. \*Significant at  $p \leq 0.05$

Table C4.6

*Paired tests on picture-naming tasks (Fisher Exact – two tailed)*

	<u>ObNam vs</u> <u>Vgen</u>	<u>ObNam vs</u> <u>FinVerb</u>	<u>ObNam vs</u> <u>ActNam</u>	<u>Vgen vs</u> <u>FinVerb</u>	<u>Vgen vs</u> <u>ActNam</u>	<u>FinVerb vs</u> <u>ActNam</u>
BL	p=0.168	p=1	p=1	p=0.168	p=0.364	p=1
CB	p=0.004*	p=0.078	p=<0.001*	p=1.000	p=<0.001*	p=<0.001*
CC	p=1	p=0.232	p=0.164	p=0.164	p=0.164	p=1
CK	p=0.003*	p=0.003*	p=0.8772	p=1	p=0.273	p=0.273
GC	p=0.04*	p=0.04*	p=0.04*	p=0.601	p=0.601	p=1
LF	p=0.731	p=0.254	p=1	p=0.04*	p=1	p=0.254
OT	p=0.333	p=0.282	p=0.028*	p=0.028*	p=0.282	p=<0.001*
PS	p=0.009*	p=0.02*	p=<0.001*	p=1	p=1	p=0.992
TG	p=0.084	p=0.084	p=<0.001*	p=1.000	p=0.109	p=0.084
UA	p=<0.001*	p=0.045*	p=0.044*	p=0.002*	p=0.186	p=0.343

*Notes.* ActNam: Action Naming; ObNam: Object Naming; FinVerb: Producing a finite verb in sentence context; Vgen: Verb generation. = Fisher exact (two tailed). We calculated false discovery rate (FDR) adjusted p-values to correct for multiple comparisons. \*Significant at  $p \leq 0.05$

Table C4.7

*Major error types per task and participant*

	<u>AL</u>	<u>BL</u>	<u>BS</u>	<u>CB</u>	<u>CC</u>	<u>CK</u>	<u>ES</u>	<u>FT</u>	<u>GC</u>	<u>GM</u>	<u>LF</u>	<u>OS</u>	<u>OT</u>	<u>PG</u>	<u>PI</u>	<u>PS</u>	<u>RA</u>	<u>RG</u>	<u>RL</u>	<u>TG</u>	<u>UA</u>	<u>TOTAL</u>
<b>ActNam</b>																						
AN	0	1	16	8	2	7	6	5	4	13	9	0	1	2	2	9	0	6	8	10	9	118
PHON	0	0	0	0	0	1	0	0	0	4	0	2	0	1	2	0	3	1	1	0	1	16
SEM	0	3	3	2	2	1	0	0	0	1	1	1	2	2	1	0	2	1	1	1	0	24
LATE	0	0	0	0	2	2	1	0	1	0	2	3	0	0	1	0	0	0	4	0	0	16
CAT/S	0	0	0	1	0	2	0	0	4	2	1	0	1	5	0	6	0	0	2	3	0	27
MSYNT	0	0	0	9	0	0	0	0	0	0	4	0	17	4	1	3	0	0	0	3	0	41
<b>ObNam</b>																						
AN	0	0	17	1	1	1	4	7	3	6	14	1	2	1	1	1	1	6	3	4	3	77
PHON	1	0	0	0	0	0	0	2	1	3	1	0	1	6	0	1	1	0	0	1	0	18
SEM	1	1	3	1	1	3	4	0	2	1	1	0	3	0	1	5	0	0	1	1	1	30
LATE	0	1	0	0	0	1	0	0	5	2	0	2	0	0	1	0	0	0	9	0	0	21
<b>FinVerb</b>																						
AN	1	0	18	4	1	10	1	5	4	6	7	2	0	1	1	5	1	6	4	5	8	90
PHON	0	0	0	0	2	1	0	1	1	1	2	1	0	6	0	0	0	0	0	2	0	17
SEM	1	2	1	2	2	5	2	0	0	3	1	0	4	0	0	1	0	0	1	1	0	26
LATE	0	1	0	0	2	0	0	0	2	0	3	1	0	0	0	0	0	0	2	0	0	11
CAT/S	0	0	0	0	0	1	0	0	0	1	0	0	0	2	0	6	1	0	0	3	0	14
MSYNT	0	0	0	2	0	0	0	0	0	1	0	0	0	0	0	2	0	0	0	0	0	5
<b>Vgen</b>																						
AN	0	0	13	6	2	9	4	3	7	12	11	1	2	1	1	11	0	2	7	3	11	106
PHON	0	0	0	0	0	1	0	0	1	6	0	0	0	0	0	0	1	1	0	1	0	11
SEM	2	3	2	0	0	0	2	1	1	0	0	0	1	0	2	1	1	3	0	2	0	21
LATE	0	1	0	0	1	0	1	0	3	0	0	2	0	0	1	0	0	0	1	1	0	11
CAT/S	0	0	0	2	0	8	1	0	0	7	8	0	8	4	0	5	0	1	0	8	5	57

*Notes.* AN: Anomia; PHON: Phonemic paraphasia; SEM: Semantic paraphasia; LATE: Answer later than 4 seconds; Cat/S: Category substitution (e.g., “chair” instead of “sitting”); MSYNT = Morpho-syntactic errors (i.e., errors of name agreement between the pronoun and the noun, e.g. “She paint” instead of “She paints”; also omission of the infinitival suffix “-re”); ActNam: Action Naming; ObNam: Object Naming; FinVerb: Finite verbs; Vgen: Verb generation.



## EPILOGUE

### ABOUT THE AUTHOR

I am a PhD candidate of the IDEALAB program (International Doctorate for Experimental Approaches to Language and Brain) at the University of Trento, the University of Groningen, and Macquarie University. I work on the neural correlates of noun and verb production in aphasia and their application to assessment of language in people with brain tumors and post-stroke aphasia. I also study non-fluent aphasia in Ibero-Romance.

During the PhD, I published 6 peer-reviewed articles and 7 abstracts, gave 17 invited talks and presented 4 posters, co-advised a master's student, and helped out in the organization of 4 events/conferences. I also took advantage of the many courses offered at my home and host universities, mastered my knowledge of Italian and Portuguese (which add up to Catalan, Spanish, and English), and I took courses at other centers (e.g., a tractography course in Padova, a neuroanatomy course in Sydney, and a virtual and classical brain dissection course in Paris). As a master's student, I graduated from Joint European Erasmus Mundus Master's Program in Clinical Linguistics (MSc) held at the University of Groningen and Potsdam University. I visited Neurosurgery Department in Montpellier to be trained in functional brain mapping, oncological neurosurgery and brain dissection. Before graduating in Catalan and English philology (double BA) at the Autonomous University of Barcelona, I was given a grant from the Center for Theoretical Linguistics to work on Comparative Linguistics. I also spent one year at the University of California in Santa Cruz, where I took courses in Syntax, Pragmatics, and Scientific Writing.

Currently, I am a member of the European Low Grade Glioma Network, and the Collaboration of Aphasia trialists. I was also very happy to be offered the opportunity to review for Aphasiology. My research interests embrace clinical linguistics (acquired disorders, assessment, awake surgery, post-stroke aphasia, rehabilitation, test standardization), brain mapping (direct electrical stimulation, functional magnetic resonance imaging, and tractography) and language processing (lexico-semantics, morpho-syntax, nouns and verbs, spontaneous speech, and time reference).

After the PhD thesis, I would be glad to continue on the academic path, become an independent researcher and a university professor, and contribute notably to the advance of functional approaches to language and cognition in awake surgery, post-stroke aphasia, language learning and treatment, and related subfields. I recently moved to Dublin (Ireland) together with my wife, and so far, I like it a lot here.

## PUBLICATIONS, PUBLISHED ABSTRACTS, AND AWARDS

### Publications

Rofes, A., Spena, G., Miozzo, A., Fontanella, M., & Miceli, G. (in press). Advantages and disadvantages of intraoperative language tasks in awake surgery: A three-task approach for prefrontal tumors. *Journal of Neurosurgical Sciences*.

Rofes, A., Capasso, R., & Miceli, G. (2015). Verb production tasks in the measurement of communicative abilities in aphasia. *Journal of Clinical and Experimental Neuropsychology*, 37(5), 483-502. doi:10.1080/13803395.2015.1025709

Rofes, A., de Aguiar, V., & Miceli, G. (2015). A minimal standardization setting for language mapping tests: An Italian example. *Neurological Sciences*, 36(7), 1113-9. doi:10.1007/s10072-015-2192-3

Martínez-Ferreiro, S., de Aguiar, V., & Rofes, A. (2015). Non-fluent aphasia in Ibero-Romance: a review of morphosyntactic deficits, *Aphasiology*, 29(1), 101-126. doi:10.1080/02687038.2014.958915

Rofes, A., & Miceli, G. (2014). Language mapping with verbs and sentences in awake surgery: A Review, *Neuropsychology Review*, 24(2), 185-99. doi:10.1007/s11065-014-9258-5

Rofes, A., Bastiaanse, R., & Martínez-Ferreiro, S. (2014). Conditional and future tense impairment in non-fluent aphasia, *Aphasiology*, 28(1), 99-115. doi:10.1080/02687038.2013.850650

### Published abstracts

Rofes, A., Santini, B., Ricciardi, G.K., Talacchi, A., Pinna, G., Nickels, L., & Miceli, G. (2015). Nouns and verbs: Perioperative comparisons in brain tumors. *Stem- Spraak- en Taalpathologie*, 20(S01), 14-16. doi:32.8310/supplement/1914

Rofes, A., Talacchi, A., Santini, B., Cappelletti, G., & Miceli, G. (2015). Intraoperative language mapping: A two-task approach. *Stem- Spraak- en Taalpathologie*, 20(S01), 191-194. doi:32.8310/supplement/1914

Rofes, A. & Miceli, G. (2014). Naming finite verbs predicts language abilities in daily living. *Stem- Spraak- en Taalpathologie*, 19(S04), 134-136. doi:32.8310/supplement/1914

Rofes, A., Santini, B., Mirtuono, P., Foroni, R., Pinna, G., Nickels, L., & Miceli, G. (2014). Lexico-semantic dissociations in patients with brain tumors in the posterior temporal lobe. *The Society for the Neurobiology of Language*, 194.

Rofes, A., De Witte, E., Mariën, P., & Bastiaanse, R. (2013). Object naming may overestimate patients' language performance after neuro-oncological surgery: A case study. *Stem- Spraak- en*

*Taalpathologie*, 18(S01), 59-63. doi:32.8310/S01/1813-59

Scientific communication/disclosure

Rofes, A. (2015). La neurocirugía con el paciente despierto [Spanish: Awake neurosurgery] Cuaderno de Cultura Científica (KZK), Cátedra de Cultura Científica, University of the Basque Country, Aug. 3 [Invited article, press here]

Posters, talks and invited presentations

Rofes, A., Santini, B., Ricciardi, G.K., Talacchi, A., Pinna, G., Nickels, L., & Miceli, G. (2015). Nouns and verbs: Perioperative comparisons in brain tumors. XV Science of Aphasia Conference, Sep. 17-22, Aveiro, Portugal [Talk]

Rofes, A., Talacchi, A., Santini, B., Cappelletti, G., & Miceli, G. (2015). Intraoperative language mapping: A two-task approach. XV Science of Aphasia Conference, Sep. 17-22, Aveiro, Portugal [Talk]

Rofes, A., & Miceli, G. (2015). When object naming may not be enough: Two single cases. European Low Grade Glioma Network Meeting. Jun. 25-27, Paris, France [Invited presentation]

Dallabona, M., & Rofes, A. (2015). Neuropsychological assessment: pre-, intra-, and postoperative. Connect Brain. Apr. 23-26, Trento, Italy [Invited presentation]

Rofes, A. (2015). Awake surgery: O papel do Terapeuta da Fala [Portuguese: Awake surgery: the role of the Speech and Language Therapist] University of Algarve, Mar. 26, Faro, Portugal [Guest Lecture at Bachelor's in Speech and Language Therapy, Prof. Susana Rodrigues and Prof. Ana Catarina Baptista]

Rofes, A. (2015). Nouns and verbs in awake surgery: needs and answers. University College London, Mar. 5, London, UK [Invited presentation at Prof. Vigliocco's lab]

Rofes, A. (2015). Awake surgery: theory and praxis. University of Groningen, Jan. 12, Groningen, The Netherlands [Guest lecture at European Master's in Clinical Linguistics]

Rofes, A. & Miceli, G. (2014). La valutazione comprensiva del linguaggio nella chirurgia da sveglio [Italian: Comprehensive language assessment in awake surgery] Habilita: La rieducazione neurocognitiva nelle malattie degenerative del sistema nervoso centrale. Nov. 22, Bergamo, Italy [Invited presentation]

Di Giacopo, R., Calanni, M., Rofes, A., Fortis, P., Malaguti, M., Ottaviani, D., & Miceli, G. (2014). La riabilitazione delle funzioni esecutive e viso-spaziali nella malattia di Parkinson: la nostra esperienza [Italian: Executive and visuo-spatial functions rehabilitation in Parkinson's Disease: our experience] III Congresso nazionale congiunto LIMPE/DISMOV-SIN, Nov. 14, Salerno, Italy [Poster]

- Rofes, A., Santini, B., Mirtuono, P., Foroni, R., Pinna, G., Nickels, L., & Miceli, G. (2014). Lexico-semantic dissociations in patients with brain tumors in the posterior temporal lobe. CIMEC Doctoral School Day, Sep. 26, Rovereto, Italy [Talk]
- Rofes, A., Capasso, R., & Miceli, G. (2014). Naming finite verbs predicts language abilities in daily living. XV Science of Aphasia Conference, Sep. 19-24, Venice, Italy [Talk]
- Rofes, A., Santini, B., Mirtuono, P., Foroni, R., Pinna, G., Nickels, L., & Miceli, G. (2014). Lexico-semantic dissociations in patients with brain tumors in the posterior temporal lobe. The Society for the Neurobiology of Language Conference, Aug. 27-29, Amsterdam, The Netherlands [Poster]
- Rofes, A., Talacchi, A., Santini, B., Zoccatelli, G., Alessandrini, F., Nickels, L., & Miceli, G. (2014). The language-role of the inferior fronto-occipital fasciculus. Center for Neurocognitive Rehabilitation, University of Trento. May 21, Rovereto, Italy [Invited presentation]
- Rofes, A., Talacchi, A., Santini, B., Zoccatelli, G., Alessandrini, F., Nickels, L., & Miceli, G. (2014). The language-role of the inferior fronto-occipital fasciculus: Evidence from awake surgery. Rovereto Workshop on Concepts, Actions, and Objects: Functional and neural perspectives. May 8-11, Rovereto, Italy [Poster]
- Martínez-Ferreiro, S. Reyes Gómez, A.F., & Rofes, A. (2014). Métodos neurocientíficos [Spanish: Neuroscientific methods]. Workshop: Cerebro y Lenguaje, El Bosque University. Apr. 9, Bogotá, Colombia [Invited presentation]
- Rofes, A. (2014). Language research in brain tumor surgery. Neuroanatomy course HLTH214, Macquarie University. Jan. 10, Sydney, Australia [Invited presentation]
- Rofes, A. (2013). Language in brain tumor patients and language testing in awake surgery. ARC Centre of Excellence in Cognition and its Disorders (CCD). Aphasia group, Macquarie University. Oct. 10, Sydney, Australia [Invited presentation]
- Rofes, A., & Miceli, G. (2013). Language mapping in awake surgery: A critical review of tests with verbs and sentences. CCD Annual Workshop, Nov. 27-28, Sydney, Australia [Poster]
- Rofes, A., De Witte, E., Mariën, P., & Bastiaanse, R. (2013). Object naming may overestimate patients' language performance after neuro-oncological surgery: A case study. XIV Science of Aphasia Conference, Sep. 20-25, Brussels, Belgium [Talk]
- Rofes, A. (2013). Functional surgical neuro-oncology: relevant language tests, better outcomes. Faculty of Medicine and Surgery, University of Verona. Apr. 17, Verona, Italy [Invited presentation]
- Arsenijević, B., Martínez-Ferreiro, S., & Rofes, A. (2011). Determiners in Catalan agrammatism: A case study. XII Science of Aphasia Conference, Sep. 01-05, Barcelona, Spain [Poster]

### Grants and awards

2012-2015 Erasmus Mundus Joint Doctorates (EMJDs) grant, IDEALAB PhD program (International Doctorate in Experimental Approaches to Language And Brain) - em-idealab.com (127.200€)

2013-2014 Cognitive Science Postgraduate Research Grant (CSPRG) R4. Macquarie University (\$600)

2010-2012 Erasmus Mundus Master Course (EMMCs) grant, EMCL program (European Master in Clinical Linguistics) - emcl-mundus.com (12.000€)

2011 Premi Extraordinari de Titulació – Special Award BA in Catalan philology and BA in English philology. Autonomous University of Barcelona (fee exemption)  
2010 Collaboration grant (COLAB). Department of Catalan philology and Center for Theoretical Linguistics at the Autonomous University of Barcelona.

2009 Grant and Grant with Honors. Becas de Movilidad Internacional Fernando Alonso 2008-

2009. Banco Santander (3.000€ + 600€)

2009 Bancaja Grants to best academic records of the Autonomous University of Barcelona International Mobility's program (1.500€)

2009 International Mobility program grants of the Autonomous University of Barcelona International Mobility's program (1.500€)

2009 General grant to undergraduates (GRAL). Agaur/Ministerio (fee exemption)

2007-2008 Four grants. Summer courses at University of Girona, symposium in Mallorca (DRAC, 250€), and general grants (GRAL) for undergraduates (fee exemption)